

Document made available under the Patent Cooperation Treaty (PCT)

International application number: PCT/BE04/000172

International filing date: 02 December 2004 (02.12.2004)

Document type: Certified copy of priority document

Document details: Country/Office: CA
Number: 2,487,529
Filing date: 15 November 2004 (15.11.2004)

Date of receipt at the International Bureau: 11 July 2005 (11.07.2005)

Remark: Priority document submitted or transmitted to the International Bureau but not in compliance with Rule 17.1(a) or (b)



World Intellectual Property Organization (WIPO) - Geneva, Switzerland
Organisation Mondiale de la Propriété Intellectuelle (OMPI) - Genève, Suisse

Federale Overheidsdienst Economie,
KMO, Middenstand en Energie
Dienst voor de Industriële Eigendom
Koning Albert II-laan 16
B-1000 Brussel

By Special Mail


Sint-Martens-Latem, June 15, 2005

Our ref. : BUNT-001-PCT
Re. : PCT Patent Application n° PCT/BE2004/000172
Filed on December 2, 2004
In the name of B&B Beheer N.V.
Titel: "Use of D4 and 5-HT2A antagonists, inverse agonists or partial agonists"


Dear Mrs. De Wulf,

- 9
- Further to our letter of March 29, 2003, we herewith submit the missing certified copy of priority document CA 2,487,529 filed on November 15, 2004 under Rule 17.1 (a) PCT in respect of the above-mentioned International patent application.

Respectfully submitted,



Mrs. De Clercq, Ann
Professional Representative before the EPO



Encl.: Certified copy (1)
Confirmation copy (please return)



Office de la propriété
intellectuelle
du Canada

Un organisme
d'Industrie Canada

Canadian
Intellectual Property
Office

An Agency of
Industry Canada

7CT/BE 2004/000172

*Bureau canadien
des brevets*
Certification

La présente atteste que les documents
ci-joints, dont la liste figure ci-dessous,
sont des copies authentiques des docu-
ments déposés au Bureau des brevets.

*Canadian Patent
Office*
Certification

This is to certify that the documents
attached hereto and identified below are
true copies of the documents on file in
the Patent Office.

Specification and drawings, as originally filed, with Application for Patent Serial No:
2,487,529, on November 15, 2004, by **B&B BEHEER NV**, assignee of Erik Buntinx, for
"Use of D4 and 5-HT_{2A} Antagonists, Inverse Agonists or Partial Agonists".

Deacy Passhuis
Agent certificateur/Certifying Officer

June 8, 2005

Date

Canada

(CIPQ 68)
31-03-04

OPIC CIPO

Abstract

The present invention relates to the use of compounds and compositions of compounds having D4 and 5-HT2A antagonistic, partial agonistic or inverse agonistic activity for the treatment of the underlying dysregulation of the emotional functionality of mental disorders (i.e. affect instability – hypersensitivity – hyperaesthesia – dissociative phenomena - etc).

5 The invention also relates to methods comprising administering to a patient diagnosed as having a neuropsychiatric disorder a pharmaceutical composition containing (i) compounds having D4 antagonistic, partial agonistic or inverse agonistic activity and (ii) compounds having 5-HT2A antagonistic, partial agonistic or inverse agonistic, and (iii) any known

10 medicinal compound and compositions of said compounds. The combined D4 and 5-HT2A antagonistic, partial agonistic or inverse agonistic effects may reside within the same chemical or biological compound or in two different chemical and/or biological compounds.

USE OF D4 AND 5-HT2A ANTAGONISTS, INVERSE AGONISTS OR PARTIAL AGONISTS

Field of the invention

The invention relates to the field of neuropsychiatry. More specifically, the invention relates to the use of compounds, which have D4 and 5-HT2A antagonist, inverse agonist or partial agonist activity, for the preparation of medicaments.

Background of the invention

Conventionally, mental disorders are divided into types based on criteria sets with defining features. DSM-IV (*American Psychiatric Association*, (1993 – ISBN 0 – 89042 – 061 – 0)) is the in the art well-known golden standard of such a categorical classification. In DSM-IV, there is no assumption that each category of mental disorder is a completely discrete entity with absolute boundaries dividing it from other mental disorders or from no mental disorder. There is also no assumption that all individuals described as having the same mental disorder are alike in all important ways. Individuals sharing a diagnosis are likely to be heterogeneous even in regard to the defining features of the diagnosis. Thus, the categorical defined mental disorders as mood and anxiety disorders are having an external and even internal variable co-incidence of symptoms concerning e.g. mood, anxiety, perception, feeding, somatic sensations, sexual functions, sleep, cognitive functioning, impulse control, attention, substance use, personality, bereavement, identity, phase of life, abuse or neglect and other aspects of behavior.

In a dimensional system, clinical presentations are classified based on quantification of attributes i.e. dysfunctions rather than the assignment to categories and works best in describing phenomena that are distributed continuously and that do not have clear boundaries.

Emotion dysregulation is known as such an attribution or dysfunction that plays an important role in the development and course of mental disorders (*Gross, J. J. & Munoz, R. F., 1995, Emotion regulation and mental health, Clinical Psychology: Science and Practice, 2, 151-164; Mennin, D.S., Heimberg, R. G., Turk, C. L. & Fresco, D. M., 2002, Applying an emotion regulation framework to integrative approaches to generalized anxiety disorder, Clinical Psychology: Science and Practice, 9, 85-90; Linehan, M. M., 1993, Cognitive-behavioral treatment of borderline personality disorder, New York, The Guilford Press; Gratz, K. L., Roemer, L., 2001 & 2004, Multidimensional assessment of emotion regulation and dysregulation: development, factor structure, and initial validation of the Difficulties in*

Emotion Regulation Scale, Annual meeting of the Association for Advancement of Behavior Therapy, Nov. 2001 & *Journal of Psychopathology and Behavioral Assessment*, Vol. 26, No. 1, March 2004) besides behavioural and cognitive dysfunctions. D4 dopamine receptors (D4DR), almost exclusively present in the mesocortical and mesolimbic systems (O'Malley, K. L., Harmon, S., Tang, L., Todd, R. D., *The rat dopamine D4 receptor: sequence, gene structure, and demonstration of expression in the cardiovascular system*, *New Biol.*, 4, 137-46, 1992), are in the art known as modulators of emotion and cognition. D4DR agonistic activity gives a behavioural sensitisation; D4DR antagonistic activity leads to an emotion modulation (Svensson, T. H., Mathé, A. A., *Monoaminergic Transmitter Systems, Biological Psychiatry* (eds. D'Haenen, H., et al.), 45-66, 2002, John Wiley & Sons, Ltd). Data demonstrate that agonism of the dopamine D4 receptors play an important role in the induction of behavioral sensitization to amphetamine and accompanying adaptations in pre- and postsynaptic neural systems associated with the mesolimbocortical dopamine projections (D. L. Feldpausch et al.; *The Journal of Pharmacology and Experimental Therapeutics* Vol. 286, Issue 1, 497-508, July 1998).

Results suggest that the antagonisms of cortical D2 dopamine receptors are a common target of traditional and atypical antipsychotics for therapeutic action. Higher *in vivo* binding to the D2 receptors in the cortex than in the basal ganglia is suggested as an indicator of favorable profile for a putative antipsychotic compound (X. Xiberas and J.L. Martinot; *The British Journal of Psychiatry* (2001) 179: 503-508). Results show that dopamine D4 receptor antagonism in the brain does not result in the same neurochemical consequences (increased dopamine metabolism or hyperprolactinemia) observed with typical neuroleptics (Smita Patel et al., *The Journal of Pharmacology and Experimental Therapeutics* Vol. 283, Issue 2, 636-647, 1997). The selective D4 dopamine receptor antagonist L-745,870 was ineffective as an antipsychotic for the treatment of neuroleptic responsive patients with acute schizophrenia (Kramer, M.S. et al., *Arch. Gen. Psychiatry* 1997 Dec; 54(12):1080).

Finally, in the biological system, mental disorders are defined on other levels of abstraction than in the categorical and dimensional system. Structural pathology (e.g. amyloid plaques in Alzheimer Disease), etiology (e.g. HIV Dementia) and deviance from a physiological norm (e.g. reduced cerebral blood flow) are often used as indicative biological markers for a mental disorder. The underlying dysregulation of various neurotransmitter systems (glutamatergic, GABAergic, cholinergic, monoaminergic (nor-adrenergic, dopaminergic, serotonergic), etc.) is the in the art used model for the explanation of the biological

determinants of the clinical presentation of mental disturbances. It is known that the Serotonin 2A Receptor (5-HT_{2A} receptor) - which is widespread in the Central Nervous System (CNS) - has a regulating role on the dysregulation of various neuro-transmitter systems. 5-HT_{2A} agonism gives several behavioural disturbances; 5-HT_{2A} antagonism leads to a governance of mood, social behaviour, anxiety, cognitive function, stress, sleep functions, nociception, sexual functions, feeding and other aspects of behaviour (*J.E. Leysen (2004) 5-HT₂ Receptors; Current Drug Targets – CNS & Neurological Disorders, 2004, 3, 11-26*).

Dysregulation of the HPA axis (hypothalamic - pituitary - adrenal axis) has frequently been reported in patients with psychiatric disorders, and is among the most robustly demonstrated neurobiological changes among psychiatric patients (*D.A. Gutman and C.B. Nemeroff, Neuroendocrinology, Biological Psychiatry (eds. D'Haenen, H., et al), 99, 2002, John Wiley & Sons, Ltd*). The resulting elevated plasma cortisol concentrations leads to an enhanced binding of serotonin for the 5-HT_{2A} receptor (*E. A. Young, Mineralocorticoid Receptor Function in Major Depression, Arch Gen Psychiatry, Jan 2003; 60: 24 - 28*) and thus agonism.

Additionally 5-HT_{2A} antagonism gives a des-inhibiting of the inhibitory effect of the 5-HT_{2A} receptor on (i) the 5-HT_{1A} receptor stimulation by serotonin (*S. M. Stahl, Newer Antidepressants and Mood Stabilizers, Essential Psychopharmacology, 265, University Press; 2 edition (June 15, 2000); ISBN: 0521646154*) and on (ii) the dopamine release in the mesocortical systems (*S. M. Stahl, Classical Antidepressants, Serotonin Selective and Noradrenargic Reuptake Inhibitors, Essential Psychopharmacology, 233, University Press; 2 edition (June 15, 2000); ISBN: 0521646154*).

Clinical or real effectiveness of psychopharma is very rare via common pooping-out; many treatment-refractory patients and up to half of patients fail to attain remission (*S. M. Stahl, Essential Psychopharmacology, Depression and Bipolar Disorders, 151, University Press; 2 edition (June 15, 2000); ISBN: 0521646154*). Implications of not attaining remission for Mental Disorders are increased relapse rates, continuing functional impairment and increased suicide rate (*S. M. Stahl, Essential Psychopharmacology, Depression and Bipolar Disorders, 152, University Press; 2 edition (June 15, 2000); ISBN: 0521646154*). Clinical causes of not attaining remission by the Current Psychopharmacological Compounds are inadequate early treatment, underlying emotion dysregulation (affecting instability – hypersensitivity – hyperaesthesia – dissociative phenomena, etc.) and competitive

antagonism. There is thus a growing need for a more efficient therapy and more efficient, selective and efficacious medicaments for treating mental disorders.

Summary of the invention

- 5 The present invention relates to the use of compounds and pharmaceutical compositions having D4 and 5-HT2A antagonistic, partial agonistic or inverse agonistic activity for the treatment of the underlying emotion dysregulation of mental disorders (e.g. affecting
- 10 Instability – hypersensitivity – hyperaesthesia – dissociative phenomena - etc.) and to methods entailing administering to a patient diagnosed as having a mental disorder a pharmaceutical composition containing (i) compounds having specific high selective D4 and 5-HT2A antagonistic, partial agonistic or inverse agonistic activity and (ii) a known medicinal compound and/or compositions of compounds. The combined D4 and 5-HT2A antagonistic, partial agonistic or inverse agonistic effects may reside within the same chemical or biological compound.
- 15 Taken into account the above mentioned (i) rare clinical or real effectiveness of psychotropics, (ii) the governance of the features and dysfunctions responsible - in a variable co-incidentally – for the clinical state of the mental disorders by D4 dopamine receptor (D4DR) and 2A serotonin receptor (5-HT2A) antagonism and (iii) the fact that 5-HT2A antagonism gives a des-inhibiting of the inhibitory effect of the 5-HT2A receptor on (a) the 5-HT1A
- 20 receptor stimulation by serotonin and on (b) the dopamine release in the mesocortical systems, the present invention relates to the use of a compound for the preparation of a medicament for treating a disease or disorder with an underlying emotion dysregulation, characterised in that said compound has (i) a selective affinity for the Dopamine-4 (D4) receptor with a pKi value equal to or higher than 8 towards the D4 receptor and less than 8
- 25 towards other Dopamine receptors, and (ii) a selective affinity for the 5-HT2A receptor with a pKi value equal to or higher than 8 towards the 5-HT2A receptor and less than 8 towards other 5-HT receptors and wherein said compound is administered to a patient in a dose ranging between 5 and 15 mg of the active ingredient. Preferably, said compound is pipamperon.
- 30 In a preferred embodiment, in a mono therapeutic context, the invention relates to the use of a compound as defined above, preferably pipamperon, for preparing a medicament for treating a disease or disorder selected from the group comprising anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders, factitious disorders, dissociative

disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, cognitive disorders, impulse control disorders, pervasive development, attention-deficit and disruptive behaviour disorders, substance-related disorders, personality disorders, psychological factors affecting medical conditions, malingering, antisocial behaviour, bereavement, occupational, identity, phase of life, academic problem, problems related to abuse or neglect.

According to a further embodiment the invention relates to the use of a first compound as defined above for the preparation of a medicament for treating a mental disease or disorder with an underlying emotion dysregulation whereby a second compound is administered simultaneously with, separate from or sequential to said first compound to augment the therapeutic effect of said second compound on said disease, or to provide a faster onset of the therapeutic effect of said second compound on said disease.

The mental diseases or disorders characterized by an underlying emotion dysregulation can be grouped into subclasses as follows: (i) non-cognitive mental disorders comprising mood disorders, anxiety disorders, psychotic disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, pervasive development disorders, attention-deficit disorders, disruptive behaviour disorders, substance-related disorders, personality disorders, psychological factors affecting medical conditions, malingering, antisocial behaviour, bereavement, occupational problems, identity problem, phase of life problem, academic problem and problems related to abuse or neglect; (ii) cognitive diseases comprising delirium, Alzheimer Disease, substance-related persisting dementia, vascular dementia, dementia due to HIV disease, dementia due to head trauma, dementia due to Parkinson Disease, dementia due to Huntington Disease, dementia due to Pick Disease, dementia due to Creutzfeldt-Jacob Disease, amnesic disorders due to a general medical condition, substance-induced persisting amnesic disorder, mild cognitive impairment disorder, other cognitive disorders; (iii) pain disorders; and (iv) Parkinson Disease.

In a preferred embodiment, the first compound is administered daily at least one day before administering said second compound.

Preferably, said second compound is characterized by the physiological property of influencing positively the activity of the Central Nervous System.

The invention also relates to a method for preparing a compound having a selective D4 and 5-HT2A antagonist, reverse agonist or partial agonist activity comprising the following steps:

(a) measuring the selective affinity of a test compound to the D4 receptor and selecting a compound that has a pKi value equal to or greater than 8 towards the D4 receptor in respect to all the other D receptors, and measuring the selective efficacy of the selected compound to the D4 receptor and selecting a compound which is a selective antagonist, inverse agonist or partial agonist of the D4 receptor; (b) measuring the selective affinity of a test compound to the 5-HT2A receptor and selecting a compound that has a pKi value equal to or greater than 8 towards the 5-HT2A receptor in respect to all the other 5HT receptors, and measuring the selective efficacy of the selected compound to the 5-HT2A receptor and selecting a compound which is a selective antagonist, inverse agonist or partial agonist of the 5-HT2A receptor; (c) identifying a compound which is selected in (a) and (b), (d) preparing the compound identified in (c).

The invention further also relates to a compound prepared by the described method.

Detailed description of the invention

The present inventors surprisingly found that compounds which have a high selective affinity towards the 5-HT2A receptor and which, at the same time have a high selective affinity towards the dopamine-4 (D4) receptor show an improved effect in treating underlying emotion dysregulation of mental disorders.

The compounds according to the invention may be chemical or biological in nature, or may be chemically synthesised. Preferably, the compounds of the invention are provided as a pharmaceutically acceptable salt.

One example of such a compound which has both a selective affinity for the 5-HT2A receptor with a pKi value equal to or higher than 8 towards the 5-HT2A receptor and less than 8 towards other 5HT receptors, and a selective affinity for the D4 receptor with a pKi value equal to or higher than 8 towards the D4 receptor and less than 8 towards other dopamine receptors is pipamperon. Pipamperon is the conventional name given for the compound of the formula 1'-[3-(p-Fluorobenzoyl)propyl]-[1,4'-bipiperidine]-4'-carboxamide. Pipamperon is also the active ingredient of for instance the commercially available Dipiperon (Janssen, Cilag B.V).

Further, the present inventors surprisingly found that the dosage of active ingredient for pipamperon in treatment (in monotherapy as well as in combination therapy as described in more detail further) could be very low compared to conventionally used dosages. Preferred dosages which, according to the invention, have been shown to be effective for treating

these mental disorders, range between 5 and 15 mg per day or between 5 and 10 mg per day. More preferably, dosages of 5, 6, 7, 8, 9, 10, 11, 12, 13, 14 or 15 mg per day are used in treatment of the diseases of the invention. In conventional pipamperon treatment, the active ingredient is available in tablets of 40 mg per tablet or in solutions of 2 mg per drop.

5 Conventional usage of high doses ranging from 40 to 360 mg is prescribed. For instance, for children up to the age of 14, doses corresponding with 2 to 6 mg per kg body weight are conventionally prescribed. The high selective affinity of pipamperon towards the 5-HT_{2A} receptor and the D₄ receptor is reflected in the low dosage which is needed for the treatment of the mental diseases listed below and also contributes to the efficacy of the treatment.

10 The mental disorders which can be treated using pipamperon in a mono therapy at such low doses are for instance anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders, factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, cognitive disorders, impulse control disorders, pervasive development, attention-deficit and disruptive behaviour disorders,
15 substance-related disorders, personality disorders, psychological factors affecting medical conditions, malingering, antisocial behaviour, bereavement, occupational, identity, phase of life, academic problem, problems related to abuse or neglect.

Mental disorders such as depression are commonly treated with serotonin re-uptake inhibitors. Unfortunately, however, these compounds can give rise to side effects in use.

20 Moreover, a substantial problem in most treatment of mental disorders is the non-response to selective serotonin re-uptake inhibitors (SSRIs). Also the onset of the therapeutic effect can be delayed undesirable.

A problem to be solved by the present invention is thus the provision of a more efficient therapy and efficient, highly selective and efficacious medicaments for treating mental
25 disorders.

The inventors found that, for instance, the non-response to selective serotonin re-uptake inhibitors (SSRIs) in depression may be declared by (partial) inhibition of the 5-HT_{1A} stimulation via 5-HT_{2A} stimulation. Des-inhibition thereof via 5-HT_{2A} antagonism seems to be an answer to this problem.

30 The present inventors found that a simultaneous or foregoing treatment with a compound having a high selective 5-HT_{2A} antagonist, inverse agonist or partial agonist activity, could lead to a greater response towards, for instance, SSRIs. However, not all compounds exhibiting 5-HT_{2A} antagonism are useful: competition between 5-HT_{2A} stimulation via serotonin and 5-HT_{2A} antagonism via the compound could be responsible for the lack of more efficacy of

compounds which have both a selective serotonin re-uptake inhibitory and 5-HT_{2A} antagonist profile, such as trazodone and nefazodone.

5 The present inventors further surprisingly found that a simultaneous or foregoing treatment with a compound having a high selective D₄ antagonist, inverse agonist or partial agonist activity in combination with a compound having a high selective 5-HT_{2A} antagonist, inverse agonist or partial agonist activity could lead to a greater response towards, for instance, SSRIs.

10 In this invention, the term "antagonist" refers to an interaction between chemicals in which one partially or completely inhibits the effect of the other, in particular agents having high affinity for a given receptor, but which do not activate this receptor.

In this invention, the term "inverse agonist" refers to a ligand which produces an effect opposite to that of the agonist by occupying the same receptor.

In this invention, the term "agonist" relates to an agent which both binds to a receptor and has an intrinsic effect.

15 In this invention, the term "partial agonist" relates to an agent with lower intrinsic activity than a full agonist, and which produces a lower maximum effect.

20 The present inventors found that a compound which binds to the 5-HT_{2A} receptor with a pK_i of at least 8 but for which the binding affinity, i.e. pK_i, towards other 5HT receptors is less than 8 in combination with a high selective affinity for the D₄ receptor, i.e. which bind to the D₄ receptor with a pK_i of at least 8 but for which the binding affinity, i.e. pK_i, towards other dopamine receptors is less than 8 also show such an improved effect in treatment. These effects, i.e. D₄ antagonism, inverse agonism or partial agonism and 5-HT_{2A} antagonism, inverse agonism or partial agonism, may reside in the same compound.

25 The term "other 5HT receptors" as used herein relate to for instance 5-HT₁ receptors (e.g. 5-HT_{1A}, 5-HT_{1B}, 5-HT_{1D}, 5-HT_{1E}, 5-HT_{1F}), 5-HT_{2B}, 5-HT_{2C}, 5-HT₆ (rat) and 5-HT₇ (rat).

By the expression "selective affinity for the 5-HT_{2A} receptor" is meant that the receptor has a higher affinity for the 5-HT_{2A} receptor than for other 5-HT receptors.

The expression "selective affinity for the D₄ receptor" means that the receptor has a higher affinity for the dopamine D₄ receptor than for other dopamine receptors.

30 The term "other dopamine receptors" are, for instance, D₁, D₂ and D₃ dopamine receptors. pK_i values of test compounds for dopamine receptors as well as 5-HT_{2A} receptors can be measured using commonly known assays.

Compounds which have a selective affinity for the D4 receptor preferably have a pKi value equal to or higher than 8 towards the D4 receptor and less than 8 towards other dopamine receptors.

Preferably, the compounds of the invention which have a selective affinity for the 5-HT_{2A} receptor (or the D4 receptor), are compounds which have a pKi value equal to or higher than 8 towards the 5-HT_{2A} receptor and the D4 receptor, and less than 8 towards other 5-HT receptors or dopamine receptors, respectively, as can be measured, for instance by methods known in the art. For instance, the "NIMH Psychoactive Drug Screening Program (PDSP)" K_i database (<http://kidb.cwru.edu/nimh/5http.php>), is a unique resource in the public domain which provides information on the abilities of drugs to interact with an expanding number of molecular targets. The PDSP K_i database serves as a data warehouse for published and internally-derived pKi, or affinity, values for a large number of drugs and drug candidates at an expanding number of G-protein coupled receptors, ion channels, transporters and enzymes. The PDSP internet site also provides for commonly used protocols and assays for measuring pKi values of 5-HT and dopamine receptors.

A preferred example of a compound which has both a selective affinity for the 5-HT_{2A} receptor with a pKi value equal to or higher than 8 towards the 5-HT_{2A} receptor and less than 8 towards other 5-HT receptors, and a selective affinity for the D4 receptor with a pKi value equal to or higher than 8 towards the D4 receptor and less than 8 towards other dopamine receptors and which is therefore useful in a combination therapy is pipamperon.

Table 1 illustrates the selective affinity of for instance pipamperon for the 5-HT_{2A} and for the D4 receptor. In addition, Table 1 also illustrates the low or absence of affinity of pipamperon for other receptors such as the adrenergic receptors Alpha 1A, Alpha 2A, Alpha 2B, Alpha 2C, Beta 1, Beta 2, and the histamine receptor H₁. As such, treating patients with pipamperon will provide for less side effects which otherwise result from simultaneous stimulation of other receptors. Therefore, and according to preferred embodiments, useful compounds according to the invention not only have a selective 5-HT_{2A} and/or D4 affinity but also a low affinity for other receptors such as the adrenergic and histamine receptors.

The low dosage which can be used in pipamperon treatment, as already described earlier, contributes to the high selective affinity of the compound towards the 5-HT_{2A} receptor and the D4 receptor and therefore also to the efficacy of the treatment.

The mental diseases or disorders characterized by an underlying emotion dysregulation can be grouped into subclasses as follows: (i) the non-cognitive mental disorders comprising mood disorders, anxiety disorders, psychotic disorders, eating disorders, premenstrual

syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, pervasive development disorders, attention-deficit disorders, disruptive behaviour disorders, substance-related disorders, personality disorders, psychological factors affecting medical conditions, malingering, antisocial behaviour, bereavement, occupational problems, identity problem, phase of life problem, academic problem and problems related to abuse or neglect; (ii) cognitive diseases comprising delirium, Alzheimer Disease, substance-related persisting dementia, vascular dementia, dementia due to HIV disease, dementia due to head trauma, dementia due to Parkinson Disease, dementia due to Huntington Disease, dementia due to Pick Disease, dementia due to Creutzfeldt-Jacob Disease, amnesic disorders due to a general medical condition, substance-induced persisting amnesic disorder, mild cognitive impairment disorder, other cognitive disorders; (iii) the pain disorders; and (iv) Parkinson Disease. In Table 5, this classification has been used for summarizing the diseases and disorders relative to known psychotropics. In Table 6, an overview of pharmacological grouping is provided, indicating the pharmacological profile numbering, the pharmacological profile, the main disease or disorder indication(s), the name of the compound, the dose range, and the company producing or selling said compound.

These diseases and their diagnosis are very clearly defined in the "Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)" published by the American Psychiatric Association. This manual sets forth diagnostic criteria, descriptions and other information to guide the classification and diagnosis of mental disorders and is commonly used in the field of neuropsychiatry. It is for instance available on the internet under: <http://www.behavenet.com/capsules/disorders/dsm4tr.htm>.

The expression "non-cognitive diseases or disorders" used in some of the embodiments of the invention comprises the following group of diseases or disorders: mood disorders, anxiety disorders, psychotic disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, pervasive development disorders, attention-deficit disorders, disruptive behaviour disorders, substance-related disorders, personality disorders, psychological factors affecting medical conditions, malingering, antisocial behaviour, bereavement, occupational problems, identity problem, phase of life problem, academic problem and problems related to abuse or neglect.

In other embodiments of the invention, the mental diseases or disorders that are characterized by an underlying emotion dysregulation belong to the group of pain disorders. For instance, the combination therapy with pipamperon is especially advantageous for management of acute pain in diseases such as, but not limited to, musculoskeletal diseases, rheumatoid arthritis, osteoarthritis and ankylosing spondylitis. For the classification of pain disorders, reference is also made to the DSM-IV where these disorders are clearly described in the section of somatoform disorders by way of internationally accepted diagnostic criteria.

In other embodiments of the invention, the 5-HT_{2A} receptor and/or Dopamine-4 receptor antagonist, inverse agonist or partial agonist (e.g. pipamperon) is used in treatment of patients having neuro-degenerative diseases or disorders, or related cognitive diseases or disorders. The diseases or disorders of the present invention are characterized by an underlying degeneration of the Central Nervous System (CNS), preferably selected from the group consisting of, but not limited to, neurodegenerative diseases such as Parkinson Disease, and in other embodiments of the invention, selected from the group of (related) cognitive diseases or disorders such as Alzheimer Disease.

For instance, Parkinson Disease, which is a chronic progressive nervous disease chiefly of later life, is linked to decreased dopamine production in the substantia nigra and is marked by tremor and weakness of resting muscles and by a shuffling gait. Dopamine agonists and even levodopa, widely used in Parkinson Disease, gives via a dopamine D₄ receptor stimulation psychiatric manifestations. The induced release of serotonin acts via 5-HT_{2A} stimulation as a "brake" on dopamine release (Young B.K., Camicioli R., Ganzini L., *Neuropsychiatric adverse effects of antiparkinsonian drugs. Characteristics, evaluation and treatment. Drugs Aging. 1997 May;10(5):367-83*). Because of the need of specific D₄ and 5-HT_{2A} antagonism in the treatment of Parkinson Disease with dopamine agonists and even levodopa, it seems reasonable to combine with a compound with a high selective D₄ and 5-HT_{2A} antagonism i.e. having merely no activity towards the other receptors especially the D₂ receptor because of the primary need of the relieve of the excessive burden of remaining dopaminergic neurons. Therefore, the use of the so-called atypical anti-psychotics or serotonin-dopamine antagonists (SDAs) is absolutely contra-indicated since their high affinity for the D₂ receptor. Even the use of serotonin releasing compounds such as SSRIs in the absence of an effective 5-HT_{2A} antagonism are contra-productive towards the Parkinson Disease symptoms although many Parkinson patients are in need for an antidepressant since major depression is a very common and disabling condition in this kind of patients.

The expression "(related) cognitive diseases or disorders" according to the invention comprises, the following group of diseases or disorders: delirium (F05), dementia (such as Alzheimer Disease (F00), vascular dementia (F01), dementia due to other general medical conditions (HIV disease (F02.4), head trauma (F06.8), Parkinson Disease (F02.3), Huntington Disease (F02.2), Pick Disease (F02.0), Creutzfeldt-Jacob Disease (F02.1) and other (F02.8)), substance-induced persisting dementia (F1x.6)), amnesic disorders due to a general medical condition (F06.8) or a substance-induced persisting amnesic disorder (F1x.6), mild cognitive impairment disorder (F06.7) and other cognitive disorders (F04). The above list of diseases is provided by way of example and is not intended to limit the invention.

For instance, Alzheimer Disease is a degenerative brain disease of unknown cause that is the most common form of dementia. Alzheimer Disease usually starts in late middle age or in old age as a memory loss for recent events spreading to memories for more distant events and progresses over the course of five to ten years to a profound intellectual decline characterized by dementia and personal helplessness. The disease is marked histologically by the degeneration of brain neurons especially in the cerebral cortex and by the presence of neurofibrillary tangles and plaques containing beta-amyloid. Because dopamine receptor D4 (DRD4) antagonism can inhibit the behavioral disturbances – merely aggression and confusion - caused by the degeneration of dopamine D2 receptors (*Esiri, M.M., The basis for behavioural disturbances in dementia, J. Neurol. Neurosurg. Psychiatry, 1996; 61(2):127–130.2*) accompanied with Alzheimer disease and 5-HT2A antagonism has an important boosting effect towards the effect of cholinesterase inhibitors such as used in the treatment by facilitating the affected dopamine release in the mesocortical dopamine pathways, a high selective D4/5-HT2A-antagonist would be a more preferable compound to combine with a cholinesterase inhibitor since this avoids the counteracting effect of the in the art used SDAs on the cognitive functioning by its dopamine receptor D2-antagonism.

These diseases and their diagnoses are very clearly defined in the "*International Statistical Classification of Diseases and Related Health Problems, 1989 Revision, Geneva, World Health Organization, 1992 (ICD-10)*". This manual sets forth diagnostic criteria, descriptions and other information to guide the classification and diagnosis of neurodegenerative disorders and is commonly used in the field of neurology. According to the ICD-10 classification, the cognitive disorders are classified under several classes of disorders, i.e. dispersed under categories F00 to F19 (see above: respective classification between parentheses). Following the DSM classification, however, they are grouped in one class of diseases or disorders.

The terms "treatment", "treating", and the like, as used herein include amelioration or elimination of a developed mental disease or condition once it has been established or alleviation of the characteristic symptoms of such disease or condition. As used herein these terms also encompass, depending on the condition of the patient, preventing the onset of a disease or condition or of symptoms associated with a disease or condition, including reducing the severity of a disease or condition or symptoms associated therewith prior to affliction with said disease or condition. Such prevention or reduction prior to affliction refers to administration of the compound or composition of the invention to a patient that is not at the time of administration afflicted with the disease or condition. "Preventing" also encompasses preventing the recurrence or relapse-prevention of a disease or condition or of symptoms associated therewith, for instance after a period of improvement. It should be clear that mental conditions may be responsible for physical complaints. In this respect, the term "treating" also includes prevention of a physical disease or condition or amelioration or elimination of the developed physical disease or condition once it has been established or alleviation of the characteristic symptoms of such conditions.

As used herein, the term "medicament" also encompasses the terms "drug", "therapeutic", "potion" or other terms which are used in the field of medicine to indicate a preparation with therapeutic or prophylactic effect.

The present inventors not only found that the selective 5-HT_{2A} and D₄ antagonists, inverse agonists or partial agonists have an effect in augmenting the therapeutic effect or in providing a faster onset of the therapeutic effect of a diversity of other pharmaceutical compounds, i.e. also named "second compounds" in the present invention, in the treatment of specific diseases or disorders. A few examples of other pharmaceutical compounds whose effects are augmented or where the onset of the effect is fastened upon simultaneous or fore-going treatment with a selective 5-HT_{2A} and D₄ antagonist, preferably pipamperon in a low dose, are nor-epinephrine re-uptake inhibitors, neuroleptic agents, dopamine antagonists, or compounds used for treating or alleviating musculoskeletal diseases or disorders. A further list of other pharmaceutical compounds or second compounds useful according to the invention is provided in Table 5. It should be clear, given the general applicable character of the invention, that this list of other pharmaceutical compounds is very brief and that the invention should not be restricted to the ones exemplified herein. It should be clear that in the present invention, pipamperon is never to be seen as a "second compound".

According to the invention, it thus has been found that the compounds having a selective 5-HT_{2A} and D₄ antagonist, inverse agonist or partial agonist activity as described above are useful for augmenting the therapeutic effect of a second compound on a disease.

According to another embodiment of the invention, it has also been found that the compounds having a selective 5-HT_{2A} and D₄ antagonist, inverse agonist or partial agonist activity as described above are useful for providing a faster onset of the therapeutic effect of a second compound on a disease.

From the above it should be clear that the selective 5-HT_{2A} and D₄ antagonist, inverse agonist or partial agonist is also named 'the first compound' in the embodiments of the invention.

According to the invention, when the 5-HT_{2A} and D₄ antagonist, inverse agonist or partial agonist activity reside in separate compounds, the term "composition" may be used. Compositions of the invention comprise a first element having (i) a selective affinity for the D₄ receptor with a pK_i value equal to or higher than 8 towards the D₄ receptor and less than 8 towards other dopamine receptors, and a second element having (ii) a selective affinity for the 5-HT_{2A} receptor with a pK_i value equal to or higher than 8 towards the 5-HT_{2A} receptor and less than 8 towards other 5-HT receptors.

The expression "the 5-HT_{2A} and D₄ antagonist, inverse agonist or partial agonist" is used herein to indicate a single compound having both activities or to indicate the composition comprising the activities in separate elements.

It should be clear that when, in the present invention, a composition of separate elements is used instead of a single compound, this composition of separate elements may be used in combination with another, i.e. a second, compound to augment the therapeutic effect of the other, i.e. the second, compound on the same or another disease.

When the 5-HT_{2A} and D₄ antagonist, inverse agonist or partial agonist or the composition comprising both elements and the second compound are administered simultaneously, the compounds or active ingredients may be present in a single pharmaceutical composition or formulation. Alternatively the compounds or active ingredients are administered in separate pharmaceutical compositions or formulations for simultaneous or separate use. The invention thus also relates to pharmaceutical compositions comprising pipamperon and a second compound of the invention and to the uses of these pharmaceutical compositions.

When the 5-HT_{2A} and D₄ antagonist, inverse agonist or partial agonist or the composition comprising both elements of the invention are administered prior to the second compound as defined, the 5-HT_{2A} and D₄ antagonist, inverse agonist or partial agonist or the composition

comprising both elements is administered at least during 1 day prior to said second compound. Preferably, the 5-HT_{2A} and D₄ antagonist, inverse agonist or partial agonist (e.g. pipamperon) or the composition comprising both elements is administered for at least 1, 2, 3, 4, 5, 6, 7, 8, 9 or 10 days prior to the administration of the second compound.

5 Preferably, the 5-HT_{2A} and D₄ antagonist, inverse agonist or partial agonist (e.g. pipamperon) or the composition comprising both elements is administered for at least 2, 3, 4 or 5 weeks prior to the administration of the second compound, or even for at least 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11 or 12 months prior to the administration of the second compound.

10 According to a preferred embodiment of the invention, the above described compounds or the composition comprising both elements having a 5-HT_{2A} and D₄ antagonist, inverse agonist or partial agonist activity are useful for augmenting the therapeutic effect of citalopram or for providing a faster onset of the therapeutic effect of citalopram.

Citalopram or citalopram hydrobromide is a selective serotonin (5-hydroxytryptamine / 5-HT) re-uptake inhibitor (SSRI) and is the conventional name given for the compound of the
15 formula (RS)-1-[3-(dimethylamino)propyl]-1-(p-fluorophenyl)-5-phthalanecarbonitrile hydrobromide.

According to an embodiment, a daily dose of active ingredient of SSRI, preferably citalopram, ranges between 10 and 40 mg per day. Preferably, daily doses of active ingredient ranging between 20 and 30 mg per day are administered. More preferably, a daily
20 dose of 10, 15, 20, 25, 30, 35 or 40 mg per day is administered.

According to another preferred embodiment of the invention, the above described compounds or the composition comprising both elements having a 5-HT_{2A} and D₄ antagonist, inverse agonist or partial agonist activity are useful for augmenting the therapeutic effect of citalopram or for providing a faster onset of the therapeutic effect of
25 fluvoxamine.

Fluvoxamine or fluvoxamine maleate (luvox, fevarin) is a selective serotonin (5-HT) reuptake inhibitor (SSRI) belonging to a new chemical series, the 2-aminoethyl oxime ethers of aralkylketones. It is chemically unrelated to other SSRIs and clomipramine. It is chemically designated as 5-methoxy-4'-(trifluoromethyl) valerophenone (E)-O-(2-aminoethyl) oxime
30 maleate (1:1).

According to an embodiment, a daily dose of active ingredient of fluvoxamine maleate ranges between 100 and 300 mg per day. Preferably, daily doses of active ingredient ranging between 150 and 200 mg per day are administered. More preferably, a daily dose of 100, 150, 200, 250 or 300 mg per day is administered.

Most of the second compounds herein described are known in the art and may be used in doses according to the supplier's or physician's prescription, or may be used according to specific embodiments described herein.

Also encompassed by the invention are pro-drugs to these second compounds or active metabolites of these compounds. For instance, for risperidone it is known that, among other products, bio transformation in the liver produces 9-hydroxyrisperidone, which is of the same pharmacological activity and intensity as parent risperidone. Therefore, also 9-hydroxyrisperidone, naturally produced or chemically synthesized may be used in the methods and uses according to the invention.

The term "active metabolite" as used herein relates to a therapeutically active compound produced by the metabolism of a parent drug. Drugs administered to treat diseases are usually transformed (metabolized) within the body into a variety of related chemical forms (metabolites), some of which may have therapeutic activity (an active metabolite).

The present invention also encompasses the use of these second compounds, administered in the form of a pharmaceutically acceptable salt in admixture with a suitable pharmaceutically acceptable excipient.

To prepare the pharmaceutical compositions, comprising the compounds or the combination of the first and second compound described herein, an effective amount of the active ingredients, in acid or base addition salt form or base form, is combined in admixture with a pharmaceutically acceptable carrier, which can take a wide variety of forms depending on the form of preparation desired for administration. These pharmaceutical compositions are desirably in unitary dosage form suitable, for administration orally, nasal, rectally, percutaneously or by parenteral injection. For example, in preparing the compositions in oral dosage form, any of the usual pharmaceutical media may be employed, such as, for example, water, glycols, oils, alcohols and the like in the case of oral liquid preparations such as suspensions, syrups, elixirs and solutions; or solid carriers such as starches, sugars, kaolin, lubricants, binders, disintegrating agents and the like in the case of powders, pills, capsules and tablets. Because of their ease in administration, tablets and capsules represent the most advantageous oral dosage unit form, in which case solid pharmaceutical carriers are obviously employed. For parenteral compositions, the carrier will usually comprise sterile water, at least in large part, though other ingredients, for example, to aid solubility, may be included.

The pharmaceutical compounds for treatment are intended for parenteral, topical, oral or local administration and generally comprise a pharmaceutically acceptable carrier and an

amount of the active ingredient sufficient to reverse or prevent the bad effects of mental disorders. The carrier may be any of those conventionally used and is limited only by chemico-physical considerations, such as solubility and lack of reactivity with the compound, and by the route of administration.

- 5 Examples of pharmaceutically acceptable acid addition salts for use in the present inventive pharmaceutical composition include those derived from mineral acids, such as hydrochloric, hydrobromic, phosphoric, metaphosphoric, nitric and sulfuric acids, and organic acids, such as tartaric, acetic, citric, malic, lactic, fumaric, benzoic, glycolic, gluconic, succinic, p-toluenesulphonic acids, and arylsulphonic, for example.
- 10 The pharmaceutically acceptable excipients described herein, for example, vehicles, adjuvants, carriers or diluents, are well-known to those who are skilled in the art and are readily available to the public. It is preferred that the pharmaceutically acceptable carrier be one that is chemically inert to the active compounds and one that has no detrimental side effects or toxicity under the conditions of use.
- 15 The following formulations for oral, aerosol, parenteral, subcutaneous, intravenous, intramuscular, interperitoneal, rectal, and vaginal administration are merely exemplary and are in no way limiting. Overall, the requirements for effective pharmaceutical carriers for parenteral compositions are well known to those of ordinary skill in the art. See
- 20 *Pharmaceutics and Pharmacy Practice*, J.B. Lippincott Company, Philadelphia, PA, Banker and Chalmers, eds., pages 238-250, (1982), and *ASHP Handbook on Injectable Drugs*, Toissel, 4th ed., pages 622-630 (1986). Topical formulations, including those that are useful for transdermal drug release, are well-known to those of skill in the art and are suitable in the context of the present invention for application to skin.

- 25 Formulations suitable for oral administration require extra considerations considering the nature of the compounds and the possible breakdown thereof if such compounds are administered orally without protecting them from the digestive secretions of the gastrointestinal tract. Such a formulation can consist of (a) liquid solutions, such as an effective amount of the compound dissolved in diluents, such as water, saline, or orange juice; (b) capsules, sachets, tablets, lozenges, and troches, each containing a predetermined
- 30 amount of the active ingredient, as solids or granules; (c) powders; (d) suspensions in an appropriate liquid; and (e) suitable emulsions. Liquid formulations may include diluents, such as water and alcohols, for example, ethanol, benzyl alcohol, and the polyethylene alcohols, either with or without the addition of a pharmaceutically acceptable surfactant, suspending agent, or emulsifying agent. Capsule forms can be of the ordinary hard- or soft-shelled
- 35 gelatin type containing, for example, surfactants, lubricants, and inert fillers, such as lactose,

sucrose, calcium phosphate, and corn starch. Tablet forms can include one or more of lactose, sucrose, mannitol, corn starch, potato starch, alginic acid, microcrystalline cellulose, acacia, gelatin, guar gum, colloidal silicon dioxide, croscarmellose sodium, talc, magnesium stearate, calcium stearate, zinc stearate, stearic acid, and other excipients, colorants, diluents, buffering agents, disintegrating agents, moistening agents, preservatives, flavoring agents, and pharmacologically compatible excipients. Lozenge forms can comprise the active ingredient in a flavor, usually sucrose and acacia or tragacanth, as well as pastilles comprising the active ingredient in an inert base, such as gelatin and glycerin, or sucrose and acacia, emulsions, gels, and the like containing, in addition to the active ingredient, such excipients as are known in the art.

The compounds of the present invention, alone or in combination with other suitable components, can be made into aerosol formulations to be administered via inhalation. For aerosol administration, the compounds are preferably supplied in finely divided form along with a surfactant and propellant. Typical percentages of compounds are 0.01%-20% by weight, preferably 1%-10%. The surfactant must, of course, be nontoxic, and preferably soluble in the propellant. Representative of such agents are the esters or partial esters of fatty acids containing from 6 to 22 carbon atoms, such as caproic, octanoic, lauric, palmitic, stearic, linoleic, linolenic, olesteric and oleic acids with an aliphatic polyhydric alcohol or its cyclic anhydride. Mixed esters, such as mixed or natural glycerides may be employed. The surfactant may constitute 0.1%-20% by weight of the compounds, preferably 0.25-5%. The balance of the compounds is ordinarily propellant. A carrier can also be included as desired, e.g., lecithin for intranasal delivery. These aerosol formulations can be placed into acceptable pressurized propellants, such as dichlorodifluoromethane, propane, nitrogen, and the like. They also may be formulated as pharmaceuticals for non-pressured preparations, such as in a nebulizer or an atomizer. Such spray formulations may be used to spray mucosa.

It will be understood that, apart from daily doses, the compounds can be administered by other schedules. For instance, the present invention also contemplates depot injection, in which a long acting form of the active compound is injected into the body, such as the muscles. From there the active compound slowly enters the rest of the body, so one injection can last from 1 to 4 weeks or even multiple months. Other form of dosage administrations relate to "once-a-week" pills, in which the ingredient is slowly released over a period of a week, and slow-release patches, e.g. a CDS (Continuous Delivery System), or Once-a-Day Transdermal Patches.

According to a further embodiment, the invention also relates to a method for preparing a compound or composition having a selective D4 and 5-HT_{2A} antagonist, reverse agonist or partial agonist. The invention also relates to the compounds prepared by the claimed method, with the proviso that said compound is not an already known compound, such as
5 pipamperon.

It should be clear that the compounds and compositions described herein are useful for treating any patient in need thereof. As used herein the term "patient" is not restricted to humans but also to other mammals, for instance, domestic animals which may also suffer from any form of a mental disease or disorder described herein.

10 The second compounds of the invention can be further grouped according to their pharmacological profile, which is summarized in Table 6.

The present invention is now described in more detail by the following embodiments. The compounds belonging to different pharmacological profiles can be further grouped according to their action on the same pathway or system as follows.

15

1: combination therapy with a 5-HT (serotonin) reuptake enhancer

The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D4 receptor, for instance pipamperon, in a combination therapy with a 5-HT (serotonin) reuptake enhancer, are chosen from the group of diseases or disorders
20 consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, substance related disorder, personality disorders, antisocial behaviour, bereavement, occupational problem, problems related to abuse or neglect and pain
25 disorders.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group
30 of diseases or disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, substance related disorder, personality disorders, antisocial behaviour, bereavement, occupational problem and problems related to
35 abuse or neglect, characterized in that pipamperon or said pharmaceutically acceptable salt

thereof is administered simultaneously with, separate from or prior to the administration of a 5-HT (serotonin) reuptake enhancer compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said 5-HT (serotonin) reuptake enhancer compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of pain disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a 5-HT (serotonin) reuptake enhancer compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said 5-HT (serotonin) reuptake enhancer compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

According to a preferred embodiment, the invention relates to the uses as described above, wherein said 5-HT (serotonin) reuptake enhancer compound is tianeptine or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof. Preferably, tianeptine is to be administered in a daily dose ranging between 25 and 50 mg of the active ingredient.

The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) a 5-HT (serotonin) reuptake enhancer, preferably tianeptine or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of a mental disease or disorder which is chosen from the group of diseases or disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, substance related disorder, personality disorder, antisocial behaviour, bereavement, occupational problem, problems related to abuse or neglect and pain disorders.

A pharmaceutical composition as described above wherein pipamperon is provided in a unitary dose of between 5 and 15 mg of the active ingredient and wherein said 5-HT

(serotonin) reuptake enhancer is tianeptine, preferably provided in a unitary dose of between 25 and 50 mg of the active ingredient.

5 2: combination therapy with a 5-HT₁ autoreceptor agonist

- 10 The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with a 5-HT₁ autoreceptor agonist, are chosen from the group of diseases or disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, substance related disorder, personality disorders, antisocial behaviour, bereavement, occupational problem, problems related to abuse or neglect and pain disorders.
- 15 The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases or disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, substance related disorder, personality disorders, antisocial behaviour, bereavement, occupational problem and problems related to abuse or neglect, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a
- 20 5-HT₁ autoreceptor agonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said 5-HT₁ autoreceptor agonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.
- 25 The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of pain disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a 5-HT₁ autoreceptor agonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said 5-HT₁ autoreceptor agonist compound, further characterized in that pipamperon is to be
- 30
- 35

administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

5 According to a preferred embodiment, the invention relates to the uses as described above, wherein said 5-HT₁ autoreceptor agonist compound is sunepitron or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof.

10 The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) a 5-HT₁ autoreceptor agonist, preferably sunepitron or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of a mental disease or disorder which is chosen from the group of diseases or disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, 15 sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, substance related disorder, personality disorder, antisocial behaviour, bereavement, occupational problem, problems related to abuse or neglect and pain disorders.

20 3: combination therapy with a 5-HT_{1A} (serotonin 1A receptor) agonist compound

The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with a 5-HT_{1A} (serotonin 1A receptor) agonist compound, are chosen from the group of diseases or disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual 25 syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender disorders, sleep disorders, adjustment disorders, impulse control disorders, attention-deficit disorders, substance-related disorder, personality disorders, antisocial behaviour, bereavement, occupational problem, problems related to abuse or neglect, pain disorders, Alzheimer Disease, substance-induced persisting 30 dementia, vascular dementia, dementia due to HIV disease, dementia due to head trauma, dementia due to Parkinson Disease, dementia due to Huntington Disease, dementia due to Pick Disease, dementia due to Creutzfeldt-Jacob Disease, amnesic disorders due to a general medical condition, substance-induced persisting amnesic disorder, mild cognitive impairment disorder, other cognitive disorders and Parkinson Disease.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender disorders, sleep disorders, adjustment disorders, impulse control disorders, attention-deficit disorders, substance-related disorder, personality disorders, antisocial behaviour, bereavement, occupational problem and problems related to abuse or neglect, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a 5-HT_{1A} (serotonin 1A receptor) agonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said 5-HT_{1A} (serotonin 1A receptor) agonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

The present invention further also relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of pain disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a 5-HT_{1A} (serotonin 1A receptor) agonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said 5-HT_{1A} (serotonin 1A receptor) agonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

The present invention further also relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a cognitive mental disease or disorder selected from the group consisting of Alzheimer Disease, substance-induced persisting dementia, vascular dementia, dementia due to HIV disease, dementia due to head trauma, dementia due to Parkinson Disease, dementia due to Huntington Disease, dementia due to Pick Disease, dementia due to Creutzfeldt-Jacob Disease, amnesic disorders due to a general medical condition, substance-induced persisting amnesic disorder, mild cognitive impairment disorder and other cognitive disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the

administration of a 5-HT_{1A} (serotonin 1A receptor) agonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said 5-HT_{1A} (serotonin 1A receptor) agonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

The present invention further also relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of Parkinson Disease, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a 5-HT_{1A} (serotonin 1A receptor) agonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said 5-HT_{1A} (serotonin 1A receptor) agonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

According to a preferred embodiment, the invention relates to the uses as described above, wherein said 5-HT_{1A} (serotonin 1A receptor) agonist compound is chosen from the group consisting of MN-305, zalospiroline, xaliproden, VPI-013 (also known as OPC-14523), tandospirone, sarizotan, PRX-00023, metanospirone, lesopitron, gepirone, flesinoxan, EMD 68843, buspirone, bupropion (preferably controlled release formulation) and alnespirone, preferably xaliproden, sarizotan, gepirone, flesinoxan and bupropion (preferably controlled release formulation) or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof. More preferably, said 5-HT_{1A} (serotonin 1A receptor) agonist is xaliproden and is to be administered in a daily dose ranging between 1 and 2 mg of the active ingredient.. Even more preferably, said 5-HT_{1A} (serotonin 1A receptor) agonist is bupropion (controlled release formulation) and is to be administered in a daily dose ranging between 150 and 450 mg of the active ingredient. Even more preferably, said 5-HT_{1A} (serotonin 1A receptor) agonist is gepirone and is to be administered in a daily dose, ranging between 20 and 80 mg of the active ingredient per day.

The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) a 5-HT_{1A} (serotonin 1A receptor) agonist, preferably chosen from the group consisting of MN-305, zalospiroline, xaliproden, VPI-013 (also known as OPC-14523), tandospirone, sarizotan, PRX-00023, metanospirone, lesopitron, gepirone, flesinoxan, EMD 68843, buspirone, bupropion (preferably controlled release formulation) and alnespirone, more

preferably xaliproden, sarizotan, gepirone, flesinoxan and bupropion (preferably controlled release formulation), or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of a mental disease or disorder which is chosen from the group of diseases or disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender disorders, sleep disorders, adjustment disorders, impulse control disorders, attention-deficit disorders, substance-related disorder, personality disorders, antisocial behaviour, bereavement, occupational problem, problems related to abuse or neglect, pain disorders, Alzheimer Disease, substance-induced persisting dementia, vascular dementia, dementia due to HIV disease, dementia due to head trauma, dementia due to Parkinson Disease, dementia due to Huntington Disease, dementia due to Pick Disease, dementia due to Creutzfeldt-Jacob Disease, amnesic disorders due to a general medical condition, substance-induced persisting amnesic disorder, mild cognitive impairment disorder, other cognitive disorders and Parkinson Disease.

The present invention also relates to a pharmaceutical composition as described above wherein pipamperon is provided in a unitary dose of between 5 and 15 mg of the active ingredient and wherein said 5-HT_{1A} (serotonin 1A receptor) agonist is xaliproden, preferably provided in a unitary dose of between 1 and 2 mg of the active ingredient.

The present invention also relates to a pharmaceutical composition as described above wherein pipamperon is provided in a unitary dose of between 5 and 15 mg of the active ingredient and wherein said 5-HT_{1A} (serotonin 1A receptor) agonist is bupropion (controlled release formulation), preferably provided in a unitary dose of between 150 and 450 mg of the active ingredient.

The present invention also relates to a pharmaceutical composition as described above wherein pipamperon is provided in a unitary dose of between 5 and 15 mg of the active ingredient and wherein said 5-HT_{1A} (serotonin 1A receptor) agonist is gepirone, preferably provided in a unitary dose of between 20 and 80 mg of the active ingredient.

4: combination therapy with a 5-HT_{1A} (serotonin 1A receptor) antagonist compound

The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with a 5-

HT1A (serotonin 1A receptor) antagonist compound, are chosen from the group of diseases or disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender disorders, adjustment disorders, impulse control disorders, substance-related disorder, personality disorders, antisocial behaviour, bereavement, occupational problem and problems related to abuse or neglect.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender disorders, adjustment disorders, impulse control disorders, attention-deficit disorders, substance-related disorder, personality disorders, antisocial behaviour, bereavement, occupational problem and problems related to abuse or neglect, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a 5-HT1A antagonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said 5-HT1A antagonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

According to a preferred embodiment, the invention relates to the uses as described above, wherein said 5-HT1A antagonist compound is chosen from the group consisting of robalzotan tartrate hydrate and NAD299 or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof.

The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) a 5-HT1A antagonist, preferably chosen from the group consisting of robalzotan tartrate hydrate and NAD299, or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder which is chosen from the group of diseases or disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender disorders, adjustment disorders, impulse control disorders, attention-deficit disorders,

substance-related disorder, personality disorders, antisocial behaviour, bereavement, occupational problem and problems related to abuse or neglect.

5: combination therapy with a 5-HT1B (serotonin 1B receptor) antagonist compound

5 The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT2A and D4 receptor, for instance pipamperon, in a combination therapy with a 5-HT1B (serotonin 1B receptor) antagonist compound, are chosen from the group of diseases or disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender disorders, sleep disorders, adjustment disorders, impulse control disorders, substance-related disorder, personality disorders, antisocial behaviour, bereavement, occupational problem and problems related to abuse or neglect.

15 The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender disorders, sleep disorders, adjustment disorders, impulse control disorders, attention-deficit disorders, substance-related disorder, personality disorders, antisocial behaviour, bereavement, occupational problem and problems related to abuse or neglect, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a 5-HT1B antagonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said 5-HT1B antagonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

30 According to a preferred embodiment, the invention relates to the use as described above, wherein said 5-HT1B antagonist compound is chosen from the group consisting of elzasonan, AZD1134 and AR-A2, preferably elzasonan, or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof.

35 The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) a 5-HT1B antagonist, preferably chosen from the group consisting of elzasonan, AZD1134 and AR-A2, preferably elzasonan, or a pro-drug or an active metabolite thereof, or

a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder which is chosen from the group of diseases or disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender disorders, sleep disorders, adjustment disorders, impulse control disorders, attention-deficit disorders, substance-related disorder, personality disorders, antisocial behaviour, bereavement, occupational problem and problems related to abuse or neglect.

6: combination therapy with a 5-HT_{2B} (serotonin 2B receptor) antagonist compound

The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with a 5-HT_{2B} (serotonin 2B receptor) antagonist compound, are chosen from the group of diseases or disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender disorders, sleep disorders, adjustment disorders, impulse control disorders, substance-related disorder, personality disorders, antisocial behaviour, bereavement, occupational problem, problems related to abuse or neglect and pain disorders.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender disorders, sleep disorders, adjustment disorders, impulse control disorders, substance-related disorder, personality disorders, antisocial behaviour, bereavement, occupational problem, problems related to abuse or neglect, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a 5-HT_{2B} antagonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said 5-HT_{2B} antagonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of pain disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from
5 or prior to the administration of a 5-HT2B antagonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said 5-HT2B antagonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

10 According to a preferred embodiment, the invention relates to the uses as described above, wherein said 5-HT2B antagonist compound is agomelatine or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof. Preferably, agomelatine is to be administered in a daily dose ranging between 25 and 50 mg of the active ingredient.

15 The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) a 5-HT2B antagonist, preferably agomelatine or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of
20 a mental disease or disorder which is chosen from the group of diseases or disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender disorders, sleep disorders, adjustment disorders, impulse control disorders, substance-related disorder, personality disorders, antisocial behaviour, bereavement, occupational problem, problems related to abuse or neglect and pain
25 disorders.

A pharmaceutical composition as described above wherein pipamperon is provided in a unitary dose of between 5 and 15 mg of the active ingredient and wherein said 5-HT2B antagonist is agomelatine, preferably provided in a unitary dose of between 25 and 50 mg of
30 the active ingredient.

7: combination therapy with a 5-HT2C (serotonin 2C receptor) antagonist compound

The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT2A and D4 receptor, for instance pipamperon, in a combination therapy with a 5-
35 HT2C (serotonin 2C receptor) antagonist compound, are chosen from the group of diseases or disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual

5 syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender disorders, sleep disorders, adjustment disorders, impulse control disorders, substance-related disorder, personality disorders, antisocial behaviour, bereavement, occupational problem, problems related to abuse or neglect and pain disorders.

10 The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender disorders, sleep disorders, adjustment disorders, impulse control disorders, substance-related disorder, personality disorders, antisocial behaviour, bereavement, occupational problem, problems related to abuse or neglect, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a 5-HT_{2C} antagonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said 5-HT_{2C} antagonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

25 The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of pain disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a 5-HT_{2C} antagonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said 5-HT_{2C} antagonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

30 According to a preferred embodiment, the invention relates to the uses as described above, wherein said 5-HT_{2C} antagonist compound is chosen from the group consisting of SB 243213 and agomelatine or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof. Preferably, agomelatine is to be administered in a daily dose ranging between 25 and 50 mg of the active ingredient.

The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) a 5-HT_{2C} antagonist, preferably chosen from the group consisting of SB 243213 and agomelatine or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of a mental disease or disorder which is chosen from the group of diseases or disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender disorders, sleep disorders, adjustment disorders, impulse control disorders, substance-related disorder, personality disorders, antisocial behaviour, bereavement, occupational problem, problems related to abuse or neglect and pain disorders.

The present invention also relates to a pharmaceutical composition as described above wherein pipamperon is provided in a unitary dose of between 5 and 15 mg of the active ingredient and wherein said 5-HT_{2C} antagonist is agomelatine, preferably provided in a unitary dose of between 25 and 50 mg of the active ingredient.

8: combination therapy with a 5-HT₃ (serotonin 3 receptor) antagonist compound

The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with a 5-HT₃ (serotonin 3 receptor) antagonist compound, are substance-related disorders.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of substance-related disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a 5-HT₃ antagonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said 5-HT₃ antagonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

According to a preferred embodiment, the invention relates to the use as described above, wherein said 5-HT₃ antagonist compound is ondansetron or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof. Preferably, ondansetron is to be administered in a daily dose ranging between 8 and 32 mg of the active ingredient.

The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) a 5-HT₃ antagonist, preferably ondansetron or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of substance-related disorders.

The invention also relates to a pharmaceutical composition as described above wherein pipamperon is provided in a unitary dose of between 5 and 15 mg of the active ingredient and wherein said 5-HT₃ antagonist is ondansetron, preferably provided in a unitary dose of between 8 and 32 mg of the active ingredient.

9: combination therapy with a 5-HT₆ (serotonin 6 receptor) antagonist compound

The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with a 5-HT₆ (serotonin 6 receptor) antagonist compound, are chosen from the group of diseases or disorders consisting of Alzheimer Disease, substance-related persisting dementia, vascular dementia, dementia due to HIV disease, dementia due to head trauma, dementia due to Parkinson Disease, dementia due to Huntington Disease, dementia due to Pick Disease, dementia due to Creutzfeldt-Jacob Disease, amnesic disorders due to a general medical condition, substance-induced persisting amnesic disorder, mild cognitive impairment disorder and other cognitive disorders.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a cognitive disorder selected from the group of diseases and disorders consisting of Alzheimer Disease, substance-related persisting dementia, vascular dementia, dementia due to HIV disease, dementia due to head trauma, dementia due to Parkinson Disease, dementia due to Huntington Disease, dementia due to Pick Disease, dementia due to Creutzfeldt-Jacob Disease, amnesic disorders due to a general medical condition, substance-induced persisting amnesic disorder, mild cognitive impairment disorder and other cognitive disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a 5-HT₆ antagonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said 5-HT₆ antagonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

According to a preferred embodiment, the invention relates to the use as described above, wherein said 5-HT6 antagonist compound is chosen from the group consisting of SB-271046, 742457 and 271046 or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof.

The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) a 5-HT6 antagonist, preferably chosen from the group consisting of SB-271046, 742457 and 271046 or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of a mental disease or disorder which is chosen from the group of diseases or disorders consisting of Alzheimer Disease, substance-related persisting dementia, vascular dementia, dementia due to HIV disease, dementia due to head trauma, dementia due to Parkinson Disease, dementia due to Huntington Disease, dementia due to Pick Disease, dementia due to Creutzfeldt-Jacob Disease, amnestic disorders due to a general medical condition, substance-induced persisting amnestic disorder, mild cognitive impairment disorder and other cognitive disorders.

10: combination therapy with an acetylcholinesterase inhibitor compound

The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with an acetylcholinesterase inhibitor compound, are chosen from the group of diseases or disorders consisting of Alzheimer Disease, substance-related persisting dementia, vascular dementia, dementia due to HIV disease, dementia due to head trauma, dementia due to Parkinson Disease, dementia due to Huntington Disease, dementia due to Pick Disease, dementia due to Creutzfeldt-Jacob Disease, amnestic disorders due to a general medical condition, substance-induced persisting amnestic disorder, mild cognitive impairment disorder and other cognitive disorders.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a cognitive disorder selected from the group of diseases and disorders consisting of Alzheimer Disease, substance-related persisting dementia, vascular dementia, dementia due to HIV disease, dementia due to head trauma, dementia due to Parkinson Disease, dementia due to Huntington Disease, dementia due to Pick Disease, dementia due to Creutzfeldt-Jacob Disease, amnestic disorders due to a general medical

condition, substance-induced persisting amnesic disorder, mild cognitive impairment disorder, other cognitive disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of an acetylcholinesterase inhibitor compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said acetylcholinesterase inhibitor compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

According to a preferred embodiment, the invention relates to the use as described above, wherein said acetylcholinesterase inhibitor compound is chosen from the group consisting of tacrine, rivastigmine tartrate, rivastigmine, physostigmine, phenserine tartrate, metrifonate, huperzine A, galantamine (preferably extended release formulation), donepezil, dichlorvos and anseculin hydrochloride, preferably tartrate, or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof. Preferably, rivastigmine tartrate is to be administered in a daily dose ranging between 3 and 12 mg of the active ingredient. Preferably, phenserine tartrate is to be administered in a daily dose ranging between 20 and 30 mg of the active ingredient. Preferably, galantamine (extended release formulation) is to be administered in a daily dose ranging between 8 and 24 mg of the active ingredient.

The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) an acetylcholinesterase inhibitor, preferably chosen from the group consisting of tacrine, rivastigmine tartrate, rivastigmine, physostigmine, phenserine tartrate, metrifonate, huperzine A, galantamine (preferably extended release formulation), donepezil, dichlorvos and anseculin hydrochloride, preferably tartrate, or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of a mental disease or disorder which is chosen from the group of diseases or disorders consisting of Alzheimer Disease, substance-related persisting dementia, vascular dementia, dementia due to HIV disease, dementia due to head trauma, dementia due to Parkinson Disease, dementia due to Huntington Disease, dementia due to Pick Disease, dementia due to Creutzfeldt-Jacob Disease, amnesic disorders due to a general medical condition, substance-induced persisting amnesic disorder, mild cognitive impairment disorder and other cognitive disorders.

The invention also relates to a pharmaceutical composition as described above wherein pipamperon is provided in a unitary dose of between 5 and 15 mg of the active ingredient and wherein said acetylcholinesterase inhibitor is rivastigmine tartrate, preferably provided in a unitary dose of between 3 and 12 mg of the active ingredient.

5

The invention further relates to a pharmaceutical composition as described above wherein pipamperon is provided in a unitary dose of between 5 and 15 mg of the active ingredient and wherein said acetylcholinesterase inhibitor is phenserine tartrate, preferably provided in a unitary dose of between 20 and 30 mg of the active ingredient.

10

In addition, the invention relates to a pharmaceutical composition as described above wherein pipamperon is provided in a unitary dose of between 5 and 15 mg of the active ingredient and wherein said acetylcholinesterase inhibitor is galantamine (preferably extended release formulation), preferably provided in a unitary dose of between 8 and 24 mg of the active ingredient.

15

11: combination therapy with an adenosine A2a receptor antagonist compound

The mental disorder which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with an adenosine A2a receptor antagonist compound, is Parkinson disease.

20

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of Parkinson disease, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of an adenosine A2a receptor antagonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said adenosine A2a receptor antagonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

25

30

According to a preferred embodiment, the invention relates to the use as described above, wherein said adenosine A2a receptor antagonist compound is KW-6002 or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof. Preferably, KW-6002 is to be administered in a daily dose ranging between 40 and 80 mg of the active ingredient.

35

The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) an adenosine A2a receptor antagonist, preferably KW-6002 or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of Parkinson disease.

The invention also relates to a pharmaceutical composition as described above wherein pipamperon is provided in a unitary dose of between 5 and 15 mg of the active ingredient and wherein said acetylcholinesterase inhibitor is KW-6002, preferably provided in a unitary dose of between 40 and 80 mg of the active ingredient.

12: combination therapy with an adrenergic transmitter releaser

The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with an adrenergic transmitter releaser, are chosen from the group of diseases or disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, adjustment disorders, impulse control disorders, personality disorders, bereavement, occupational problem, problems related to abuse or neglect and pain disorders.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, adjustment disorders, impulse control disorders, personality disorders, bereavement, occupational problem and problems related to abuse or neglect, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of an adrenergic transmitter releaser compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said adrenergic transmitter releaser compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of pain disorders, characterized in that pipamperon or said
5 pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of an adrenergic transmitter releaser compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said adrenergic transmitter releaser compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active
10 ingredient.

According to a preferred embodiment, the invention relates to the uses as described above, wherein said adrenergic transmitter releaser compound is pipoxazole or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof. Preferably,
15 pipoxazole is to be administered in a daily dose ranging between 30 and 60 mg of the active ingredient.

The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) an adrenergic transmitter releaser, preferably pipoxazole, or a pro-drug or an active
20 metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of a mental disease or disorder which is chosen from the group of diseases or disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual
25 syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, adjustment disorders, impulse control disorders, personality disorders, bereavement, occupational problem, problems related to abuse or neglect and pain disorders.

The invention also relates to a pharmaceutical composition as described above wherein
30 pipamperon is provided in a unitary dose of between 5 and 15 mg of the active ingredient and wherein said adrenergic transmitter releaser is pipoxazole, preferably provided in a unitary dose of between 30 and 60 mg of the active ingredient.

13: combination therapy with an alpha 1 adrenoreceptor antagonist

35 The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with an

alpha 1 adrenoreceptor antagonist, are chosen from the group of diseases or disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, personality disorders, bereavement, occupational problem, problems related to abuse or neglect, pain disorders and Parkinson disease.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, personality disorders, bereavement, occupational problem and problems related to abuse or neglect, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a alpha 1 adrenoreceptor antagonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said alpha 1 adrenoreceptor antagonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of pain disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of an alpha 1 adrenoreceptor antagonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said alpha 1 adrenoreceptor antagonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of Parkinson disease, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from

or prior to the administration of an alpha 1 adrenoreceptor antagonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said alpha 1 adrenoreceptor antagonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

According to a preferred embodiment, the invention relates to the uses as described above, wherein said alpha 1 adrenoreceptor antagonist compound is chosen from the group consisting of SDZ NVI 085 and flesinoxan or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof.

The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) an alpha 1 adrenoreceptor antagonist, preferably chosen from the group consisting of SDZ NVI 085 and flesinoxan or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of a mental disease or disorder which is chosen from the group of diseases or disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, personality disorders, bereavement, occupational problem, problems related to abuse or neglect, pain disorders and Parkinson disease.

14: combination therapy with an alpha 2 adrenoreceptor antagonist

The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with an alpha 2 adrenoreceptor antagonist, are chosen from the group of diseases or disorders consisting of mood disorders, anxiety disorders, psychotic disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, substance related disorders, personality disorders, bereavement, occupational problem, problems related to abuse or neglect and pain disorders.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying

emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting of mood disorders, anxiety disorders, psychotic disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, substance related disorders, personality disorders, bereavement, occupational problem and problems related to abuse or neglect, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of an alpha 2 adrenoreceptor antagonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said alpha 2 adrenoreceptor antagonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of pain disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of an alpha 2 adrenoreceptor antagonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said alpha 2 adrenoreceptor antagonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

According to a preferred embodiment, the invention relates to the uses as described above, wherein said alpha 2 adrenoreceptor antagonist compound is chosen from the group consisting of UK-14304, sunepitron, mirtazepine, idazoxan, fluparoxan, A75200 and (R)-A 75200, preferably sunepitron or idazoxan, or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof. Preferably, idazoxan is to be administered in a daily dose ranging between 5 and 40 mg of the active ingredient.

The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) an alpha 2 adrenoreceptor antagonist, preferably chosen from the group consisting of UK-14304, sunepitron, mirtazepine, idazoxan, fluparoxan, A75200 and (R)-A 75200, preferably sunepitron or idazoxan, or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of a mental

disease or disorder which is chosen from the group of diseases or disorders consisting of mood disorders, anxiety disorders, psychotic disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, substance related disorders, personality disorders, bereavement, occupational problem, problems related to abuse or neglect and pain disorders.

The invention also relates to a pharmaceutical composition as described above wherein pipamperon is provided in a unitary dose of between 5 and 15 mg of the active ingredient and wherein said alpha 2 adrenoreceptor antagonist is Idazoxan, preferably provided in a unitary dose of between 5 and 40 mg of the active ingredient.

15: combination therapy with an AMPA receptor mediator compound

The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with an AMPA (alpha-amino-3-hydroxy-5-methyl-4-isoxazole propionate) receptor mediator compound, are chosen from the group of diseases or disorders consisting of mood disorders, anxiety disorders, psychotic disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sleep disorders, adjustment disorders, impulse control disorders, pervasive development disorders, disruptive behaviour disorders, substance-related disorders, personality disorders, psychological factors affecting medical conditions, malingering, antisocial behaviour, bereavement, occupational problem, identity problem, problems related to abuse or neglect, pain disorders, delirium, Alzheimer Disease, substance-related persisting dementia, vascular dementia, dementia due to HIV disease, dementia due to head trauma, dementia due to Parkinson Disease, dementia due to Huntington Disease, dementia due to Pick Disease, dementia due to Creutzfeldt-Jacob Disease, amnesic disorders due to a general medical condition, substance-induced persisting amnesic disorder, mild cognitive impairment disorder and other cognitive disorders.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting of mood disorders, anxiety disorders, psychotic disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sleep disorders, adjustment disorders,

impulse control disorders, pervasive development disorders, disruptive behaviour disorders, substance-related disorders, personality disorders, psychological factors affecting medical conditions, malingering, antisocial behaviour, bereavement, occupational problem, identity problem and problems related to abuse or neglect, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of an AMPA receptor mediator compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said AMPA receptor mediator compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of pain disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of an AMPA receptor mediator compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said AMPA receptor mediator compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a cognitive mental disease or disorder selected from the group of diseases and disorders consisting of delirium, Alzheimer Disease, substance-related persisting dementia, vascular dementia, dementia due to HIV disease, dementia due to head trauma, dementia due to Parkinson Disease, dementia due to Huntington Disease, dementia due to Pick Disease, dementia due to Creutzfeldt-Jacob Disease, amnesic disorders due to a general medical condition, substance-induced persisting amnesic disorder, mild cognitive impairment disorder and other cognitive disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of an AMPA receptor mediator compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said AMPA receptor mediator compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

According to a preferred embodiment, the invention relates to the uses as described above, wherein said AMPA receptor mediator compound is chosen from the group consisting of

ampakine ORG 24448/CX-619, ampakine CX-717, ampakine CX-691 and ampakine CX-516, preferably ampakine ORG 24448/CX-619, ampakine CX-717 or ampakine CX-691, or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof.

5 The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and
(b) an AMPA receptor mediator, preferably chosen from the group consisting of ampakine
ORG 24448/CX-619, ampakine CX-717, ampakine CX-691 and ampakine CX-516,
preferably ampakine ORG 24448/CX-619, ampakine CX-717 or ampakine CX-691, or a pro-
10 drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a
combined preparation for simultaneous, separate or sequential use for treating the
underlying emotion dysregulation of a mental disease or disorder which is chosen from the
group of diseases or disorders consisting of mood disorders, anxiety disorders, psychotic
disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain
15 disorders), factitious disorders, dissociative disorders, sleep disorders, adjustment disorders,
impulse control disorders, pervasive development disorders, disruptive behaviour disorders,
substance-related disorders, personality disorders, psychological factors affecting medical
conditions, malingering, antisocial behaviour, bereavement, occupational problem, identity
problem, problems related to abuse or neglect, pain disorders, delirium, Alzheimer Disease,
20 substance-related persisting dementia, vascular dementia, dementia due to HIV disease,
dementia due to head trauma, dementia due to Parkinson Disease, dementia due to
Huntington Disease, dementia due to Pick Disease, dementia due to Creutzfeldt-Jacob
Disease, amnesic disorders due to a general medical condition, substance-induced
persisting amnesic disorder, mild cognitive impairment disorder and other cognitive
disorders.

25
16: combination therapy with an amphetamine compound
The mental disorders which can be treated using compounds having a high selective affinity
for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with an
amphetamine compound, are attention-deficit disorders.

30
The present invention thus relates to the use of pipamperon or a pharmaceutically
acceptable salt thereof for the preparation of a medicament for treating the underlying
emotion dysregulation of attention deficit disorders, characterized in that pipamperon or said
pharmaceutically acceptable salt thereof is administered simultaneously with, separate from
35 or prior to the administration of an amphetamine compound to augment the therapeutic
effect or to provide a faster onset of the therapeutic effect of said amphetamine compound,

further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

According to a preferred embodiment, the invention relates to the use as described above, wherein said amphetamine compound is methylphenidate (preferably administered by the transdermal system) or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof.

The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) an amphetamine, preferably methylphenidate (preferably administered by the transdermal system) or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of attention deficit disorders.

17: combination therapy with an amyloid aggregation-inhibitor compound

The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with an amyloid aggregation-inhibitor compound, are chosen from the group of diseases or disorders consisting of Alzheimer Disease, substance-related persisting dementia, vascular dementia, dementia due to HIV disease, dementia due to head trauma, dementia due to Parkinson Disease, dementia due to Huntington Disease, dementia due to Pick Disease, dementia due to Creutzfeldt-Jacob Disease, amnesic disorders due to a general medical condition, substance-induced persisting amnesic disorder, mild cognitive impairment disorder and other cognitive disorders.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a cognitive mental disease or disorder selected from the group of diseases and disorders consisting of Alzheimer Disease, substance-related persisting dementia, vascular dementia, dementia due to HIV disease, dementia due to head trauma, dementia due to Parkinson Disease, dementia due to Huntington Disease, dementia due to Pick Disease, dementia due to Creutzfeldt-Jacob Disease, amnesic disorders due to a general medical condition, substance-induced persisting amnesic disorder, mild cognitive impairment disorder and other cognitive disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of an amyloid aggregation-inhibitor compound to augment the

therapeutic effect or to provide a faster onset of the therapeutic effect of said amyloid aggregation-inhibitor compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

5

According to a preferred embodiment, the invention relates to the use as described above, wherein said amyloid aggregation-inhibitor compound is chosen from the group consisting of APAN and Alzhemed, or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof. Preferably, Alzhemed is to be administered in a daily dose of

10

The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) an amyloid aggregation-inhibitor, preferably chosen from the group consisting of APAN and Alzhemed, or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of a mental disease or disorder which is chosen from the group of diseases or disorders consisting of Alzheimer Disease, substance-related persisting dementia, vascular dementia, dementia due to HIV disease, dementia due to head trauma, dementia due to Parkinson Disease, dementia due to Huntington Disease, dementia due to Pick Disease, dementia due to Creutzfeldt-Jacob Disease, amnesic disorders due to a general medical condition, substance-induced persisting amnesic disorder, mild cognitive impairment disorder and other cognitive disorders.

15

20

25

The invention also relates to a pharmaceutical composition as described above wherein pipamperon is provided in a unitary dose of between 5 and 15 mg of the active ingredient and wherein said amyloid aggregation-inhibitor is Alzhemed, preferably provided in a unitary dose of between 200 and 300 mg of the active ingredient.

30

18: combination therapy with an androgen receptor modulator compound

The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with an androgen receptor modulator compound, are sexual and gender identity disorders.

35

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying

emotion dysregulation of sexual and gender identity disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of an androgen receptor modulator compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said androgen receptor modulator compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

According to a preferred embodiment, the invention relates to the use as described above, wherein said androgen receptor modulator compound is LGD2226 or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof.

The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) an androgen receptor modulator, preferably LGD2226 or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of sexual and gender identity disorders.

19: combination therapy with an beta 3 adrenoreceptor agonist

The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with an beta 3 adrenoreceptor agonist, are chosen from the group of diseases or disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, substance related disorders, personality disorders, bereavement, occupational problem, problems related to abuse or neglect and pain disorders.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, substance related disorders, personality disorders, bereavement, occupational problem and problems related to abuse or neglect, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is

administered simultaneously with, separate from or prior to the administration of a beta 3 adrenoreceptor agonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said beta 3 adrenoreceptor agonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging
5 between 5 and 15 mg of the active ingredient.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of pain disorders, characterized in that pipamperon or said
10 pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of an beta 3 adrenoreceptor agonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said beta 3 adrenoreceptor agonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active
15 ingredient.

According to a preferred embodiment, the invention relates to the uses as described above, wherein said beta 3 adrenoreceptor agonist compound is SR 58611 or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof.

The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) a beta 3 adrenoreceptor agonist, preferably SR 58611 or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion
20 dysregulation of a mental disease or disorder which is chosen from the group of diseases or disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, substance related disorders, personality disorders, bereavement, occupational problem, problems related to abuse or neglect and pain disorders.
25
30

20: combination therapy with a calcium channel modulator compound

The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with a
35 calcium channel modulator compound, are chosen from the group of diseases or disorders consisting of Alzheimer Disease, substance-related persisting dementia, vascular dementia,

dementia due to HIV disease, dementia due to head trauma, dementia due to Parkinson Disease, dementia due to Huntington Disease, dementia due to Pick Disease, dementia due to Creutzfeldt-Jacob Disease, amnestic disorders due to a general medical condition, substance-induced persisting amnestic disorder, mild cognitive impairment disorder, other cognitive disorders and Parkinson disease.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a cognitive mental disease or disorder selected from the group of diseases and disorders consisting of Alzheimer Disease, substance-related persisting dementia, vascular dementia, dementia due to HIV disease, dementia due to head trauma, dementia due to Parkinson Disease, dementia due to Huntington Disease, dementia due to Pick Disease, dementia due to Creutzfeldt-Jacob Disease, amnestic disorders due to a general medical condition, substance-induced persisting amnestic disorder, mild cognitive impairment disorder and other cognitive disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a calcium channel modulator compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said calcium channel modulator compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of Parkinson disease, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a calcium channel modulator compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said calcium channel modulator compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

According to a preferred embodiment, the invention relates to the uses as described above, wherein said calcium channel modulator compound is chosen from the group consisting of safinamide and MEM 1003, or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof.

The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) a calcium channel modulator, preferably chosen from the group consisting of safinamide and MEM 1003, or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of a mental disease or disorder which is chosen from the group of diseases or disorders consisting of Alzheimer Disease, substance-related persisting dementia, vascular dementia, dementia due to HIV disease, dementia due to head trauma, dementia due to Parkinson Disease, dementia due to Huntington Disease, dementia due to Pick Disease, dementia due to Creutzfeldt-Jacob Disease, amnesic disorders due to a general medical condition, substance-induced persisting amnesic disorder, mild cognitive impairment disorder, other cognitive disorders and Parkinson disease.

21: combination therapy with a cannaboid receptor 1 (CB1) antagonist

The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with a cannaboid receptor 1 (CB1) antagonist, are chosen from the group of diseases or disorders consisting of mood disorders, anxiety disorders, psychotic disorders, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sleep disorders, adjustment disorders, impulse control disorders, pervasive development disorders, disruptive behaviour disorders, substance-related disorders, personality disorders, psychological factors affecting medical conditions, malingering, antisocial behaviour, bereavement, occupational problem, identity problem, problems related to abuse or neglect, pain disorders and delirium.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting mood disorders, anxiety disorders, psychotic disorders, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sleep disorders, adjustment disorders, impulse control disorders, pervasive development disorders, disruptive behaviour disorders, substance-related disorders, personality disorders, psychological factors affecting medical conditions, malingering, antisocial behaviour, bereavement, occupational problem, identity problem and problems related to abuse or neglect, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a

cannaboid receptor 1 antagonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said cannaboid receptor antagonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

5

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of pain disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a cannaboid receptor 1 antagonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said cannaboid receptor 1 antagonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

10

15

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of delirium, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a cannaboid receptor 1 antagonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said cannaboid receptor 1 antagonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

20

25

According to a preferred embodiment, the invention relates to the uses as described above, wherein said cannaboid receptor antagonist compound is SR 141716 or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof.

30

35

The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) a cannaboid receptor antagonist, preferably SR 141716 or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of a mental disease or disorder which is chosen from the group of diseases or disorders consisting of mood disorders, anxiety disorders, psychotic disorders, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sleep disorders, adjustment disorders, impulse control disorders, pervasive development

disorders, disruptive behaviour disorders, substance-related disorders, personality disorders, psychological factors affecting medical conditions, malingering, antisocial behaviour, bereavement, occupational problem, identity problem, problems related to abuse or neglect, pain disorders and delirium.

22: combination therapy with a cathepsin K inhibitor compound

The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with a cathepsin K inhibitor compound, are pain disorders.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of pain disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a cathepsin K inhibitor compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said cathepsin K inhibitor compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

According to a preferred embodiment, the invention relates to the use as described above, wherein said cathepsin K inhibitor compound is 462795 or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof.

The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) a cathepsin K inhibitor, preferably 462795 or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of pain disorders.

23: combination therapy with a choline uptake enhancer compound

The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with a choline uptake enhancer compound, are chosen from the group of diseases or disorders consisting of Alzheimer Disease, substance-related persisting dementia, vascular dementia, dementia due to HIV disease, dementia due to head trauma, dementia due to Parkinson Disease, dementia due to Huntington Disease, dementia due to Pick Disease, dementia due

to Creutzfeldt-Jacob Disease, amnesic disorders due to a general medical condition, substance-induced persisting amnesic disorder, mild cognitive impairment disorder and other cognitive disorders.

- 5 The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a cognitive mental disease or disorder selected from the group of diseases and disorders consisting of Alzheimer Disease, substance-related persisting dementia, vascular dementia, dementia due to HIV disease, dementia due to head trauma, 10 dementia due to Parkinson Disease, dementia due to Huntington Disease, dementia due to Pick Disease, dementia due to Creutzfeldt-Jacob Disease, amnesic disorders due to a general medical condition, substance-induced persisting amnesic disorder, mild cognitive impairment disorder and other cognitive disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from 15 or prior to the administration of a choline uptake enhancer compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said choline uptake enhancer compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.
- 20 According to a preferred embodiment, the invention relates to the use as described above, wherein said choline uptake enhancer compound is MKC-231 or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof. Preferably, MKC-231 is to be administered in a daily dose of between 20 and 160 mg of the active ingredient.
- 25 The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) a choline uptake enhancer, preferably MKC-231 or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of a mental disease or disorder which is chosen from the group of diseases or disorders 30 consisting of Alzheimer Disease, substance-related persisting dementia, vascular dementia, dementia due to HIV disease, dementia due to head trauma, dementia due to Parkinson Disease, dementia due to Huntington Disease, dementia due to Pick Disease, dementia due to Creutzfeldt-Jacob Disease, amnesic disorders due to a general medical condition, substance-induced persisting amnesic disorder, mild cognitive impairment disorder and 35 other cognitive disorders.

The invention also relates to a pharmaceutical composition as described above wherein pipamperon is provided in a unitary dose of between 5 and 15 mg of the active ingredient and wherein said choline uptake enhancer is MKC-231, preferably provided in a unitary dose of between 20 and 160 mg of the active ingredient.

5

24: combination therapy with a COX-2 inhibitor compound

The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with a COX-2 inhibitor compound, are pain disorders.

10

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of pain disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a COX-2 inhibitor compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said COX-2 inhibitor compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

15

According to a preferred embodiment, the invention relates to the use as described above, wherein said COX-2 inhibitor compound is chosen from the group consisting of valdecoxib, rofecoxib, parecoxib, etoricoxib, COX 189, celecoxib and ABT-963, preferably parecoxib, etoricoxib or COX 189, or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof. Preferably, parecoxib is to be administered in a daily dose of between 20 and 80 mg of the active ingredient. Preferably, etoricoxib is to be administered in a daily dose of between 20 and 120 mg of the active ingredient. Preferably, COX 189 is to be administered in a daily dose of between 100 and 800 mg of the active ingredient.

25

The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) a COX-2 inhibitor, preferably chosen from the group consisting of valdecoxib, rofecoxib, parecoxib, etoricoxib, COX 189, celecoxib and ABT-963, preferably parecoxib, etoricoxib or COX 189, or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of pain disorders.

30

35

The invention also relates to a pharmaceutical composition as described above wherein pipamperon is provided in a unitary dose of between 5 and 15 mg of the active ingredient and wherein said COX-2 inhibitor is parecoxib, preferably provided in a unitary dose of between 20 and 80 mg of the active ingredient.

5

The invention also relates to a pharmaceutical composition as described above wherein pipamperon is provided in a unitary dose of between 5 and 15 mg of the active ingredient and wherein said COX-2 inhibitor is etoricoxib, preferably provided in a unitary dose of between 20 and 120 mg of the active ingredient.

10

The invention also relates to a pharmaceutical composition as described above wherein pipamperon is provided in a unitary dose of between 5 and 15 mg of the active ingredient and wherein said COX-2 inhibitor is COX 189, preferably provided in a unitary dose of between 100 and 800 mg of the active ingredient.

15

25: combination therapy with a COX-inhibiting nitric oxide donator (CINOD) compound

The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with a COX-inhibiting nitric oxide donator (CINOD) compound, are pain disorders.

20

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of pain disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a COX-inhibiting nitric oxide donator (CINOD) compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said COX-inhibiting nitric oxide donator (CINOD) compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

25

According to a preferred embodiment, the invention relates to the use as described above, wherein said COX-inhibiting nitric oxide donator (CINOD) compound is chosen from the group consisting of AZD4717 and AZD3582 or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof. Preferably, AZD3582 is to be administered in a daily dose ranging between 93.75 and 750 mg of the active ingredient.

30

35

The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) a COX-inhibiting nitric oxide donator (CINOD), preferably chosen from the group consisting of AZD4717 and AZD3582 or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of pain disorders.

The invention also relates to a pharmaceutical composition as described above wherein pipamperon is provided in a unitary dose of between 5 and 15 mg of the active ingredient and wherein said COX-inhibiting nitric oxide donator (CINOD) is AZD3582, preferably provided in a unitary dose of between 93.75 and 750 mg of the active ingredient.

26: combination therapy with a CRF1 (corticotropin-releasing factor receptor 1) antagonist

The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with a CRF1 (Corticotropin-Releasing Factor receptor 1) antagonist, are chosen from the group of diseases or disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, substance related disorders, personality disorders, bereavement, occupational problem, problems related to abuse or neglect and pain disorders.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, substance related disorders, personality disorders, bereavement, occupational problem and problems related to abuse or neglect, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a CRF1 antagonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said CRF1 antagonist compound, further characterized in that

pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

5 The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of pain disorders, characterized in that pipamperon or said
10 pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a CRF1 antagonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said CRF1 antagonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

According to a preferred embodiment, the invention relates to the uses as described above, wherein said CRF1 antagonist compound is chosen from the group consisting of R121919,
15 NBI-34041, elzasonan, CP-448,187, CP-154-526, AAG 561 and 723620, preferably R121919, elzasonan or AAG 561, or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof. Preferably, R121919 is to be administered in a daily dose of between 5 and 80 mg of the active ingredient.

20 The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) a CRF1 antagonist, preferably chosen from the group consisting of R121919, NBI-34041, elzasonan, CP-448,187, CP-154-526, AAG 561 and 723620, preferably R121919, elzasonan or AAG 561, or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for
25 treating the underlying emotion dysregulation of a mental disease or disorder which is chosen from the group of diseases or disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, substance related
30 disorders, personality disorders, bereavement, occupational problem, problems related to abuse or neglect and pain disorders.

The invention also relates to a pharmaceutical composition as described above wherein pipamperon is provided in a unitary dose of between 5 and 15 mg of the active ingredient
35 and wherein said CRF1 antagonist is R121919, preferably provided in a unitary dose of between 5 and 80 mg of the active ingredient.

27: combination therapy with a D1 (dopamine 1) receptor agonist

The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with a D1 (dopamine 1) receptor agonist, are chosen from the group of diseases or disorders consisting of substance related disorders and Parkinson disease.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of substance related disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a D1 receptor agonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said D1 receptor agonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of Parkinson disease, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a D1 receptor agonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said D1 receptor agonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

According to a preferred embodiment, the invention relates to the uses as described above, wherein said D1 receptor agonist compound is DAS-431 or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof.

The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) a D1 receptor agonist, preferably DAS-431 or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of a mental disease or disorder which is chosen from the group of diseases or disorders consisting of substance related disorders and Parkinson disease.

28: combination therapy with D2 (dopamine 2) receptor antagonist

The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with D₂ (dopamine 2) receptor antagonist, are chosen from the group of diseases or disorders consisting of mood disorders, psychotic disorders, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sleep disorders, adjustment disorders, impulse control disorders, pervasive development disorders, disruptive behaviour disorders, substance-related disorders, personality disorders, psychological factors affecting medical conditions, malingering, antisocial behaviour, bereavement, occupational problem, identity problem, problems related to abuse or neglect, pain disorders and delirium.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting mood disorders, psychotic disorders, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sleep disorders, adjustment disorders, impulse control disorders, pervasive development disorders, disruptive behaviour disorders, substance-related disorders, personality disorders, psychological factors affecting medical conditions, malingering, antisocial behaviour, bereavement, occupational problem, identity problem and problems related to abuse or neglect, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a D₂ receptor antagonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said D₂ receptor antagonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of pain disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a D₂ receptor antagonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said D₂ receptor antagonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of delirium, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a D2 receptor antagonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said D2 receptor antagonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

According to a preferred embodiment, the invention relates to the uses as described above, wherein said D2 receptor antagonist compound is chosen from the group consisting of bifeprunox, amisulpride aminosultopride and amisulpride, preferably bifeprunox, or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof.

The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) a D2 receptor antagonist, preferably chosen from the group consisting of bifeprunox, amisulpride aminosultopride and amisulpride, preferably bifeprunox, or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of a mental disease or disorder which is chosen from the group of diseases or disorders consisting of mood disorders, psychotic disorders, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sleep disorders, adjustment disorders, impulse control disorders, pervasive development disorders, disruptive behaviour disorders, substance-related disorders, personality disorders, psychological factors affecting medical conditions, malingering, antisocial behaviour, bereavement, occupational problem, identity problem, problems related to abuse or neglect, pain disorders and delirium.

29: combination therapy with D3 (dopamine 3) receptor antagonist

The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with D₃ (dopamine 3) receptor antagonist, are chosen from the group of diseases or disorders consisting of psychotic disorders, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sleep disorders, adjustment disorders, impulse control disorders, pervasive development disorders, disruptive behaviour disorders, substance-related disorders, personality disorders, psychological factors affecting medical conditions,

malinger, antisocial behaviour, bereavement, occupational problem, identity problem, problems related to abuse or neglect, pain disorders, delirium and Parkinson disease.

5 The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting of psychotic disorders, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sleep disorders, adjustment disorders, impulse control disorders, pervasive development disorders, disruptive
10 behaviour disorders, substance-related disorders, personality disorders, psychological factors affecting medical conditions, malingering, antisocial behaviour, bereavement, occupational problem, identity problem, problems related to abuse or neglect, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a D3 receptor antagonist
15 compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said D3 receptor antagonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

20 The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of pain disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a D3 receptor antagonist compound to augment the
25 therapeutic effect or to provide a faster onset of the therapeutic effect of said D3 receptor antagonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

30 The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of delirium, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a D3 receptor antagonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said D3 receptor antagonist compound,
35 further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of Parkinson disease, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a D3 receptor agonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said D3 receptor agonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

According to a preferred embodiment, the invention relates to the uses as described above, wherein said D3 receptor antagonist compound is chosen from the group consisting of BSF-201640 and PD 58491, or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof.

The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) a D3 receptor antagonist, preferably chosen from the group consisting of BSF-201640 and PD 58491, or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of a mental disease or disorder which is chosen from the group of diseases or disorders consisting of psychotic disorders, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sleep disorders, adjustment disorders, impulse control disorders, pervasive development disorders, disruptive behaviour disorders, substance-related disorders, personality disorders, psychological factors affecting medical conditions, malingering, antisocial behaviour, bereavement, occupational problem, identity problem, problems related to abuse or neglect, pain disorders, delirium and Parkinson disease.

30: combination therapy with a DA (dopamine) uptake inhibitor

The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with a DA (dopamine) uptake inhibitor, are chosen from the group of diseases or disorders consisting of substance related disorders and Parkinson disease.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying

emotion dysregulation of substance related disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a DA uptake inhibitor compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said DA uptake inhibitor compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of Parkinson disease, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a DA uptake inhibitor compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said DA uptake inhibitor compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

According to a preferred embodiment, the invention relates to the uses as described above, wherein said DA uptake inhibitor compound is chosen from the group consisting of safinamide and GBR 12909, or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof.

The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) a D2 receptor antagonist, preferably chosen from the group consisting of safinamide and GBR 12909, or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of a mental disease or disorder which is chosen from the group of diseases or disorders consisting of substance related disorders and Parkinson disease.

31: combination therapy with an dopamine (receptor) agonist

The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with an dopamine (receptor) agonist, are chosen from the group of diseases or disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, adjustment disorders, impulse control disorders, attention-deficit disorders, substance-related disorders,

personality disorders, problems related to abuse or neglect, pain disorders and Parkinson disease.

5 The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, adjustment disorders, impulse control disorders, attention-deficit disorders, substance-related disorders, personality disorders and problems related to abuse or neglect,, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a dopamine (receptor) agonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said dopamine (receptor) agonist compound, further
10 characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

20 The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of pain disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a dopamine (receptor) agonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said dopamine (receptor) agonist compound, further characterized in that pipamperon is to be administered
25 to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

30 The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of Parkinson disease, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a dopamine (receptor) agonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said dopamine (receptor) agonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

According to a preferred embodiment, the invention relates to the uses as described above, wherein said dopamine (receptor) agonist compound is chosen from the group consisting of sumanirole, SLV 308, sarizotan, S32504, rotigotine (preferably a Once-a-Day Transdermal Patch), ropinirole HCL (preferably controlled-release formulation), pramipexole, DAB452, cabergoline, bromocriptine, alaptide amantadine, bromocriptine, cabergoline lisuride and pergolide, preferably sumanirole, rotigotine (preferably a Once-a-Day Transdermal Patch), pergolide or ropinirole HCL (preferably controlled-release formulation), or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof. Preferably, sumanirole is to be administered in a daily dose of between 4 and 16 mg of the active ingredient. Preferably, rotigotine (Once-a-Day Transdermal Patch) is to be administered in a daily dose of between 4.5 and 13.5 mg of the active ingredient. Preferably, ropinirole HCL (controlled-release formulation) is to be administered in a daily dose of between 0.75 and 24 mg of the active ingredient. Preferably, pergolide is to be administered in a daily dose of between 0.5 and 10 mg of the active ingredient.

The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) a dopamine (receptor) agonist, preferably chosen from the group consisting of sumanirole, SLV 308, sarizotan, S32504, rotigotine (preferably a Once-a-Day Transdermal Patch), ropinirole HCL (preferably controlled-release formulation), pramipexole, DAB452, cabergoline, bromocriptine, alaptide amantadine, bromocriptine, cabergoline lisuride and pergolide, more preferably sumanirole, rotigotine (preferably a Once-a-Day Transdermal Patch), ropinirole HCL (preferably controlled-release formulation) or pergolide, or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of a mental disease or disorder which is chosen from the group of diseases or disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, adjustment disorders, impulse control disorders, attention-deficit disorders, substance-related disorders, personality disorders, problems related to abuse or neglect, pain disorders and Parkinson disease.

The invention also relates to a pharmaceutical composition as described above wherein pipamperon is provided in a unitary dose of between 5 and 15 mg of the active ingredient and wherein said dopamine (receptor) agonist is sumanirole, preferably provided in a unitary dose of between 4 and 16 mg of the active ingredient.

The invention also relates to a pharmaceutical composition as described above wherein pipamperon is provided in a unitary dose of between 5 and 15 mg of the active ingredient and wherein said dopamine (receptor) agonist is rotigotine (Once-a-Day Transdermal Patch), preferably provided in a unitary dose of between 4.5 and 13.5 mg of the active ingredient.

The invention also relates to a pharmaceutical composition as described above wherein pipamperon is provided in a unitary dose of between 5 and 15 mg of the active ingredient and wherein said dopamine (receptor) agonist is ropinirole HCL (controlled-release formulation), preferably provided in a unitary dose of between 0.75 and 24 mg of the active ingredient.

The invention also relates to a pharmaceutical composition as described above wherein pipamperon is provided in a unitary dose of between 5 and 15 mg of the active ingredient and wherein said dopamine (receptor) agonist is pergolide, preferably provided in a unitary dose of between 0.5 and 10 mg of the active ingredient.

32: combination therapy with a compound activating ERK (extracellular signal-related kinase)

The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with a compound that activates ERK (extracellular signal-related kinase), are chosen from the group of diseases or disorders consisting of Alzheimer Disease, substance-related persisting dementia, vascular dementia, dementia due to HIV disease, dementia due to head trauma, dementia due to Parkinson Disease, dementia due to Huntington Disease, dementia due to Pick Disease, dementia due to Creutzfeldt-Jacob Disease, amnestic disorders due to a general medical condition, substance-induced persisting amnestic disorder, mild cognitive impairment disorder and other cognitive disorders.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a cognitive mental disease or disorder selected from the group of diseases and disorders consisting of Alzheimer Disease, substance-related persisting dementia, vascular dementia, dementia due to HIV disease, dementia due to head trauma, dementia due to Parkinson Disease, dementia due to Huntington Disease, dementia due to Pick Disease, dementia due to Creutzfeldt-Jacob Disease, amnestic disorders due to a

general medical condition, substance-induced persisting amnesic disorder, mild cognitive impairment disorder and other cognitive disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a compound that activates ERK (extracellular signal-related kinase) to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said compound that activates ERK (extracellular signal-related kinase), further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

According to a preferred embodiment, the invention relates to the use as described above, wherein said compound that activates ERK (extracellular signal-related kinase) is CPI-1189 or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof. Preferably, CPI-1189 is to be administered in a daily dose of between 50 and 100 mg of the active ingredient.

The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) a compound that activates ERK (extracellular signal-related kinase), preferably CPI-1189 or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of a mental disease or disorder which is chosen from the group of diseases or disorders consisting of Alzheimer Disease, substance-related persisting dementia, vascular dementia, dementia due to HIV disease, dementia due to head trauma, dementia due to Parkinson Disease, dementia due to Huntington Disease, dementia due to Pick Disease, dementia due to Creutzfeldt-Jacob Disease, amnesic disorders due to a general medical condition, substance-induced persisting amnesic disorder, mild cognitive impairment disorder and other cognitive disorders.

The invention also relates to a pharmaceutical composition as described above wherein pipamperon is provided in a unitary dose of between 5 and 15 mg of the active ingredient and wherein said compound that activates ERK (extracellular signal-related kinase) is CPI-1189, preferably provided in a unitary dose of between 50 and 100 mg of the active ingredient.

33: combination therapy with a GABA (gamma-aminobutyric acid) agonist compound

The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with a

GABA (gamma-aminobutyric acid) agonist compound, are chosen from the group of diseases or disorders consisting of Alzheimer Disease, substance-related persisting dementia, vascular dementia, dementia due to HIV disease, dementia due to head trauma, dementia due to Parkinson Disease, dementia due to Huntington Disease, dementia due to Pick Disease, dementia due to Creutzfeldt-Jacob Disease, amnestic disorders due to a general medical condition, substance-induced persisting amnestic disorder, mild cognitive impairment disorder and other cognitive disorders.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a cognitive mental disease or disorder selected from the group of diseases and disorders consisting of Alzheimer Disease, substance-related persisting dementia, vascular dementia, dementia due to HIV disease, dementia due to head trauma, dementia due to Parkinson Disease, dementia due to Huntington Disease, dementia due to Pick Disease, dementia due to Creutzfeldt-Jacob Disease, amnestic disorders due to a general medical condition, substance-induced persisting amnestic disorder, mild cognitive impairment disorder and other cognitive disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a GABA agonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said GABA agonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

According to a preferred embodiment, the invention relates to the use as described above, wherein said GABA agonist compound is nefiracetam or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof.

The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) a GABA agonist, preferably nefiracetam or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of a mental disease or disorder which is chosen from the group of diseases or disorders consisting of Alzheimer Disease, substance-related persisting dementia, vascular dementia, dementia due to HIV disease, dementia due to head trauma, dementia due to Parkinson Disease, dementia due to Huntington Disease, dementia due to Pick Disease, dementia due to Creutzfeldt-Jacob Disease, amnestic disorders due to a general medical condition,

substance-induced persisting amnesic disorder, mild cognitive impairment disorder and other cognitive disorders.

34: combination therapy with a GABA-A agonist compound

5 The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with a GABA-A (gamma-aminobutyric acid receptor A) agonist compound, are sleep disorders.

10 The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of sleep disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a GABA-A agonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said GABA-A agonist
15 compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

According to a preferred embodiment, the invention relates to the use as described above, wherein said GABA-A agonist compound is gaboxadol or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof. Preferably, gaboxadol is to be
20 administered in a daily dose of between 5 and 20 mg of the active ingredient.

The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) an GABA-A agonist, preferably gaboxadol or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous,
25 separate or sequential use for treating the underlying emotion dysregulation of sleep disorders.

The invention also relates to a pharmaceutical composition as described above wherein pipamperon is provided in a unitary dose of between 5 and 15 mg of the active ingredient
30 and wherein said GABA-A agonist is Gaboxadol, preferably provided in a unitary dose of between 5 and 20 mg of the active ingredient.

35: combination therapy with a GABA-A modulator compound

35 The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with a

GABA-A (gamma-aminobutyric acid receptor A) modulator compound, are chosen from the group of diseases or disorders consisting of anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, personality disorders, bereavement, occupational problem and problems related to abuse or neglect.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting of anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, personality disorders, bereavement, occupational problem and problems related to abuse or neglect, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a GABA-A modulator compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said GABA-A modulator compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

According to a preferred embodiment, the invention relates to the use as described above, wherein said GABA-A modulator compound is chosen from the group consisting of zolpidem (preferably MR sustained-release version), zaleplon (preferably extended-release formulation), SL 65.1498, SEP174559, pagoclone, NGD 96-3, indiplon, eszopiclone, CP-730,330 (NGD 96-3) and ocinaplon, preferably zolpidem (preferably MR sustained-release version), zaleplon (preferably extended-release formulation), pagoclone, NGD 96-3, indiplon or eszopiclone, or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof. Preferably, zolpidem MR sustained-release version is to be administered in a daily dose of between 10 and 20 mg of the active ingredient. Preferably, zaleplon extended-release is to be administered in a daily dose ranging between 2.5 and 20 mg of the active ingredient. Preferably, pagoclone is to be administered in a daily dose ranging between 7.5 and 60 mg of the active ingredient. Preferably, indiplon is to be administered in a daily dose of between 10 and 20 mg of the active ingredient. Preferably, eszopiclone is to be administered in a daily dose of between 2 and 3 mg of the active

ingredient. Preferably, ocinaplon is to be administered in a daily dose of between 10 and 60 mg of the active ingredient.

The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) a GABA-A modulator, preferably chosen from the group consisting of zolpidem (preferably MR sustained-release version), zaleplon (preferably extended-release formulation), SL 65.1498, SEP174559, pagoclone, NGD 96-3, indiplon, eszopiclone, CP-730,330 (NGD 96-3) and ocinaplon, preferably zolpidem (preferably MR sustained-release version), zaleplon (preferably extended-release formulation), pagoclone, NGD 96-3, indiplon or eszopiclone, or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of a mental disease or disorder which is chosen from the group of diseases or disorders consisting of anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, personality disorders, bereavement, occupational problem and problems related to abuse or neglect.

The invention also relates to a pharmaceutical composition as described above wherein pipamperon is provided in a unitary dose of between 5 and 15 mg of the active ingredient and wherein said GABA-A modulator is zolpidem MR sustained-release version, preferably provided in a unitary dose of between 10 and 20 mg of the active ingredient.

The invention also relates to a pharmaceutical composition as described above wherein pipamperon is provided in a unitary dose of between 5 and 15 mg of the active ingredient and wherein said GABA-A modulator is zaleplon extended-release, preferably provided in a unitary dose of between 2.5 and 20 mg of the active ingredient.

The invention also relates to a pharmaceutical composition as described above wherein pipamperon is provided in a unitary dose of between 5 and 15 mg of the active ingredient and wherein said GABA-A modulator is Pagoclone, preferably provided in a unitary dose of between 7.5 and 60 mg of the active ingredient.

The invention also relates to a pharmaceutical composition as described above wherein pipamperon is provided in a unitary dose of between 5 and 15 mg of the active ingredient

and wherein said GABA-A modulator is indiplon, preferably provided in a unitary dose of between 10 and 20 mg of the active ingredient.

5 The invention also relates to a pharmaceutical composition as described above wherein pipamperon is provided in a unitary dose of between 5 and 15 mg of the active ingredient and wherein said GABA-A modulator is eszopiclone, preferably provided in a unitary dose of between 2 and 3 mg of the active ingredient.

10 The invention also relates to a pharmaceutical composition as described above wherein pipamperon is provided in a unitary dose of between 5 and 15 mg of the active ingredient and wherein said GABA-A modulator is ocinaplon, preferably provided in a unitary dose of between 10 and 60 mg of the active ingredient.

36: combination therapy with a GABA-B antagonist compound

15 The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with a GABA-B (gamma-aminobutyric acid receptor B) antagonist compound, are chosen from the group of diseases or disorders consisting of anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, personality disorders, bereavement, occupational problem and problems related to abuse or neglect.

25 The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting of anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, personality disorders, bereavement, occupational problem and problems related to abuse or neglect, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a GABA-B antagonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said GABA-B antagonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

According to a preferred embodiment, the invention relates to the use as described above, wherein said GABA-B antagonist compound is AVE 7398 or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof.

5 The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) a GABA-B antagonist, preferably AVE 7398 or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of a mental
10 disease or disorder which is chosen from the group of diseases or disorders consisting of anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, personality disorders, bereavement, occupational problem and problems related to abuse or
15 neglect.

37: combination therapy with a Glial-cell Line Derived Neurotrophic Factor compound

The mental disorder which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with a
20 Glial-cell Line Derived Neurotrophic Factor compound, is Parkinson disease.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of Parkinson disease, characterized in that pipamperon or said
25 pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a Glial-cell Line Derived Neurotrophic Factor compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said Glial-cell Line Derived Neurotrophic Factor compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg
30 of the active ingredient.

According to a preferred embodiment, the invention relates to the use as described above, wherein said Glial-cell Line Derived Neurotrophic Factor compound is GDNF or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof. Preferably,
35 GDNF is to be administered in a daily dose ranging between 3.75 and 30 mg of the active ingredient.

The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) a Glial-cell Line Derived Neurotrophic Factor, preferably GDNF or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of Parkinson disease.

The invention also relates to a pharmaceutical composition as described above wherein pipamperon is provided in a unitary dose of between 5 and 15 mg of the active ingredient and wherein said Glial-cell Line Derived Neurotrophic Factor is GDNF, preferably provided in a unitary dose of between 3.75 and 30 mg of the active ingredient.

38: combination therapy with a glucocorticoid synthesis inhibitor compound

The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with a glucocorticoid synthesis inhibitor compound, are chosen from the group of diseases or disorders consisting of substance related disorders and Parkinson disease.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of substance related disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a glucocorticoid synthesis inhibitor compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said glucocorticoid synthesis inhibitor compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of Parkinson disease, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a glucocorticoid synthesis inhibitor compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said glucocorticoid synthesis inhibitor compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

According to a preferred embodiment, the invention relates to the uses as described above, wherein said glucocorticoid synthesis inhibitor compound is metyrapone or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof.

5

The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) a glucocorticoid synthesis inhibitor, preferably metyrapone or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of a mental disease or disorder which is chosen from the group of diseases or disorders consisting of substance related disorders and Parkinson disease.

10

39: combination therapy with a glutamate receptor antagonist compound

15

The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with a glutamate receptor antagonist compound, are chosen from the group of diseases or disorders consisting of anxiety disorders, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, adjustment disorders, impulse control disorders, substance-related disorders, personality disorders, bereavement, occupational problem, problems related to abuse or neglect and pain disorders.

20

25

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting of anxiety disorders, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, adjustment disorders, impulse control disorders, substance-related disorders, personality disorders, bereavement, occupational problem, and problems related to abuse or neglect, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a glutamate receptor antagonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said glutamate receptor antagonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

30

35

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of pain disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a glutamate receptor antagonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said glutamate receptor antagonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

According to a preferred embodiment, the invention relates to the uses as described above, wherein said glutamate receptor antagonist compound is LY354740 or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof.

The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) a glutamate receptor antagonist, preferably LY354740 or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of a mental disease or disorder which is chosen from the group of diseases or disorders consisting of anxiety disorders, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, adjustment disorders, impulse control disorders, substance-related disorders, personality disorders, bereavement, occupational problem, problems related to abuse or neglect and pain disorders.

40: combination therapy with an GPCR (G-protein-coupled receptor) modulator

The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with an GPCR (G-protein-coupled receptor) modulator, are chosen from the group of diseases or disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, substance-related disorders, personality disorders, bereavement, occupational problem, problems related to abuse or neglect and pain disorders.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying

emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, substance-related disorders, personality disorders, bereavement, occupational problem and problems related to abuse or neglect, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a GPCR modulator compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said GPCR modulator compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of pain disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a GPCR modulator compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said GPCR modulator compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

According to a preferred embodiment, the invention relates to the uses as described above, wherein said GPCR modulator compound is R1204 or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof.

The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) a GPCR modulator, preferably R1204 or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of a mental disease or disorder which is chosen from the group of diseases or disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, substance-related disorders, personality disorders, bereavement, occupational problem, problems related to abuse or neglect and pain disorders.

41: combination therapy with an GR (glucocorticoid receptor) antagonist

The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with an GR (glucocorticoid receptor) antagonist, are chosen from the group of diseases or disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, substance-related disorders, personality disorders, bereavement, occupational problem, problems related to abuse or neglect and pain disorders.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, substance-related disorders, personality disorders, bereavement, occupational problem and problems related to abuse or neglect, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a GR antagonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said GR antagonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of pain disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a GR antagonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said GR antagonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

According to a preferred embodiment, the invention relates to the use as described above, wherein said GR antagonist compound is chosen from the group consisting of ORG 34517/34850 and mifepristone, preferably mifepristone, or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof. Preferably, mifepristone is to be administered in a daily dose of between 600 and 1200 mg of the active ingredient.

The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) a GR antagonist, preferably chosen from the group consisting of ORG 34517/34850 and mifepristone, preferably mifepristone, or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of a mental disease or disorder which is chosen from the group of diseases or disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, substance-related disorders, personality disorders, bereavement, occupational problem, problems related to abuse or neglect and pain disorders.

The invention also relates to a pharmaceutical composition as described above wherein pipamperon is provided in a unitary dose of between 5 and 15 mg of the active ingredient and wherein said GR antagonist is Mifepristone, preferably provided in a unitary dose of between 600 and 1200 mg of the active ingredient.

42: combination therapy with a histamine H3-receptor antagonist

The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with a histamine H₃-receptor antagonist, are chosen from the group of diseases or disorders consisting of Alzheimer Disease, substance-related persisting dementia, vascular dementia, dementia due to HIV disease, dementia due to head trauma, dementia due to Parkinson Disease, dementia due to Huntington Disease, dementia due to Pick Disease, dementia due to Creutzfeldt-Jacob Disease, amnesic disorders due to a general medical condition, substance-induced persisting amnesic disorder, mild cognitive impairment disorder and other cognitive disorders.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying

- emotion dysregulation of a cognitive mental disease or disorder selected from the group of diseases and disorders consisting of Alzheimer Disease, substance-related persisting dementia, vascular dementia, dementia due to HIV disease, dementia due to head trauma, dementia due to Parkinson Disease, dementia due to Huntington Disease, dementia due to Pick Disease, dementia due to Creutzfeldt-Jacob Disease, amnestic disorders due to a general medical condition, substance-induced persisting amnestic disorder, mild cognitive impairment disorder and other cognitive disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a histamine H3-receptor antagonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said histamine H3-receptor antagonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.
- According to a preferred embodiment, the invention relates to the uses as described above, wherein said histamine H3-receptor antagonist compound is chosen from the group of compounds consisting of ABT-834 and ABT-239, or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof.
- The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) a histamine H3-receptor antagonist, preferably chosen from the group consisting of ABT-834 and ABT-239 or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of a cognitive mental disease or disorder which is chosen from the group consisting of Alzheimer Disease, substance-related persisting dementia, vascular dementia, dementia due to HIV disease, dementia due to head trauma, dementia due to Parkinson Disease, dementia due to Huntington Disease, dementia due to Pick Disease, dementia due to Creutzfeldt-Jacob Disease, amnestic disorders due to a general medical condition, substance-induced persisting amnestic disorder, mild cognitive impairment disorder and other cognitive disorders.

43: combination therapy with a hormonal substance

- The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with a hormonal substance, are chosen from the group of diseases or disorders consisting of premenstrual syndrome and sexual and gender identity disorders.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a cognitive mental disease or disorder selected from the group of diseases and disorders consisting of premenstrual syndrome and sexual and gender identity disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a hormonal substance to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said hormonal substance, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

According to a preferred embodiment, the invention relates to the uses as described above, wherein said hormonal substance is chosen from the group consisting of a testosterone transdermal spray, a testosterone gel, a female testosterone patch, synthetic conjugated estrogen A, methyltestosterone, a estrogens/methyltestosterone and a drosiperone/ethinyl estradiol composition, or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof. More preferably, said hormonal substance is synthetic conjugated estrogen A and is to be administered in a daily dose ranging between 0.075 and 0.6 mg of the active ingredient. More preferably, said hormonal substance is a drosiperone/ethinyl estradiol composition and is to be administered as a daily dose in tablets, preferably comprising 3mg drosiperone and 0.02 mg ethinyl estradiol of the active ingredients, respectively.

The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) a hormonal substance, preferably chosen from the group consisting of a testosterone transdermal spray, a testosterone gel, a female testosterone patch, synthetic conjugated estrogen A, methyltestosterone, a estrogens/methyltestosterone and a drosiperone/ethinyl estradiol composition, or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of a cognitive mental disease or disorder which is chosen from the group consisting of premenstrual syndrome and sexual and gender identity disorders.

The invention also relates to a pharmaceutical composition as described above wherein pipamperon is provided in a unitary dose of between 5 and 15 mg of the active ingredient

and wherein said hormonal substance is synthetic conjugated estrogen A, preferably provided in a unitary dose of between 0.075 and 0.6 mg of the active ingredient.

The invention also relates to a pharmaceutical composition as described above wherein
5 pipamperon is provided in a unitary dose of between 5 and 15 mg of the active ingredient
and wherein said hormonal substance is a drosiperone/ethinyl estradiol composition,
preferably provided in tablets comprising a unitary dose of 3mg drosiperone and 0.02 mg
ethinyl estradiol of the active ingredients, respectively.

10 44: combination therapy with a compound which increases brain concentrations of 5-HT

The mental disorders which can be treated using compounds having a high selective affinity
for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with a
compound which increases brain concentrations of 5-HT (serotonin), are chosen from the
group of diseases or disorders consisting of mood disorders, anxiety disorders, eating
15 disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders),
factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep
disorders, adjustment disorders, impulse control disorders, substance-related disorders,
personality disorders, bereavement, occupational problem, problems related to abuse or
neglect and pain disorders.

20 The present invention thus relates to the use of pipamperon or a pharmaceutically
acceptable salt thereof for the preparation of a medicament for treating the underlying
emotion dysregulation of a non cognitive mental disease or disorder selected from the group
of diseases and disorders consisting of mood disorders, anxiety disorders, eating disorders,
25 premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious
disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders,
adjustment disorders, impulse control disorders, substance-related disorders, personality
disorders, bereavement, occupational problem and problems related to abuse or neglect,
characterized in that pipamperon or said pharmaceutically acceptable salt thereof is
30 administered simultaneously with, separate from or prior to the administration of a compound
which increases brain concentrations of 5-HT (serotonin) to augment the therapeutic effect
or to provide a faster onset of the therapeutic effect of said compound which increases brain
concentrations of 5-HT (serotonin), further characterized in that pipamperon is to be
administered to a patient in a daily dose ranging between 5 and 15 mg of the active
35 ingredient.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of pain, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a compound which increases brain concentrations of 5-HT (serotonin) to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said compound which increases brain concentrations of 5-HT (serotonin), further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

According to a preferred embodiment, the invention relates to the uses as described above, wherein said compound which increases brain concentrations of 5-HT (serotonin) is chosen from the group consisting of triptosine, tramadol, SP 186, PMD 145 and KW 6055, or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof.

The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) a compound which increases brain concentrations of 5-HT (serotonin), preferably chosen from the group consisting of triptosine, tramadol, SP 186, PMD 145 and KW 6055, or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of a mental disease or disorder which is chosen from the group of diseases and disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, substance-related disorders, personality disorders, bereavement, occupational problem, problems related to abuse or neglect and pain disorders.

45: combination therapy with a compound which increases insulin sensitivity

The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with a compound which increases insulin sensitivity, are chosen from the group of diseases or disorders consisting of Alzheimer Disease, substance-related persisting dementia, vascular dementia, dementia due to HIV disease, dementia due to head trauma, dementia due to Parkinson Disease, dementia due to Huntington Disease, dementia due to Pick Disease, dementia due to Creutzfeldt-Jacob Disease, amnesic disorders due to a general medical

condition, substance-induced persisting amnesic disorder, mild cognitive impairment disorder and other cognitive disorders.

5 The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a cognitive mental disease or disorder selected from the group of diseases and disorders consisting of Alzheimer Disease, substance-related persisting dementia, vascular dementia, dementia due to HIV disease, dementia due to head trauma, dementia due to Parkinson Disease, dementia due to Huntington Disease, dementia due to
10 Pick Disease, dementia due to Creutzfeldt-Jacob Disease, amnesic disorders due to a general medical condition, substance-induced persisting amnesic disorder, mild cognitive impairment disorder and other cognitive disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a compound which increases insulin sensitivity to augment
15 the therapeutic effect or to provide a faster onset of the therapeutic effect of said compound which increases insulin sensitivity, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

20 According to a preferred embodiment, the invention relates to the use as described above, wherein said compound which increases insulin sensitivity is rosiglitazone maleate, or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof.

25 The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) a compound which increases insulin sensitivity, preferably rosiglitazone maleate or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of a cognitive mental disease or disorder which is chosen from the group consisting of Alzheimer Disease, substance-related persisting dementia, vascular dementia, dementia due to HIV disease, dementia due to head trauma, dementia due to Parkinson Disease, dementia due to Huntington Disease, dementia due to Pick
30 Disease, dementia due to Creutzfeldt-Jacob Disease, amnesic disorders due to a general medical condition, substance-induced persisting amnesic disorder, mild cognitive impairment disorder and other cognitive disorders.

46: combination therapy with a compound inhibiting the mixed lineage kinase family

The mental disorder which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with a compound which is an inhibitor of the mixed lineage kinase family is Parkinson Disease.

5

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of Parkinson Disease, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a compound which is an inhibitor of the mixed lineage kinase family to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said compound which is an inhibitor of the mixed lineage kinase family, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

10

According to a preferred embodiment, the invention relates to the use as described above, wherein said compound which is an inhibitor of the mixed lineage kinase family is CEP-1347 or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof.

15

The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) a compound which is an inhibitor of the mixed lineage kinase family, preferably CEP-1347 or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of Parkinson Disease.

20

47: combination therapy with an interleukin-1 beta converting enzyme inhibitor compound

The mental disorder which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with an interleukin-1 beta converting enzyme inhibitor compound, is a pain disorder.

25

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a pain disorder, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of an interleukin-1 beta converting enzyme inhibitor compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said

30

35

interleukin-1 beta converting enzyme inhibitor compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

- 5 According to a preferred embodiment, the invention relates to the use as described above, wherein said interleukin-1 beta converting enzyme inhibitor is pralnacasan or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof.

- 10 The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) an interleukin-1 beta converting enzyme inhibitor, preferably pralnacasan or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of a pain disorder,.

15 48: combination therapy with a levodopa/decarboxylase inhibitor compound

The mental disorder which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with a levodopa/decarboxylase inhibitor compound, is Parkinson Disease.

- 20 The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of Parkinson Disease, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a levodopa/decarboxylase inhibitor compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said
- 25 levodopa/decarboxylase inhibitor compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

- 30 According to a preferred embodiment, the invention relates to the use as described above, wherein said levodopa/decarboxylase inhibitor compound is levodopa/carbidopa, levodopa/benserazide, etilevodopa/carbidopa or etilevodopa/benserazide, or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof. According to a further preferred embodiment, the invention relates to the use as described above, wherein said
- 35 levodopa/decarboxylase inhibitor compound is (eti)levodopa/carbidopa, or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof in combination with

entacapone, which is an inhibitor of catechol-O-methyltransferase (COMT), or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof. Preferably said levodopa/decarboxylase inhibitor compound is levodopa/carbidopa and is to be administered in a dose ranging between 2000 mg/ 50 mg and 100 mg/ 10 mg of the active ingredients.

5 Preferably said entacapone is to be administered in a dose ranging between 1000 mg/ 50 mg, more preferably between 500 mg/ 100 mg, and most preferably 200 mg of the active ingredients per day.

10 The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) a levodopa/decarboxylase inhibitor compound, preferably levodopa/carbidopa, levodopa/benserazide, etilevodopa/carbidopa or etilevodopa/benserazide, or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of Parkinson Disease. The invention also relates to a pharmaceutical
15 composition comprising (a) pipamperon, and (b) a levodopa/decarboxylase inhibitor compound, preferably is (eti)levodopa/carbidopa, or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof in combination with entacapone, which is an inhibitor of catechol-O-methyltransferase (COMT), or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for
20 simultaneous, separate or sequential use for treating the underlying emotion dysregulation of Parkinson Disease.

The invention also relates to a pharmaceutical composition as described above wherein pipamperon is provided in a unitary dose of between 5 and 15 mg of the active ingredient and wherein said levodopa/decarboxylase inhibitor compound is levodopa/carbidopa,
25 preferably provided in a unitary dose of between 100 mg and 10 mg of the active ingredient.

The invention also relates to a pharmaceutical composition as described above wherein pipamperon is provided in a unitary dose of between 5 and 15 mg of the active ingredient and wherein said levodopa/decarboxylase inhibitor compound is levodopa/carbidopa or
30 etilevodopa/ carbidopa in combination with entacapone, of which the latter is preferably provided in a unitary dose of between 500 mg and 100 mg of the active ingredient.

49: combination therapy with a lipid-DNA complex

35 The mental disorder which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with a lipid-DNA complex, is Parkinson Disease.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of Parkinson Disease, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of lipid-DNA complex to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said lipid-DNA complex, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

According to a preferred embodiment, the invention relates to the use as described above, wherein said lipid-DNA complex is GR213487B or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof.

The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) a lipid-DNA complex, preferably GR213487B or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of Parkinson Disease.

50: combination therapy with a monoamine oxidase (MAO) reuptake inhibitor

The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with a monoamine oxidase (MAO) reuptake inhibitor, are chosen from the group of diseases or disorders consisting of substance related disorders and attention-deficit disorders (ADHD).

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of non-cognitive mental disease or disorder which are substance related disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a monoamine oxidase (MAO) reuptake inhibitor compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said monoamine oxidase (MAO) reuptake inhibitor compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of attention-deficit disorders (ADHD), characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a monoamine oxidase (MAO) reuptake inhibitor compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said monoamine oxidase (MAO) reuptake inhibitor compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

According to a preferred embodiment, the invention relates to the uses as described above, wherein said monoamine oxidase (MAO) reuptake inhibitor compound is NS 2359 or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof.

The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) a monoamine oxidase (MAO) reuptake inhibitor, preferably NS 2359 or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of a mental disease or disorder which is chosen from the group consisting of substance related disorders and attention-deficit disorders (ADHD).

51: combination therapy with a MAO-A and a MAO-B reuptake inhibitor

The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with a monoamine oxidase A (MAO-A) and a monoamine oxidase B (MAO-B) reuptake inhibitor, wherein said disorders are attention-deficit disorders.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of attention-deficit disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a monoamine oxidase A (MAO-A) and a monoamine oxidase B (MAO-B) reuptake inhibitor compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said monoamine oxidase A (MAO-A) and a monoamine oxidase B (MAO-B) reuptake inhibitor compound, further characterized in that

pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

According to a preferred embodiment, the invention relates to the uses as described above, wherein said monoamine oxidase A (MAO-A) and a monoamine oxidase B (MAO-B) reuptake inhibitor compound is SPD473 or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof.

The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) a monoamine oxidase A (MAO-A) and a monoamine oxidase B (MAO-B) reuptake inhibitor, preferably SPD473 or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of attention-deficit disorders.

52: combination therapy with a monoamine oxidase B (MAO-B) inhibitor

The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with a monoamine oxidase B (MAO-B) inhibitor, are chosen from the group of diseases or disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, adjustment disorders, impulse control disorders, attention-deficit disorders, substance-related disorders, personality disorders, problems related to abuse or neglect, pain disorder and Parkinson Disease.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, adjustment disorders, impulse control disorders, attention-deficit disorders, substance-related disorders, personality disorders, problems related to abuse or neglect, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a monoamine oxidase B (MAO-B) inhibitor compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said monoamine oxidase B (MAO-B) inhibitor compound, further characterized in that pipamperon is to be

administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

5 The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of pain disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a monoamine oxidase B (MAO-B) inhibitor compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said
10 monoamine oxidase B (MAO-B) inhibitor compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

15 The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of Parkinson Disease, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a monoamine oxidase B (MAO-B) inhibitor compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said
20 monoamine oxidase B (MAO-B) inhibitor compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

25 According to a preferred embodiment, the invention relates to the uses as described above, wherein said monoamine oxidase B (MAO-B) inhibitor compound is chosen from the group consisting of selegiline, rasagiline (TVP-1012) and EmSam (transdermal selegiline), or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof. More preferably, said monoamine oxidase B (MAO-B) inhibitor is selegiline and is to be administered in a daily dose ranging between 5 and 10 mg of the active ingredient. More
30 preferably, said monoamine oxidase B (MAO-B) inhibitor is rasagiline (TVP-1012) and is to be administered in a daily dose ranging between 1 and 2 mg of the active ingredient.

The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) a monoamine oxidase B (MAO-B) inhibitor, preferably chosen from the group consisting
35 of selegiline, rasagiline (TVP-1012) and EmSam (transdermal selegiline), or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined

preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of a mental disease or disorder which is chosen from the group consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, adjustment disorders, impulse control disorders, attention-deficit disorders, substance-related disorders, personality disorders, problems related to abuse or neglect, pain disorder and Parkinson Disease.

The invention also relates to a pharmaceutical composition as described above wherein pipamperon is provided in a unitary dose of between 5 and 15 mg of the active ingredient and wherein said monoamine oxidase B (MAO-B) inhibitor is selegiline, preferably provided in a unitary dose of between 5 and 10 mg of the active ingredient.

The invention also relates to a pharmaceutical composition as described above wherein pipamperon is provided in a unitary dose of between 5 and 15 mg of the active ingredient and wherein said monoamine oxidase B (MAO-B) inhibitor is rasagiline (TVP-1012), preferably provided in a unitary dose of between 1 and 2 mg of the active ingredient.

53: combination therapy with a monoamine oxidase B (MAO-B) reuptake inhibitor

The mental disorder which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with a monoamine oxidase B (MAO-B) reuptake inhibitor, is Parkinson Disease.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of Parkinson Disease, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a monoamine oxidase B (MAO-B) reuptake inhibitor to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said monoamine oxidase B (MAO-B) reuptake inhibitor, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

According to a preferred embodiment, the invention relates to the use as described above, wherein said monoamine oxidase B (MAO-B) reuptake inhibitor is safinamide or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof.

The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) a monoamine oxidase B (MAO-B) reuptake inhibitor, preferably safinamide or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of Parkinson Disease.

54: combination therapy with a melanocortin-4 (MC4) receptor antagonist compound

The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with a melanocortin-4 (MC4) receptor antagonist compound, are chosen from the group of diseases or disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, substance-related disorders, personality disorders, bereavement, occupational problem, problems related to abuse or neglect and pain disorders.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, substance-related disorders, personality disorders, bereavement, occupational problem and problems related to abuse or neglect, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a melanocortin-4 (MC4) receptor antagonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said melanocortin-4 (MC4) receptor antagonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of pain disorders, characterized in that pipamperon or said

pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a melanocortin-4 (MC4) receptor antagonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said melanocortin-4 (MC4) receptor antagonist compound, further characterized in that
5 pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

According to a preferred embodiment, the invention relates to the uses as described above, wherein said melanocortin-4 (MC4) receptor antagonist compound is MCL0129, or a pro-
10 drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof.

The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) a melanocortin-4 (MC4) receptor antagonist compound, preferably MCL0129 or a pro-
15 drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of a mental disease or disorder which is chosen from the group consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders,
20 impulse control disorders, substance-related disorders, personality disorders, bereavement, occupational problem, problems related to abuse or neglect and pain disorders.

55: combination therapy with a MCH receptor antagonist compound

The mental disorders which can be treated using compounds having a high selective affinity
25 for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with a melanin concentrating hormone (MCH) receptor antagonist compound, are chosen from the group of diseases or disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep
30 disorders, adjustment disorders, impulse control disorders, substance-related disorders, personality disorders, bereavement, occupational problem, problems related to abuse or neglect and pain disorders.

The present invention thus relates to the use of pipamperon or a pharmaceutically
35 acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the

group of diseases and disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, substance-related disorders, personality disorders, bereavement, occupational problem and problems related to abuse or neglect, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a melanin concentrating hormone (MCH) receptor antagonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said melanin concentrating hormone (MCH) receptor antagonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of pain disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a melanin concentrating hormone (MCH) receptor antagonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said melanin concentrating hormone (MCH) receptor antagonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

According to a preferred embodiment, the invention relates to the uses as described above, wherein said melanin concentrating hormone (MCH) receptor antagonist compound is SNAP-7941 or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof.

The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) a melanin concentrating hormone (MCH) receptor antagonist compound, preferably SNAP-7941 or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of a mental disease or disorder which is chosen from the group consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders,

adjustment disorders, impulse control disorders, substance-related disorders, personality disorders, bereavement, occupational problem, problems related to abuse or neglect and pain disorders.

5 56: combination therapy with a melatonin receptor (MT) agonist compound

10 The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with a melatonin receptor (MT) agonist compound, are chosen from the group of diseases or disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, substance-related disorders, personality disorders, bereavement, occupational problem, problems related to abuse or neglect and pain disorders.

15 The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders),
20 factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, substance-related disorders, personality disorders, bereavement, occupational problem and problems related to abuse or neglect, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a melatonin
25 receptor (MT) agonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said melatonin receptor (MT) agonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

30 The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of pain disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a melatonin receptor (MT) agonist compound to augment the
35 therapeutic effect or to provide a faster onset of the therapeutic effect of said melatonin receptor (MT) agonist compound, further characterized in that pipamperon is to be

administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

According to a preferred embodiment, the invention relates to the uses as described above, wherein said melatonin receptor (MT) agonist compound is chosen from the group consisting of ramelteon and agomelatine, or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof. More preferably, said melatonin receptor (MT) agonist compound is agomelatine and is to be administered in a daily dose ranging between 25 and 50 mg of the active ingredient.

The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) a melatonin receptor (MT) agonist compound, preferably ramelteon or agomelatine or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of a mental disease or disorder which is chosen from the group consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, substance-related disorders, personality disorders, bereavement, occupational problem, problems related to abuse or neglect and pain disorders.

The invention also relates to a pharmaceutical composition as described above wherein pipamperon is provided in a unitary dose of between 5 and 15 mg of the active ingredient and wherein said melatonin receptor (MT) agonist compound is agomelatine, preferably provided in a unitary dose of between 25 and 50 mg of the active ingredient.

57: combination therapy with a metabotropic glutamate receptor (MgluR) agonist compound

The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with a metabotropic glutamate receptor (MgluR) agonist compound, are chosen from the group of diseases or disorders consisting of anxiety disorders, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, adjustment disorders, impulse control disorders, substance-related disorders, personality disorders, bereavement, occupational problem, problems related to abuse or neglect and pain disorders.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting of anxiety disorders, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, adjustment disorders, impulse control disorders, substance-related disorders, personality disorders, bereavement, occupational problem and problems related to abuse or neglect, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a metabotropic glutamate receptor (MgluR) agonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said metabotropic glutamate receptor (MgluR) agonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of pain disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a metabotropic glutamate receptor (MgluR) agonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said metabotropic glutamate receptor (MgluR) agonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

According to a preferred embodiment, the invention relates to the uses as described above, wherein said metabotropic glutamate receptor (MgluR) agonist compound is PRE703 or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof.

The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) a metabotropic glutamate receptor (MgluR) agonist, preferably PRE703 or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of a mental disease or disorder which is chosen from the group consisting of anxiety disorders, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, adjustment disorders, impulse control disorders, substance-related

disorders, personality disorders, bereavement, occupational problem, problems related to abuse or neglect and pain disorders.

58: combination therapy with a compound mimicking the effect of nerve growth factor (NGF)

5 The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with a compound which mimics the effect of nerve growth factor (NGF), are chosen from the group of diseases or disorders consisting of Alzheimer Disease, substance-related persisting dementia, vascular dementia, dementia due to HIV disease, dementia due to head trauma, 10 dementia due to Parkinson Disease, dementia due to Huntington Disease, dementia due to Pick Disease, dementia due to Creutzfeldt-Jacob Disease, amnestic disorders due to a general medical condition, substance-induced persisting amnestic disorder, mild cognitive impairment disorder, other cognitive disorders and Parkinson Disease.

15 The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a cognitive mental disease or disorder selected from the group of diseases and disorders consisting of Alzheimer Disease, substance-related persisting dementia, vascular dementia, dementia due to HIV disease, dementia due to head trauma, 20 dementia due to Parkinson Disease, dementia due to Huntington Disease, dementia due to Pick Disease, dementia due to Creutzfeldt-Jacob Disease, amnestic disorders due to a general medical condition, substance-induced persisting amnestic disorder, mild cognitive impairment disorder and other cognitive disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from 25 or prior to the administration of a compound which mimics the effect of nerve growth factor (NGF) to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said compound which mimics the effect of nerve growth factor (NGF), further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

30 The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of Parkinson Disease, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from 35 or prior to the administration of a compound which mimics the effect of nerve growth factor (NGF) to augment the therapeutic effect or to provide a faster onset of the therapeutic effect

of said compound which mimics the effect of nerve growth factor (NGF), further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

5 According to a preferred embodiment, the invention relates to the uses as described above, wherein said compound which mimics the effect of nerve growth factor (NGF) is xaliproden or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof. More preferably, said compound which mimics the effect of nerve growth factor (NGF) is xaliproden and is to be administered in a daily dose ranging between 1 and 2 mg of the
10 active ingredient.

The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) a compound which mimics the effect of nerve growth factor (NGF), preferably xaliproden or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof,
15 as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of a mental disease or disorder which is chosen from the group consisting of Alzheimer Disease, substance-related persisting dementia, vascular dementia, dementia due to HIV disease, dementia due to head trauma, dementia due to Parkinson Disease, dementia due to Huntington Disease, dementia due to Pick Disease,
20 dementia due to Creutzfeldt-Jacob Disease, amnesic disorders due to a general medical condition, substance-induced persisting amnesic disorder, mild cognitive impairment disorder, other cognitive disorders and Parkinson Disease.

The invention also relates to a pharmaceutical composition as described above wherein
25 pipamperon is provided in a unitary dose of between 5 and 15 mg of the active ingredient and wherein said compound which mimics the effect of nerve growth factor (NGF) is xaliproden, preferably provided in a unitary dose of between 1 and 2 mg of the active ingredient.

30 59: combination therapy with a muscarinic receptor partial agonist compound

The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with a muscarinic receptor partial agonist compound, are chosen from the group of diseases or disorders consisting of Alzheimer Disease, substance-related persisting dementia, vascular
35 dementia, dementia due to HIV disease, dementia due to head trauma, dementia due to Parkinson Disease, dementia due to Huntington Disease, dementia due to Pick Disease,

dementia due to Creutzfeldt-Jacob Disease, amnestic disorders due to a general medical condition, substance-induced persisting amnestic disorder, mild cognitive impairment disorder and other cognitive disorders.

5 The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a cognitive mental disease or disorder selected from the group of diseases and disorders consisting of Alzheimer Disease, substance-related persisting dementia, vascular dementia, dementia due to HIV disease, dementia due to head trauma, 10 dementia due to Parkinson Disease, dementia due to Huntington Disease, dementia due to Pick Disease, dementia due to Creutzfeldt-Jacob Disease, amnestic disorders due to a general medical condition, substance-induced persisting amnestic disorder, mild cognitive impairment disorder and other cognitive disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from 15 or prior to the administration of a muscarinic receptor partial agonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said muscarinic receptor partial agonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

20 According to a preferred embodiment, the invention relates to the use as described above, wherein said muscarinic receptor partial agonist compound is sevimeline or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof.

25 The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) a muscarinic receptor partial agonist compound, preferably sevimeline or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of a mental disease or disorder which is chosen from the group consisting of 30 Alzheimer Disease, substance-related persisting dementia, vascular dementia, dementia due to HIV disease, dementia due to head trauma, dementia due to Parkinson Disease, dementia due to Huntington Disease, dementia due to Pick Disease, dementia due to Creutzfeldt-Jacob Disease, amnestic disorders due to a general medical condition, substance-induced persisting amnestic disorder, mild cognitive impairment disorder and 35 other cognitive disorders.

60: combination therapy with a selective nor-adrenaline re-uptake inhibitor (NARI) compound

The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with a selective nor-adrenaline re-uptake inhibitor (NARI) compound, are chosen from the group of diseases or disorders consisting of mood disorders, anxiety disorders, adjustment disorders, attention-deficit disorders, personality disorders, antisocial behaviour, bereavement, occupational problem, problems related to abuse or neglect and pain disorders.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting of mood disorders, anxiety disorders, adjustment disorders, attention-deficit disorders, personality disorders, antisocial behaviour, bereavement, occupational problem and problems related to abuse or neglect, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a selective nor-adrenaline receptor inhibitor (NARI) compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said selective nor-adrenaline re-uptake inhibitor (NARI) compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of pain disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a selective nor-adrenaline re-uptake inhibitor (NARI) compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said selective nor-adrenaline re-uptake inhibitor (NARI) compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

According to a preferred embodiment, the invention relates to the uses as described above, wherein said selective nor-adrenaline re-uptake inhibitor (NARI) compound is chosen from the group consisting of reboxetine, atomoxetine hydrochloride, A 75200, 155U88, (S)-A 75200, tandamine, pirandamine, ciclazindol, fluparoxan, lortalamine, talsupram, talopram, prindamine, nomifensine, viloxazine, tomoxetine, duloxetine, venlafaxine and milnacipran or

a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof. More preferably, said selective nor-adrenaline re-uptake inhibitor (NARI) compound is reboxetine and is to be administered in a daily dose ranging between 8 and 12 mg of the active ingredient. More preferably, said selective nor-adrenaline re-uptake inhibitor (NARI) compound is atomoxetine hydrochloride and is to be administered in a daily dose ranging between 40 and 100 mg of the active ingredient.

The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) a selective nor-adrenaline re-uptake inhibitor (NARI) compound, preferably chosen from the group consisting of reboxetine, atomoxetine hydrochloride, A 75200, 155U88, (S)-A 75200, tandamine, pirandamine, ciclazindol, fluparoxan, lortalamine, talsupram, talopram, prindamine, nomifensine, viloxazine, tomoxetine, duloxetine, venlafaxine and milnacipran, or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of mental disease or disorder which is chosen from the group consisting of mood disorders, anxiety disorders, adjustment disorders, attention-deficit disorders, personality disorders, antisocial behaviour, bereavement, occupational problem, problems related to abuse or neglect and pain disorders.

The invention also relates to a pharmaceutical composition as described above wherein pipamperon is provided in a unitary dose of between 5 and 15 mg of the active ingredient and wherein said selective nor-adrenaline re-uptake inhibitor (NARI) compound is reboxetine, preferably provided in a unitary dose of between 8 and 12 mg of the active ingredient.

The invention also relates to a pharmaceutical composition as described above wherein pipamperon is provided in a unitary dose of between 5 and 15 mg of the active ingredient and wherein said selective nor-adrenaline re-uptake inhibitor (NARI) compound is atomoxetine hydrochloride, preferably provided in a unitary dose of between 40 and 100 mg of the active ingredient.

61: combination therapy with a NaSSA compound

The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with a noradrenergic/specific serotonergic antidepressant (NaSSA) compound, are chosen from the group of diseases or disorders consisting of mood disorders, anxiety disorders, eating

disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders),
factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep
disorders, adjustment disorders, impulse control disorders, personality disorders, antisocial
behaviour, bereavement, occupational problem, problems related to abuse or neglect and
5 pain disorders.

The present invention thus relates to the use of pipamperon or a pharmaceutically
acceptable salt thereof for the preparation of a medicament for treating the underlying
emotion dysregulation of a non-cognitive mental disease or disorder selected from the
10 group of diseases and disorders consisting of mood disorders, anxiety disorders, eating
disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders),
factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep
disorders, adjustment disorders, impulse control disorders, personality disorders, antisocial
behaviour, bereavement, occupational problem and problems related to abuse or neglect,
15 characterized in that pipamperon or said pharmaceutically acceptable salt thereof is
administered simultaneously with, separate from or prior to the administration of a
noradrenergic/specific serotonergic antidepressant (NaSSA) compound to augment the
therapeutic effect or to provide a faster onset of the therapeutic effect of said
noradrenergic/specific serotonergic antidepressant (NaSSA) compound, further
20 characterized in that pipamperon is to be administered to a patient in a daily dose ranging
between 5 and 15 mg of the active ingredient.

The present invention thus relates to the use of pipamperon or a pharmaceutically
acceptable salt thereof for the preparation of a medicament for treating the underlying
25 emotion dysregulation of pain disorders, characterized in that pipamperon or said
pharmaceutically acceptable salt thereof is administered simultaneously with, separate from
or prior to the administration of a noradrenergic/specific serotonergic antidepressant
(NaSSA) compound to augment the therapeutic effect or to provide a faster onset of the
therapeutic effect of said noradrenergic/specific serotonergic antidepressant (NaSSA)
30 compound, further characterized in that pipamperon is to be administered to a patient in a
daily dose ranging between 5 and 15 mg of the active ingredient.

According to a preferred embodiment, the invention relates to the uses as described above,
wherein said noradrenergic/specific serotonergic antidepressant (NaSSA) compound is ORG
35 4420 or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt
thereof.

The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) a noradrenergic/specific serotonergic antidepressant (NaSSA) compound, preferably ORG 4420 or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of mental disease or disorder which is chosen from the group consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, personality disorders, antisocial behaviour, bereavement, occupational problem, problems related to abuse or neglect and pain disorders.

62: combination therapy with a selective NDRI compound

The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with a selective nor-adrenaline and dopamine re-uptake inhibitor (NDRI) compound, are chosen from the group of diseases or disorders consisting of mood disorders, anxiety disorders, adjustment disorders, attention-deficit disorders, personality disorders, antisocial behaviour, bereavement, occupational problem, problems related to abuse or neglect, pain disorders, delirium, Alzheimer Disease, substance-related persisting dementia, vascular dementia, dementia due to HIV disease, dementia due to head trauma, dementia due to Parkinson Disease, dementia due to Huntington Disease, dementia due to Pick Disease, dementia due to Creutzfeldt-Jacob Disease, amnesic disorders due to a general medical condition, substance-induced persisting amnesic disorder, mild cognitive impairment disorder and other cognitive disorders.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting of mood disorders, anxiety disorders, adjustment disorders, attention-deficit disorders, personality disorders, antisocial behaviour, bereavement, occupational problem and problems related to abuse or neglect, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a selective nor-adrenaline and dopamine re-uptake inhibitor (NDRI) compound to augment the therapeutic effect or to

provide a faster onset of the therapeutic effect of said selective nor-adrenaline and dopamine re-uptake inhibitor (NDRI) compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

5

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of pain disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a selective nor-adrenaline and dopamine re-uptake inhibitor (NDRI) compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said selective nor-adrenaline and dopamine re-uptake inhibitor (NDRI) compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

10

15

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a cognitive mental disease or disorder selected from the group of diseases and disorders consisting of delirium, Alzheimer Disease, substance-related persisting dementia, vascular dementia, dementia due to HIV disease, dementia due to head trauma, dementia due to Parkinson Disease, dementia due to Huntington Disease, dementia due to Pick Disease, dementia due to Creutzfeldt-Jacob Disease, amnesic disorders due to a general medical condition, substance-induced persisting amnesic disorder, mild cognitive impairment disorder and other cognitive disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a selective nor-adrenaline and dopamine re-uptake inhibitor (NDRI) compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said selective nor-adrenaline and dopamine re-uptake inhibitor (NDRI) compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

20

25

30

According to a preferred embodiment, the invention relates to the uses as described above, wherein said selective nor-adrenaline and dopamine re-uptake inhibitor (NDRI) compound is GW353162 or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof. More preferably, said selective nor-adrenaline and dopamine re-uptake inhibitor

35

(NDRI) compound is GW353162 and is to be administered in a daily dose ranging between 20 and 60 mg of the active ingredient.

5 The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and
(b) a selective nor-adrenaline and dopamine re-uptake inhibitor (NDRI) compound,
preferably GW353162 or a pro-drug or an active metabolite thereof, or a pharmaceutically
acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential
10 use for treating the underlying emotion dysregulation of mental disease or disorder which is
chosen from the group consisting of mood disorders, anxiety disorders, adjustment
disorders, attention-deficit disorders, personality disorders, antisocial behaviour,
bereavement, occupational problem, problems related to abuse or neglect, pain disorders,
delirium, Alzheimer Disease, substance-related persisting dementia, vascular dementia,
15 dementia due to HIV disease, dementia due to head trauma, dementia due to Parkinson
Disease, dementia due to Huntington Disease, dementia due to Pick Disease, dementia due
to Creutzfeldt-Jacob Disease, amnesic disorders due to a general medical condition,
substance-induced persisting amnesic disorder, mild cognitive impairment disorder and
other cognitive disorders.

20 The invention also relates to a pharmaceutical composition as described above wherein
pipamperon is provided in a unitary dose of between 5 and 15 mg of the active ingredient
and wherein said selective nor-adrenaline and dopamine re-uptake inhibitor (NDRI)
compound is GW353162, preferably provided in a unitary dose of between 20 and 60 mg of
the active ingredient.

25 63: combination therapy with a compound which is a neuroimmunophilin ligand

The mental disorder which can be treated using compounds having a high selective affinity
for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with a
compound which is a neuroimmunophilin ligand, is Parkinson Disease.

30 The present invention thus relates to the use of pipamperon or a pharmaceutically
acceptable salt thereof for the preparation of a medicament for treating the underlying
emotion dysregulation of Parkinson Disease, characterized in that pipamperon or said
pharmaceutically acceptable salt thereof is administered simultaneously with, separate from
or prior to the administration of a compound which is a neuroimmunophilin ligand to augment
35 the therapeutic effect or to provide a faster onset of the therapeutic effect of said compound
which is a neuroimmunophilin ligand, further characterized in that pipamperon is to be

administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

According to a preferred embodiment, the invention relates to the use as described above, wherein said a compound which is a neuroimmunophilin ligand is GPI 1485 or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof. More preferably, said a compound which is a neuroimmunophilin ligand is GPI 1485 and is to be administered in a daily dose ranging between 200 and 1000 mg of the active ingredient.

The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) a compound which is a neuroimmunophilin ligand, preferably GPI 1485 or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of Parkinson Disease.

The invention also relates to a pharmaceutical composition as described above wherein pipamperon is provided in a unitary dose of between 5 and 15 mg of the active ingredient and wherein said a compound which is a neuroimmunophilin ligand is GPI 1485, preferably provided in a unitary dose of between 200 and 1000 mg of the active ingredient.

64: combination therapy with a neuromodulator compound

The mental disorder which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with a neuromodulator compound, is Parkinson Disease.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of Parkinson Disease, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a neuromodulator compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said neuromodulator compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

According to a preferred embodiment, the invention relates to the use as described above, wherein said neuromodulator compound is adenosine or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof.

- 5 The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) a neuromodulator compound, preferably adenosine or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of Parkinson Disease.

10

65: combination therapy with a neurotensin receptor antagonist compound

- The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with a neurotensin receptor antagonist compound, are chosen from the group of diseases or disorders consisting of mood disorders, anxiety disorders, psychotic disorders, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sleep disorders, adjustment disorders, impulse control disorders, pervasive development disorders, disruptive behaviour disorders, substance-related disorders, personality disorders, psychological factors affecting medical conditions, malingering, antisocial behaviour, bereavement, occupational problem, identity problem, problems related to abuse or neglect, pain disorders and delirium.

- The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting of mood disorders, anxiety disorders, psychotic disorders, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sleep disorders, adjustment disorders, impulse control disorders, pervasive development disorders, disruptive behaviour disorders, substance-related disorders, personality disorders, psychological factors affecting medical conditions, malingering, antisocial behaviour, bereavement, occupational problem, identity problem and problems related to abuse or neglect, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a neurotensin receptor antagonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said neurotensin receptor

antagonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

5 The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of pain disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a neurotensin receptor antagonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said neurotensin receptor antagonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

15 The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a cognitive mental disease or disorder which is delirium, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a neurotensin receptor antagonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said neurotensin receptor antagonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

25 According to a preferred embodiment, the invention relates to the use as described above, wherein said neurotensin receptor antagonist compound is SR 48692 or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof. More preferably, said neurotensin receptor antagonist compound is SR 48692 and is to be administered in a daily dose ranging between 90 and 300 mg of the active ingredient.

30 The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) a neurotensin receptor antagonist compound, preferably SR 48692 or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of a mental disease or disorder which is chosen from the group consisting mood disorders, anxiety disorders, psychotic disorders, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sleep disorders, adjustment

disorders, impulse control disorders, pervasive development disorders, disruptive behaviour disorders, substance-related disorders, personality disorders, psychological factors affecting medical conditions, malingering, antisocial behaviour, bereavement, occupational problem, identity problem, problems related to abuse or neglect, pain disorders and delirium.

5

The invention also relates to a pharmaceutical composition as described above wherein pipamperon is provided in a unitary dose of between 5 and 15 mg of the active ingredient and wherein said neurotensin receptor antagonist compound is SR 48692, preferably provided in a unitary dose of between 90 and 300 mg of the active ingredient.

10

66: combination therapy with nerve growth factor (NGF) gene therapy

15

The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with nerve growth factor (NGF) gene therapy, are chosen from the group of diseases or disorders consisting of Alzheimer Disease, substance-related persisting dementia, vascular dementia, dementia due to HIV disease, dementia due to head trauma, dementia due to Parkinson Disease, dementia due to Huntington Disease, dementia due to Pick Disease, dementia due to Creutzfeldt-Jacob Disease, amnesic disorders due to a general medical condition, substance-induced persisting amnesic disorder, mild cognitive impairment disorder, other cognitive disorders and Parkinson Disease.

20

25

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a cognitive mental disease or disorder selected from the group of diseases and disorders consisting of Alzheimer Disease, substance-related persisting dementia, vascular dementia, dementia due to HIV disease, dementia due to head trauma, dementia due to Parkinson Disease, dementia due to Huntington Disease, dementia due to Pick Disease, dementia due to Creutzfeldt-Jacob Disease, amnesic disorders due to a general medical condition, substance-induced persisting amnesic disorder, mild cognitive impairment disorder and other cognitive disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from nerve growth factor (NGF) gene therapy, to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said nerve growth factor (NGF) gene therapy, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

35

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of Parkinson Disease, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from
5 or prior to nerve growth factor (NGF) gene therapy, to augment the therapeutic effect or to provide a faster onset of nerve growth factor (NGF) gene therapy, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

10 The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) a compound useful in nerve growth factor (NGF) gene therapy, preferably xaliproden or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of a mental disease or disorder which is chosen from the
15 group consisting of Alzheimer Disease, substance-related persisting dementia, vascular dementia, dementia due to HIV disease, dementia due to head trauma, dementia due to Parkinson Disease, dementia due to Huntington Disease, dementia due to Pick Disease, dementia due to Creutzfeldt-Jacob Disease, amnesic disorders due to a general medical condition, substance-induced persisting amnesic disorder, mild cognitive impairment
20 disorder, other cognitive disorders and Parkinson Disease.

It should be understood that "nerve growth factor gene therapy" is well known in the art, and the compounds, for instance nucleic acids used in nerve growth factor gene therapy are well described (see e.g. Tuszynski *et al.*, (2002) Journal of Molecular Neuroscience
25 Volume 19, Issue 1-2, pps. 207-208).

67: combination therapy with a nicotinic acetylcholine receptor antagonist compound

The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with a
30 nicotinic acetylcholine receptor antagonist compound, are chosen from the group of diseases or disorders consisting of anxiety disorders, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, adjustment disorders, impulse control disorders, substance-related disorders, personality disorders, bereavement, occupational problem, problems related to abuse or neglect and pain disorders.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting of anxiety disorders, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, adjustment disorders, impulse control disorders, substance-related disorders, personality disorders, bereavement, occupational problem and problems related to abuse or neglect, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a nicotinic acetylcholine receptor antagonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said nicotinic acetylcholine receptor antagonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of pain disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a nicotinic acetylcholine receptor antagonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said nicotinic acetylcholine receptor antagonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

According to a preferred embodiment, the invention relates to the uses as described above, wherein said nicotinic acetylcholine receptor antagonist compound is SEP174559 or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof.

The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) a nicotinic acetylcholine receptor antagonist compound, preferably SEP174559 or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of a mental disease or disorder which is chosen from the group consisting of anxiety disorders, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, adjustment disorders, impulse control disorders,

substance-related disorders, personality disorders, bereavement, occupational problem, problems related to abuse or neglect and pain disorders.

68: combination therapy with a nicotinic receptor agonist compound

- 5 The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with a nicotinic receptor agonist compound, are chosen from the group of diseases or disorders consisting of Alzheimer Disease, substance-related persisting dementia, vascular dementia, dementia due to HIV disease, dementia due to head trauma, dementia due to Parkinson
- 10 Disease, dementia due to Huntington Disease, dementia due to Pick Disease, dementia due to Creutzfeldt-Jacob Disease, amnesic disorders due to a general medical condition, substance-induced persisting amnesic disorder, mild cognitive impairment disorder and other cognitive disorders.
- 15 The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a cognitive mental disease or disorder selected from the group of diseases and disorders consisting of Alzheimer Disease, substance-related persisting dementia, vascular dementia, dementia due to HIV disease, dementia due to head trauma,
- 20 dementia due to Parkinson Disease, dementia due to Huntington Disease, dementia due to Pick Disease, dementia due to Creutzfeldt-Jacob Disease, amnesic disorders due to a general medical condition, substance-induced persisting amnesic disorder, mild cognitive impairment disorder and other cognitive disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from
- 25 or prior to the administration of a nicotinic receptor agonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said nicotinic receptor agonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.
- 30 According to a preferred embodiment, the invention relates to the use as described above, wherein said nicotinic receptor agonist compound is ABT-089, or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof. More preferably, said nicotinic receptor agonist compound is ABT-089 and is to be administered in a daily dose ranging between 4 and 40 mg of the active ingredient.

The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) a nicotinic receptor agonist compound, preferably ABT-089 or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of a cognitive mental disease or disorder which is chosen from the group consisting of Alzheimer Disease, substance-related persisting dementia, vascular dementia, dementia due to HIV disease, dementia due to head trauma, dementia due to Parkinson Disease, dementia due to Huntington Disease, dementia due to Pick Disease, dementia due to Creutzfeldt-Jacob Disease, amnesic disorders due to a general medical condition, substance-induced persisting amnesic disorder, mild cognitive impairment disorder and other cognitive disorders.

The invention also relates to a pharmaceutical composition as described above, wherein pipamperon is provided in a unitary dose of between 5 and 15 mg of the active ingredient and wherein said nicotinic receptor agonist compound is ABT-089, preferably provided in a unitary dose of between 4 and 40 mg of the active ingredient.

69: combination therapy with a neurokinin 2 receptor (NK2) antagonist compound

The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with a neurokinin 2 receptor (NK2) antagonist compound, are chosen from the group of diseases or disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, substance-related disorders, personality disorders, bereavement, occupational problem, problems related to abuse or neglect and pain disorders.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, substance-related disorders, personality disorders, bereavement, occupational problem and problems related to abuse or neglect, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is

115

administered simultaneously with, separate from or prior to the administration of a neurokinin 2 receptor (NK2) antagonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said neurokinin 2 receptor (NK2) antagonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of pain disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a neurokinin 2 receptor (NK2) antagonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said neurokinin 2 receptor (NK2) antagonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

According to a preferred embodiment, the invention relates to the uses as described above, wherein said neurokinin 2 receptor (NK2) antagonist compound is saredutant or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof. More preferably, said neurokinin 2 receptor (NK2) antagonist compound is saredutant and is to be administered in a daily dose ranging between 25 and 200 mg of the active ingredient.

The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) a neurokinin 2 receptor (NK2) antagonist compound, preferably saredutant or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of a mental disease or disorder which is chosen from the group consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, substance-related disorders, personality disorders, bereavement, occupational problem, problems related to abuse or neglect and pain disorders.

The invention also relates to a pharmaceutical composition as described above wherein pipamperon is provided in a unitary dose of between 5 and 15 mg of the active ingredient

and wherein said neurokinin 2 receptor (NK2) antagonist compound is saredutant, preferably provided in a unitary dose of between 25 and 200 mg of the active ingredient.

70: combination therapy with a neurokinin 3 receptor (NK3) antagonist compound

5 The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with a neurokinin 3 receptor (NK3) antagonist compound, are chosen from the group of diseases or disorders consisting of mood disorders, anxiety disorders, psychotic disorders, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sleep
10 disorders, adjustment disorders, impulse control disorders, pervasive development disorders, disruptive behaviour disorders, substance-related disorders, personality disorders, psychological factors affecting medical conditions, malingering, antisocial behaviour, bereavement, occupational problem, identity problem, problems related to abuse or neglect, pain disorders and delirium.

15 The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting of mood disorders, anxiety disorders, psychotic disorders, somatoform disorders (excluding pain disorders), factitious disorders, dissociative
20 disorders, sleep disorders, adjustment disorders, impulse control disorders, pervasive development disorders, disruptive behaviour disorders, substance-related disorders, personality disorders, psychological factors affecting medical conditions, malingering, antisocial behaviour, bereavement, occupational problem, identity problem and problems related to abuse or neglect, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the
25 administration of a neurokinin 3 receptor (NK3) antagonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said neurokinin 3 receptor (NK3) antagonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active
30 ingredient.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying
35 emotion dysregulation of pain disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from

or prior to the administration of a neurokinin 3 receptor (NK3) antagonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said neurokinin 3 receptor (NK3) antagonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a cognitive mental disease or disorder which is delirium, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a neurokinin 3 receptor (NK3) antagonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said neurokinin 3 receptor (NK3) antagonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

According to a preferred embodiment, the invention relates to the uses as described above, wherein said neurokinin 3 receptor (NK3) antagonist compound is talnetant or osanetant, or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof. More preferably, said neurokinin 3 receptor (NK3) antagonist compound is talnetant and is to be administered in a daily dose ranging between 1.5 and 12 mg of the active ingredient.

The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) a neurokinin 3 receptor (NK3) antagonist compound, preferably talnetant or osanetant, or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of a mental disease or disorder which is chosen from the group consisting mood disorders, anxiety disorders, psychotic disorders, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sleep disorders, adjustment disorders, impulse control disorders, pervasive development disorders, disruptive behaviour disorders, substance-related disorders, personality disorders, psychological factors affecting medical conditions, malingering, antisocial behaviour, bereavement, occupational problem, identity problem, problems related to abuse or neglect, pain disorders and delirium.

The invention also relates to a pharmaceutical composition as described above wherein pipamperon is provided in a unitary dose of between 5 and 15 mg of the active ingredient and wherein said neurokinin 3 receptor (NK3) antagonist compound is talnetant, preferably provided in a unitary dose of between 1.5 and 12 mg of the active ingredient.

5

71: combination therapy with an N-Methyl-D-aspartate (NMDA) antagonist compound

The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with an N-Methyl-D-aspartate (NMDA) antagonist compound, are chosen from the group of diseases or disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, adjustment disorders, impulse control disorders, personality disorders, bereavement, occupational problem, problems related to abuse or neglect, pain disorders, Alzheimer Disease, substance-related persisting dementia, vascular dementia, dementia due to HIV disease, dementia due to head trauma, dementia due to Parkinson Disease, dementia due to Huntington Disease, dementia due to Pick Disease, dementia due to Creutzfeldt-Jacob Disease, amnesic disorders due to a general medical condition, substance-induced persisting amnesic disorder, mild cognitive impairment disorder and other cognitive disorders.

20

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, adjustment disorders, impulse control disorders, personality disorders, bereavement, occupational problem and problems related to abuse or neglect, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of an N-Methyl-D-aspartate (NMDA) antagonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said N-Methyl-D-aspartate (NMDA) antagonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

35

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of pain disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from
5 or prior to the administration of an N-Methyl-D-aspartate (NMDA) antagonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said N-Methyl-D-aspartate (NMDA) antagonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

10 The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a cognitive mental disease or disorder selected from the group of diseases and disorders consisting of Alzheimer Disease, substance-related persisting
15 dementia, vascular dementia, dementia due to HIV disease, dementia due to head trauma, dementia due to Parkinson Disease, dementia due to Huntington Disease, dementia due to Pick Disease, dementia due to Creutzfeldt-Jacob Disease, amnestic disorders due to a general medical condition, substance-induced persisting amnestic disorder, mild cognitive impairment disorder and other cognitive disorders, characterized in that pipamperon or said
20 pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of an N-Methyl-D-aspartate (NMDA) antagonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said N-Methyl-D-aspartate (NMDA) antagonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active
25 ingredient.

According to a preferred embodiment, the invention relates to the uses as described above, wherein said N-Methyl-D-aspartate (NMDA) antagonist compound is chosen from the group consisting of SEP174559, memantine, delucemine, or a pro-drug or an active metabolite
30 thereof, or a pharmaceutically acceptable salt thereof. More preferably, said N-Methyl-D-aspartate (NMDA) antagonist compound is memantine and is to be administered in a daily dose ranging between 5 and 40 mg of the active ingredient.

The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and
35 (b) an N-Methyl-D-aspartate (NMDA) antagonist compound, preferably chosen from the group consisting of SEP174559, memantine, delucemine or a pro-drug or an active

metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of mental disease or disorder which is chosen from the group consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, adjustment disorders, impulse control disorders, personality disorders, bereavement, occupational problem, problems related to abuse or neglect, pain disorders, Alzheimer Disease, substance-related persisting dementia, vascular dementia, dementia due to HIV disease, dementia due to head trauma, dementia due to Parkinson Disease, dementia due to Huntington Disease, dementia due to Pick Disease, dementia due to Creutzfeldt-Jacob Disease, amnesic disorders due to a general medical condition, substance-induced persisting amnesic disorder, mild cognitive impairment disorder and other cognitive disorders.

The invention also relates to a pharmaceutical composition as described above wherein pipamperon is provided in a unitary dose of between 5 and 15 mg of the active ingredient and wherein said N-Methyl-D-aspartate (NMDA) antagonist compound is memantine, preferably provided in a unitary dose of between 5 and 40 mg of the active ingredient.

72: combination therapy with a non-steroidal anti-inflammatory drug

The mental disorder which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with a non-steroidal anti-inflammatory drug, is a pain disorder or Alzheimer Disease.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a pain disorder, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a non-steroidal anti-inflammatory drug to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said a non-steroidal anti-inflammatory drug, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a cognitive disease, such as Alzheimer Disease, characterized in

that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a non-steroidal anti-inflammatory drug to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said a non-steroidal anti-inflammatory drug, further characterized in that
5 pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

According to a preferred embodiment, the invention relates to the uses as described above, wherein said a non-steroidal anti-inflammatory drug is chosen from the group consisting of
10 piroxicam , MX-1094, meloxicam and flurizan (pure R-enantiomer form of flurbiprofen), or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof.

The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) a non-steroidal anti-inflammatory drug, preferably chosen from the group consisting of
15 piroxicam, MX-1094, meloxicam and flurizan (pure R-enantiomer form of flurbiprofen), or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of a pain disorder or Alzheimer Disease .

20 73: combination therapy with an opoid antagonist compound

The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with an opoid antagonist compound, are substance related disorders.

25 It will be appreciated that the terms "opoid" and "opioid" may be used interchangeably.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of substance related disorders, characterized in that pipamperon or
30 said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a opoid antagonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said opoid antagonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

According to a preferred embodiment, the invention relates to the use as described above, wherein said opioid antagonist compound is naltrexone, preferably as a depot formulation, more preferably in the form of microcapsules, or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof. Preferably, said naltrexone is to be administered in the form of a depot, preferably a depot of microcapsules comprising a daily dose of between 192 and 384 mg.

The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) an opioid antagonist, preferably naltrexone, or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of substance related disorders.

The invention also relates to a pharmaceutical composition as described above wherein pipamperon is provided in a unitary dose of between 5 and 15 mg of the active ingredient and wherein said opioid antagonist compound is naltrexone, preferably provided in a unitary dose of between 192 and 384 mg of the active ingredient.

74: combination therapy with an opioid agonist compound

The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with an opioid agonist compound, are chosen from the group of diseases or disorders consisting of anxiety disorders, psychotic disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, substance-related disorders, personality disorders, bereavement, occupational problem and problems related to abuse or neglect.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting of anxiety disorders, psychotic disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, substance-related disorders, personality disorders, bereavement, occupational problem and problems related to abuse or

123

neglect, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of an opoid agonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said opoid agonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

According to a preferred embodiment, the invention relates to the use as described above, wherein said opoid agonist compound is chosen from the group consisting of siramesine, E-5842 and cyclazocine, preferably siramesine, or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof.

The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) an opoid agonist compound, preferably chosen from the group consisting of siramesine, E-5842 and cyclazocine, preferably siramesine, or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder which is chosen from the group consisting of anxiety disorders, psychotic disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, substance-related disorders, personality disorders, bereavement, occupational problem and problems related to abuse or neglect.

75: combination therapy with a phosphodiesterase-4 (PDE4) inhibitor compound

The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with a phosphodiesterase-4 (PDE4) inhibitor compound, are chosen from the group of diseases or disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, adjustment disorders, impulse control disorders, personality disorders, bereavement, occupational problem, problems related to abuse or neglect, pain disorders, Alzheimer Disease, substance-related persisting dementia, vascular dementia, dementia due to HIV disease, dementia due to head trauma, dementia due to Parkinson Disease, dementia due to Huntington Disease, dementia due to Pick Disease, dementia due to Creutzfeldt-Jacob Disease, amnesic disorders due to a general

medical condition, substance-induced persisting amnesic disorder, mild cognitive impairment disorder and other cognitive disorders.

5 The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, adjustment disorders, impulse control disorders, personality disorders, bereavement, occupational problem and problems related to abuse or neglect, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of phosphodiesterase-4 (PDE4) inhibitor compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said phosphodiesterase-4 (PDE4) inhibitor compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

20 The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of pain disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a phosphodiesterase-4 (PDE4) inhibitor compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said phosphodiesterase-4 (PDE4) inhibitor compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

30 The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a cognitive mental disease or disorder selected from the group of diseases and disorders consisting of Alzheimer Disease, substance-related persisting dementia, vascular dementia, dementia due to HIV disease, dementia due to head trauma, dementia due to Parkinson Disease, dementia due to Huntington Disease, dementia due to Pick Disease, dementia due to Creutzfeldt-Jacob Disease, amnesic disorders due to a general medical condition, substance-induced persisting amnesic disorder, mild cognitive

impairment disorder and other cognitive disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a phosphodiesterase-4 (PDE4) inhibitor compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said phosphodiesterase-4 (PDE4) inhibitor compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

According to a preferred embodiment, the invention relates to the uses as described above, wherein said phosphodiesterase-4 (PDE4) inhibitor compound is chosen from the group consisting of ND1251 and MEM 1917 (R1497), or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof.

The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) a phosphodiesterase-4 (PDE4) inhibitor antagonist compound, preferably chosen from the group consisting of ND1251 and MEM 1917 (R1497), or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of mental disease or disorder which is chosen from the group consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, adjustment disorders, impulse control disorders, personality disorders, bereavement, occupational problem, problems related to abuse or neglect, pain disorders, Alzheimer Disease, substance-related persisting dementia, vascular dementia, dementia due to HIV disease, dementia due to head trauma, dementia due to Parkinson Disease, dementia due to Huntington Disease, dementia due to Pick Disease, dementia due to Creutzfeldt-Jacob Disease, amnesic disorders due to a general medical condition, substance-induced persisting amnesic disorder, mild cognitive impairment disorder and other cognitive disorders.

76: combination therapy with a peptidic compound

The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with a peptidic compound, are chosen from the group of diseases or disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and

gender identity disorders, adjustment disorders, impulse control disorders, personality disorders, bereavement, occupational problem, problems related to abuse or neglect, pain disorders, Alzheimer Disease, substance-related persisting dementia, vascular dementia, dementia due to HIV disease, dementia due to head trauma, dementia due to Parkinson Disease, dementia due to Huntington Disease, dementia due to Pick Disease, dementia due to Creutzfeldt-Jacob Disease, amnesic disorders due to a general medical condition, substance-induced persisting amnesic disorder, mild cognitive impairment disorder and other cognitive disorders.

10 The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders),
15 factitious disorders, dissociative disorders, sexual and gender identity disorders, adjustment disorders, impulse control disorders, personality disorders, bereavement, occupational problem and problems related to abuse or neglect, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a peptidic compound to augment the therapeutic effect or to
20 provide a faster onset of the therapeutic effect of said peptidic compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of pain disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a peptidic compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said peptidic compound, further
25 characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a cognitive mental disease or disorder selected from the group of diseases and disorders consisting of Alzheimer Disease, substance-related persisting
35

dementia, vascular dementia, dementia due to HIV disease, dementia due to head trauma, dementia due to Parkinson Disease, dementia due to Huntington Disease, dementia due to Pick Disease, dementia due to Creutzfeldt-Jacob Disease, amnesic disorders due to a general medical condition, substance-induced persisting amnesic disorder, mild cognitive impairment disorder and other cognitive disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a peptidic compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said peptidic compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

According to a preferred embodiment, the invention relates to the uses as described above, wherein said peptidic compound is chosen from the group consisting of secretin, PT-141, INN 00835 and beta sheet breaker peptide, or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof. More preferably, said peptidic compound is secretin and is to be administered in a daily dose ranging between 0.2 and 0.4 mg/kg of the active ingredient. More preferably, said peptidic compound is INN 00835 and is to be administered in a daily dose ranging between 18 and 160 mg of the active ingredient.

The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) a peptidic compound, preferably chosen from the group consisting of secretin, PT-141, INN 00835 and beta sheet breaker peptide, or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of mental disease or disorder which is chosen from the group consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, adjustment disorders, impulse control disorders, personality disorders, bereavement, occupational problem, problems related to abuse or neglect, pain disorders, Alzheimer Disease, substance-related persisting dementia, vascular dementia, dementia due to HIV disease, dementia due to head trauma, dementia due to Parkinson Disease, dementia due to Huntington Disease, dementia due to Pick Disease, dementia due to Creutzfeldt-Jacob Disease, amnesic disorders due to a general medical condition, substance-induced persisting amnesic disorder, mild cognitive impairment disorder and other cognitive disorders.

128

The invention also relates to a pharmaceutical composition as described above wherein pipamperon is provided in a unitary dose of between 5 and 15 mg of the active ingredient and wherein said peptidic compound is secretin, preferably provided in a unitary dose of 0.2 and 0.4 mg/kg of the active ingredient.

5

The invention also relates to a pharmaceutical composition as described above wherein pipamperon is provided in a unitary dose of between 5 and 15 mg of the active ingredient and wherein said peptidic compound is INN 00835, preferably provided in a unitary dose of 18 and 160 mg of the active ingredient.

10

77: combination therapy with a phospholipase A2 inhibitor compound

15

The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with a phospholipase A2 inhibitor compound which has caspase inhibitor activity, are chosen from the group of diseases or disorders consisting of mood disorders, anxiety disorders, psychotic disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, pervasive development disorders, disruptive behaviour disorders, substance-related disorders, personality disorders, bereavement, occupational problem, problems related to abuse or neglect, pain disorders and delirium.

20

25

30

35

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting of mood disorders, anxiety disorders, psychotic disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, pervasive development disorders, disruptive behaviour disorders, substance-related disorders, personality disorders, bereavement, occupational problem and problems related to abuse or neglect,, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a phospholipase A2 inhibitor compound which has caspase inhibitor activity to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said phospholipase A2 inhibitor compound which has caspase inhibitor activity, further characterized in that

pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

5 The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of pain disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a phospholipase A2 inhibitor compound which has caspase inhibitor activity to augment the therapeutic effect or to provide a faster onset of the
10 therapeutic effect of said phospholipase A2 inhibitor compound which has caspase inhibitor activity, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

15 The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a cognitive mental disease or disorder which is delirium, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a phospholipase A2 inhibitor compound which has caspase inhibitor activity to augment the
20 therapeutic effect or to provide a faster onset of the therapeutic effect of said phospholipase A2 inhibitor compound which has caspase inhibitor activity, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

25 According to a preferred embodiment, the invention relates to the uses as described above, wherein said phospholipase A2 inhibitor compound which has caspase inhibitor activity is chosen from the group consisting of LAX-101a, LAX-101b and LAX-101c, preferably LAX-101a, or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof.

30 The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) a phospholipase A2 inhibitor compound which has caspase inhibitor activity, preferably chosen from the group consisting of LAX-101a, LAX-101b and LAX-101c, more preferably LAX-101a, or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable
35 salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of a mental disease or disorder which is

chosen from the group consisting mood disorders, anxiety disorders, psychotic disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, pervasive development disorders, disruptive behaviour disorders, substance-related disorders, personality disorders, bereavement, occupational problem, problems related to abuse or neglect, pain disorders and delirium.

78: combination therapy with a compound which is a prodrug of uridine

The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with a compound which is a prodrug of uridine, are chosen from the group of diseases or disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, personality disorders, bereavement, occupational problem, problems related to abuse or neglect and pain disorders.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, substance-related disorders, personality disorders, bereavement, occupational problem and problems related to abuse or neglect, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a compound which is a prodrug of uridine to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said compound which is a prodrug of uridine, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of pain disorders, characterized in that pipamperon or said

pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a compound which is a prodrug of uridine to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said compound which is a prodrug of uridine, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

According to a preferred embodiment, the invention relates to the uses as described above, wherein said compound which is a prodrug of uridine is RG2133 (triacetyluridine) or a prodrug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof.

The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) a compound which is a prodrug of uridine, preferably RG2133 (triacetyluridine) or a prodrug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of a mental disease or disorder which is chosen from the group consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, personality disorders, bereavement, occupational problem, problems related to abuse or neglect and pain disorders.

79: combination therapy with prostaglandin E1 compound

The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with prostaglandin E1 compound, are sexual and gender identity disorders.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of sexual and gender identity disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a prostaglandin E1 compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said prostaglandin E1 compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

According to a preferred embodiment, the invention relates to the use as described above, wherein said prostaglandin E1 is alprostadil or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof. More preferably, said prostaglandin E1 compound is alprostadil, preferably in the form of cream or gel, preferably a topical gel, and is to be administered in a daily dose ranging between 50 and 300 microgram per application of the active ingredient.

The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) a prostaglandin E1 compound, preferably alprostadil or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of a mental disease or disorder which is chosen from the group of sexual and gender identity disorders.

The invention also relates to a pharmaceutical composition as described above wherein pipamperon is provided in a unitary dose of between 5 and 15 mg of the active ingredient and wherein said prostaglandin E1 compound is alprostadil, preferably provided in the form of a cream or gel, preferably a topical gel, wherein a unitary dose comprises between 50 and 300 microgram of the active ingredient per application.

80: combination therapy with a compound protecting dopaminergic and cholinergic neurons

The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with a compound which protects dopaminergic and cholinergic neurons, are chosen from the group of diseases or disorders consisting of Alzheimer Disease, substance-related persisting dementia, vascular dementia, dementia due to HIV disease, dementia due to head trauma, dementia due to Parkinson Disease, dementia due to Huntington Disease, dementia due to Pick Disease, dementia due to Creutzfeldt-Jacob Disease, amnestic disorders due to a general medical condition, substance-induced persisting amnestic disorder, mild cognitive impairment disorder, other cognitive disorders and Parkinson Disease.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a cognitive mental disease or disorder selected from the group of diseases and disorders consisting of Alzheimer Disease, substance-related persisting dementia, vascular dementia, dementia due to HIV disease, dementia due to head trauma,

dementia due to Parkinson Disease, dementia due to Huntington Disease, dementia due to Pick Disease, dementia due to Creutzfeldt-Jacob Disease, amnestic disorders due to a general medical condition, substance-induced persisting amnestic disorder, mild cognitive impairment disorder and other cognitive disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a compound which protects dopaminergic and cholinergic neurons to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said compound which protects dopaminergic and cholinergic neurons, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of Parkinson Disease, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a compound which protects dopaminergic and cholinergic neurons to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said compound which protects dopaminergic and cholinergic neurons, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

According to a preferred embodiment, the invention relates to the uses as described above, wherein said compound which protects dopaminergic and cholinergic neurons is SR 57667 or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof.

The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) a compound which protects dopaminergic and cholinergic neurons, preferably SR 57667 or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of a mental disease or disorder which is chosen from the group consisting of Alzheimer Disease, substance-related persisting dementia, vascular dementia, dementia due to HIV disease, dementia due to head trauma, dementia due to Parkinson Disease, dementia due to Huntington Disease, dementia due to Pick Disease, dementia due to Creutzfeldt-Jacob Disease, amnestic disorders due to a general medical condition, substance-induced persisting amnestic disorder, mild cognitive impairment disorder, other cognitive disorders and Parkinson Disease.

81: combination therapy with a psychostimulant

The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with a psychostimulant, are chosen from the group of diseases or disorders consisting of sleep disorders, attention-deficit disorders and substance-related disorders.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting of sleep disorders, attention-deficit disorders and substance-related disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a psychostimulant to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said psychostimulant, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

According to a preferred embodiment, the invention relates to the uses as described above, wherein said psychostimulant is chosen from the group consisting of SPD 503, r-modafinil and modafinil, or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof. More preferably, said psychostimulant is SPE 503, more preferably said psychostimulant is modafinil and is to be administered in a daily dose ranging between 200 and 600 mg of the active ingredient.

The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) a psychostimulant, preferably chosen from the group consisting of SPD 503, r-modafinil and modafinil, more preferably said SPC 503 or modafinil or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder which is chosen from the group consisting of sleep disorders, attention-deficit disorders and substance-related disorders.

The invention also relates to a pharmaceutical composition as described above wherein pipamperon is provided in a unitary dose of between 5 and 15 mg of the active ingredient

and wherein said psychostimulant is modafinil, preferably provided in a unitary dose of between 200 and 600 mg of the active ingredient.

82: combination therapy with a compound which is a Reversible Inhibitor of Mono-Amine oxydase A (RIMA)

The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with a compound which is a reversible inhibitor of mono-amine oxydase A (RIMA), are chosen from the group of diseases or disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, adjustment disorders, impulse control disorders, personality disorders, antisocial behaviour, bereavement, occupational problem, problems related to abuse or neglect and pain disorders.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, adjustment disorders, impulse control disorders, personality disorders, antisocial behaviour, bereavement, occupational problem and problems related to abuse or neglect, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a compound which is a reversible inhibitor of mono-amine oxydase A (RIMA) to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said compound which is a reversible inhibitor of mono-amine oxydase A (RIMA), further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of pain disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a compound which is a reversible inhibitor of mono-amine

oxydase A (RIMA) to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said compound which is a reversible inhibitor of mono-amine oxydase A (RIMA), further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

5

According to a preferred embodiment, the invention relates to the uses as described above, wherein said compound which is a reversible inhibitor of mono-amine oxydase A (RIMA) is chosen from the group consisting of toloxatone, RS 8359, moclobemide, cimoxatone, caroxazone (F.I 6654) and befloxatone, or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof. More preferably, said compound which is a reversible inhibitor of mono-amine oxydase A (RIMA) is befloxatone and is to be administered in a daily dose ranging between 2.5 and 20 mg of the active ingredient.

10

The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) a compound which is a reversible inhibitor of mono-amine oxydase A (RIMA), preferably chosen from the group consisting of toloxatone, RS 8359, moclobemide, cimoxatone, caroxazone (F.I 6654) and befloxatone, or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of mental disease or disorder which is chosen from the group consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, adjustment disorders, impulse control disorders, personality disorders, antisocial behaviour, bereavement, occupational problem, problems related to abuse or neglect and pain disorders.

15

20

25

The invention also relates to a pharmaceutical composition as described above wherein pipamperon is provided in a unitary dose of between 5 and 15 mg of the active ingredient and wherein said compound which is a reversible inhibitor of mono-amine oxydase A (RIMA) is befloxatone, preferably provided in a unitary dose of between 2.5 and 20 mg of the active ingredient.

30

83: combination therapy with a compound which modulates SCT-11

The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with a compound which modulates SCT-11 (i.e. SCT-11 is a G protein-coupled receptor), are

35

chosen from the group of diseases or disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, personality disorders, bereavement, occupational problem, problems related to abuse or neglect and pain disorders.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, substance-related disorders, personality disorders, bereavement, occupational problem and problems related to abuse or neglect, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a compound which modulates SCT-11 to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said compound which modulates SCT-11, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of pain disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a compound which modulates SCT-11 to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said compound which modulates SCT-11, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

According to a preferred embodiment, the invention relates to the use as described above, wherein said compound which modulates SCT-11 is SNEC-2 or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof.







The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) a compound which modulates SCT-11, preferably SNE-2 or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of a mental disease or disorder which is chosen from the group consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, personality disorders, bereavement, occupational problem, problems related to abuse or neglect and pain disorders.

84: combination therapy with a serotonin/dopamine antagonist compound (SDA)

The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with a serotonin/dopamine antagonist compound (SDA), are chosen from the group of diseases or disorders consisting of mood disorders, anxiety disorders, psychotic disorders, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sleep disorders, adjustment disorders, impulse control disorders, pervasive development disorders, disruptive behaviour disorders, substance-related disorders, personality disorders, psychological factors affecting medical conditions, malingering, antisocial behaviour, bereavement, occupational problem, identity problem, problems related to abuse or neglect, pain disorders and delirium.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting of mood disorders, anxiety disorders, psychotic disorders, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sleep disorders, adjustment disorders, impulse control disorders, pervasive development disorders, disruptive behaviour disorders, substance-related disorders, personality disorders, psychological factors affecting medical conditions, malingering, antisocial behaviour, bereavement, occupational problem, identity problem and problems related to abuse or neglect, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a serotonin/dopamine antagonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said serotonin/dopamine

antagonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

5 The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of pain disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a serotonin/dopamine antagonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said
10 serotonin/dopamine antagonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

15 The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a cognitive mental disease or disorder which is delirium, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a serotonin/dopamine antagonist compound to augment the therapeutic effect or to provide a
20 faster onset of the therapeutic effect of said serotonin/dopamine antagonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

25 According to a preferred embodiment, the invention relates to the uses as described above, wherein said serotonin/dopamine antagonist compound is chosen from the group consisting of zotepine, ziprasidone, SM-13496, SL 91.0177, sertindole, S-18327, risperidone, quetiapine fumarate (preferably sustained release formulation), quetiapine fumarate (preferably granules), quetiapine, perospirone, paliperidone, olanzapine, ocaperidone, LU 31-131, iloperidone, clozapine, BSF-190555, blonanserin, bifeprunox, asenapine and
30 aripiprazole, or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof. Even more preferably, said serotonin/dopamine antagonist compound is chosen from the group consisting of SL 91.0177, sertindole, perospirone, paliperidone, blonanserin, bifeprunox and asenapine, or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof. More preferably, said serotonin/dopamine
35 antagonist compound is sertindole and is to be administered in a daily dose ranging between 12 and 24 mg of the active ingredient. More preferably, said serotonin/dopamine antagonist

compound is paliperidone and is to be administered in a daily dose ranging between 3 and 15 mg of the active ingredient. More preferably, said serotonin/dopamine antagonist compound is asenapine and is to be administered in a daily dose ranging between 2.5 and 20 mg of the active ingredient.

5

The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) a serotonin/dopamine antagonist compound, preferably chosen from the group consisting of SL 91.0177, sertindole, perospirone, paliperidone, blonanserin, bifeprunox and asenapine, or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of a mental disease or disorder which is chosen from the group consisting mood disorders, anxiety disorders, psychotic disorders, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sleep disorders, adjustment disorders, impulse control disorders, pervasive development disorders, disruptive behaviour disorders, substance-related disorders, personality disorders, psychological factors affecting medical conditions, malingering, antisocial behaviour, bereavement, occupational problem, identity problem, problems related to abuse or neglect, pain disorders and delirium.

10

15

20

The invention also relates to a pharmaceutical composition as described above wherein pipamperon is provided in a unitary dose of between 5 and 15 mg of the active ingredient and wherein said serotonin/dopamine antagonist compound is sertindole, preferably provided in a unitary dose of between 12 and 24 mg of the active ingredient.

25

The invention also relates to a pharmaceutical composition as described above wherein pipamperon is provided in a unitary dose of between 5 and 15 mg of the active ingredient and wherein said serotonin/dopamine antagonist compound is paliperidone, preferably provided in a unitary dose of between 3 and 15 mg of the active ingredient.

30

The invention also relates to a pharmaceutical composition as described above wherein pipamperon is provided in a unitary dose of between 5 and 15 mg of the active ingredient and wherein said serotonin/dopamine antagonist compound is asenapine, preferably provided in a unitary dose of between 2.5 and 20 mg of the active ingredient.

85: combination therapy with a selective SDRI compound

The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with a selective serotonin and dopamine re-uptake inhibitor (SDRI) compound, are chosen from the group of diseases or disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sleep disorders, adjustment disorders, impulse control disorders, substance-related disorders, personality disorders, antisocial behaviour, bereavement, occupational problem, problems related to abuse or neglect, pain disorders, delirium, Alzheimer Disease, substance-related persisting dementia, vascular dementia, dementia due to HIV disease, dementia due to head trauma, dementia due to Parkinson Disease, dementia due to Huntington Disease, dementia due to Pick Disease, dementia due to Creutzfeldt-Jacob Disease, amnesic disorders due to a general medical condition, substance-induced persisting amnesic disorder, mild cognitive impairment disorder and other cognitive disorders.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sleep disorders, adjustment disorders, impulse control disorders, substance-related disorders, personality disorders, antisocial behaviour, bereavement, occupational problem and problems related to abuse or neglect, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a selective serotonin and dopamine reuptakeinhibitor (SDRI) compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said selective serotonin and dopamine reuptake inhibitor (SDRI) compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of pain disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from

or prior to the administration of a selective serotonin and dopamine reuptake inhibitor (SDRI) compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said selective serotonin and dopamine reuptake inhibitor (SDRI) compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging
5 between 5 and 15 mg of the active ingredient.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a cognitive mental disease or disorder selected from the group of
10 diseases and disorders consisting of delirium, Alzheimer Disease, substance-related persisting dementia, vascular dementia, dementia due to HIV disease, dementia due to head trauma, dementia due to Parkinson Disease, dementia due to Huntington Disease, dementia due to Pick Disease, dementia due to Creutzfeldt-Jacob Disease, amnestic disorders due to a general medical condition, substance-induced persisting amnestic disorder, mild cognitive
15 impairment disorder and other cognitive disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a selective serotonin and dopamine reuptake inhibitor (SDRI) compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said selective serotonin and dopamine reuptake inhibitor (SDRI) compound, further
20 characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

According to a preferred embodiment, the invention relates to the uses as described above, wherein said selective serotonin and dopamine reuptake inhibitor (SDRI) compound is
25 bazineprine, or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof.

The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) a selective serotonin and dopamine reuptake inhibitor (SDRI) compound, preferably
30 bazineprine or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of mental disease or disorder which is chosen from the group consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious
35 disorders, dissociative disorders, sleep disorders, adjustment disorders, impulse control disorders, substance-related disorders, personality disorders, antisocial behaviour,

bereavement, occupational problem, problems related to abuse or neglect, pain disorders, delirium, Alzheimer Disease, substance-related persisting dementia, vascular dementia, dementia due to HIV disease, dementia due to head trauma, dementia due to Parkinson Disease, dementia due to Huntington Disease, dementia due to Pick Disease, dementia due to Creutzfeldt-Jacob Disease, amnesic disorders due to a general medical condition, substance-induced persisting amnesic disorder, mild cognitive impairment disorder and other cognitive disorders.

86: combination therapy with a second messenger beta agonist compound

The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with a second messenger beta agonist compound, are chosen from the group of diseases or disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, substance-related disorders, personality disorders, bereavement, occupational problem, problems related to abuse or neglect and pain disorders.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, substance-related disorders, personality disorders, bereavement, occupational problem and problems related to abuse or neglect, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a second messenger beta agonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said second messenger beta agonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of pain disorders, characterized in that pipamperon or said

pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a second messenger beta agonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said second messenger beta agonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

According to a preferred embodiment, the invention relates to the uses as described above, wherein said second messenger beta agonist compound is chosen from the group consisting of SR 57227, rolipram and eplivanserin, or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof. More preferably, said second messenger beta agonist compound is rolipram and is to be administered in a daily dose ranging between 1.5 and 3 mg of the active ingredient.

The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) a second messenger beta agonist compound, preferably chosen from the group consisting of SR 57227, rolipram and eplivanserin or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of a mental disease or disorder which is chosen from the group consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, substance-related disorders, personality disorders, bereavement, occupational problem, problems related to abuse or neglect and pain disorders.

The invention also relates to a pharmaceutical composition as described above wherein pipamperon is provided in a unitary dose of between 5 and 15 mg of the active ingredient and wherein said second messenger beta agonist compound is rolipram, preferably provided in a unitary dose of between 1.5 and 3 mg of the active ingredient.

87: combination therapy with a secretin pancreatic hormone

The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with a secretin pancreatic hormone, are chosen from the group of diseases or disorders consisting of anxiety disorders, psychotic disorders, somatoform disorders (excluding pain disorders),

factitious disorders, dissociative disorders, sleep disorders, adjustment disorders, impulse control disorders, pervasive development disorders, disruptive behaviour disorders, substance-related disorders, personality disorders, psychological factors affecting medical conditions, malingering, antisocial behaviour, bereavement, occupational problem, identity problem, problems related to abuse or neglect, pain disorders and delirium.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting of anxiety disorders, psychotic disorders, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sleep disorders, adjustment disorders, impulse control disorders, pervasive development disorders, disruptive behaviour disorders, substance-related disorders, personality disorders, psychological factors affecting medical conditions, malingering, antisocial behaviour, bereavement, occupational problem, identity problem and problems related to abuse or neglect, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a secretin pancreatic hormone to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said secretin pancreatic hormone, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of pain disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a secretin pancreatic hormone to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said secretin pancreatic hormone, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a cognitive mental disease or disorder which is delirium, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a secretin

pancreatic hormone to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said secretin pancreatic hormone, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

5

According to a preferred embodiment, the invention relates to the uses as described above, wherein said secretin pancreatic hormone is RG1068 or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof.

10 The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) a secretin pancreatic hormone, preferably RG1068, or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of a mental disease or disorder which is chosen from the group consisting of anxiety disorders, 15 psychotic disorders, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sleep disorders, adjustment disorders, impulse control disorders, pervasive development disorders, disruptive behaviour disorders, substance-related disorders, personality disorders, psychological factors affecting medical conditions, malingering, antisocial behaviour, bereavement, occupational problem, identity problem, 20 problems related to abuse or neglect, pain disorders and delirium.

88: combination therapy with a sigma receptor agonist compound

The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with a 25 sigma receptor agonist compound, are chosen from the group of diseases or disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, attention-deficit disorders, personality disorders, bereavement, occupational 30 problem, problems related to abuse or neglect and pain disorders.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the 35 group of diseases and disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders),

factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, attention-deficit disorders, substance-related disorders, personality disorders, bereavement, occupational problem and problems related to abuse or neglect, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a sigma receptor agonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said sigma receptor agonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of pain disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a sigma receptor agonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said sigma receptor agonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

According to a preferred embodiment, the invention relates to the uses as described above, wherein said sigma receptor agonist compound is VPI-013 (also known as OPC-14523) or PRX-00023, preferably VPI-013 (also known as OPC-14523), or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof.

The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) a sigma receptor agonist compound, preferably VPI-013 (also known as OPC-14523) or PRX-00023, preferably VPI-013 (also known as OPC-14523), or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of a mental disease or disorder which is chosen from the group consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, attention-deficit disorders, personality disorders, bereavement, occupational problem, problems related to abuse or neglect and pain disorders.

89: combination therapy with a sigma receptor antagonist compound

The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with a sigma receptor antagonist compound, are chosen from the group of diseases or disorders consisting of mood disorders, anxiety disorders, psychotic disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, pervasive development disorders, disruptive behaviour disorders, substance-related disorders, personality disorders, bereavement, occupational problem, problems related to abuse or neglect, pain disorders and delirium.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting of mood disorders, anxiety disorders, psychotic disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, pervasive development disorders, disruptive behaviour disorders, substance-related disorders, personality disorders, bereavement, occupational problem and problems related to abuse or neglect,, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a sigma receptor antagonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said sigma receptor antagonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of pain disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a sigma receptor antagonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said sigma receptor antagonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a cognitive mental disease or disorder which is delirium, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a sigma receptor antagonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said sigma receptor antagonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

According to a preferred embodiment, the invention relates to the uses as described above, wherein said sigma receptor antagonist compound is chosen from the group consisting of SR 31742 and EMD 68843, or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof. More preferably, said sigma receptor antagonist compound is EMD 68843 and is to be administered in a daily dose ranging between 5 and 40 mg of the active ingredient.

The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) a sigma receptor antagonist compound, preferably chosen from the group consisting of SR 31742 and EMD 68843, or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of a mental disease or disorder which is chosen from the group consisting mood disorders, anxiety disorders, psychotic disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, pervasive development disorders, disruptive behaviour disorders, substance-related disorders, personality disorders, bereavement, occupational problem, problems related to abuse or neglect, pain disorders and delirium.

The invention also relates to a pharmaceutical composition as described above wherein pipamperon is provided in a unitary dose of between 5 and 15 mg of the active ingredient and wherein said sigma receptor antagonist compound is EMD 68843, preferably provided in a unitary dose of between 5 and 40 mg of the active ingredient.

90: combination therapy with a selective SNDRI compound

The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with a selective serotonin, nor-adrenaline and dopamine re-uptake inhibitor (SNDRI) compound, are chosen from the group of diseases or disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sleep disorders, adjustment disorders, impulse control disorders, attention-deficit disorders, substance-related disorders, personality disorders, antisocial behaviour, bereavement, occupational problem, problems related to abuse or neglect, pain disorders, delirium, Alzheimer Disease, substance-related persisting dementia, vascular dementia, dementia due to HIV disease, dementia due to head trauma, dementia due to Parkinson Disease, dementia due to Huntington Disease, dementia due to Pick Disease, dementia due to Creutzfeldt-Jacob Disease, amnesic disorders due to a general medical condition, substance-induced persisting amnesic disorder, mild cognitive impairment disorder and other cognitive disorders.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sleep disorders, adjustment disorders, impulse control disorders, attention-deficit disorders, substance-related disorders, personality disorders, antisocial behaviour, bereavement, occupational problem and problems related to abuse or neglect, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a selective serotonin, nor-adrenaline and dopamine re-uptake inhibitor (SNDRI) compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said selective serotonin, nor-adrenaline and dopamine re-uptake inhibitor (SNDRI) compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of pain disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from

or prior to the administration of a selective serotonin, nor-adrenaline and dopamine re-uptake inhibitor (SNDRI) compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said selective serotonin, nor-adrenaline and dopamine re-uptake inhibitor (SNDRI) compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a cognitive mental disease or disorder selected from the group of diseases and disorders consisting of delirium, Alzheimer Disease, substance-related persisting dementia, vascular dementia, dementia due to HIV disease, dementia due to head trauma, dementia due to Parkinson Disease, dementia due to Huntington Disease, dementia due to Pick Disease, dementia due to Creutzfeldt-Jacob Disease, amnestic disorders due to a general medical condition, substance-induced persisting amnestic disorder, mild cognitive impairment disorder and other cognitive disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a selective serotonin, nor-adrenaline and dopamine re-uptake inhibitor (SNDRI) compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said selective serotonin, nor-adrenaline and dopamine re-uptake inhibitor (SNDRI) compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

According to a preferred embodiment, the invention relates to the uses as described above, wherein said selective serotonin, nor-adrenaline and dopamine re-uptake inhibitor (SNDRI) compound is selected from the group consisting of NS 2330; McN 5652; DOV 216,303 and DOV 21,947; more preferably NS 2330 or DOV 216,303; or a pro-drug or an active metabolite thereof; or a pharmaceutically acceptable salt thereof.

The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) a selective serotonin, nor-adrenaline and dopamine re-uptake inhibitor (SNDRI) compound, preferably selected from the group consisting of NS 2330; McN 5652; DOV 216,303 and DOV 21,947, more preferably NS 2330 or DOV 216,303, or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion

dysregulation of mental disease or disorder which is chosen from the group consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sleep disorders, adjustment disorders, impulse control disorders, attention-deficit disorders, substance-related disorders, personality disorders, antisocial behaviour, bereavement, occupational problem, problems related to abuse or neglect, pain disorders, delirium, Alzheimer Disease, substance-related persisting dementia, vascular dementia, dementia due to HIV disease, dementia due to head trauma, dementia due to Parkinson Disease, dementia due to Huntington Disease, dementia due to Pick Disease, dementia due to Creutzfeldt-Jacob Disease, amnesic disorders due to a general medical condition, substance-induced persisting amnesic disorder, mild cognitive impairment disorder and other cognitive disorders.

91: combination therapy with a selective SNRI compound

The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with a selective serotonin and nor-adrenaline re-uptake inhibitor (SNRI) compound, are chosen from the group of diseases or disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sleep disorders, adjustment disorders, impulse control disorders, attention-deficit disorders, substance-related disorders, personality disorders, antisocial behaviour, bereavement, occupational problem, problems related to abuse or neglect and pain disorders.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sleep disorders, adjustment disorders, impulse control disorders, attention-deficit disorders, substance-related disorders, personality disorders, antisocial behaviour, bereavement, occupational problem and problems related to abuse or neglect, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a selective serotonin and nor-adrenaline re-uptake inhibitor (SNRI) compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said selective

serotonin and nor-adrenaline re-uptake inhibitor (SNRI) compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

- 5 The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of pain disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a selective serotonin and nor-adrenaline re-uptake inhibitor
- 10 (SNRI) compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said selective serotonin and nor-adrenaline re-uptake inhibitor (SNRI) compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.
- 15 According to a preferred embodiment, the invention relates to the uses as described above, wherein said selective serotonin and nor-adrenaline re-uptake inhibitor (SNRI) compound is selected from the group consisting of venlafaxine, tomoxetine, tandamine, talsupram, talopram, nefazodone, milnacipran, LY 113.821, duloxetine, desvenlafaxine and amoxapine, or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof.
- 20 Even more preferably, said selective serotonin and nor-adrenaline re-uptake inhibitor (SNRI) compound is chosen from the group consisting of venlafaxine, tomoxetine, milnacipran, duloxetine and desvenlafaxine, or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof. More preferably, said selective serotonin and nor-adrenaline re-uptake inhibitor (SNRI) compound is venlafaxine and is to be administered in a
- 25 daily dose ranging between 75 and 300 mg of the active ingredient. More preferably, said selective serotonin and nor-adrenaline re-uptake inhibitor (SNRI) compound is tomoxetine and is to be administered in a daily dose ranging between 0.475 and 3.8 mg/kg of the active ingredient. More preferably, said selective serotonin and nor-adrenaline re-uptake inhibitor (SNRI) compound is milnacipran and is to be administered in a daily dose ranging between
- 30 50 and 200 mg of the active ingredient. More preferably, said selective serotonin and nor-adrenaline re-uptake inhibitor (SNRI) compound is duloxetine and is to be administered in a daily dose ranging between 40 and 60 mg of the active ingredient.

- The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and
- 35 (b) a selective serotonin and nor-adrenaline re-uptake inhibitor (SNRI) compound, preferably selected from the group consisting of venlafaxine, tomoxetine, tandamine, talsupram,

154

talopram, nefazodone, milnacipran, LY 113.821, duloxetine, desvenlafaxine and amoxapine, or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of mental disease or disorder which is chosen from the group consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sleep disorders, adjustment disorders, impulse control disorders, attention-deficit disorders, substance-related disorders, personality disorders, antisocial behaviour, bereavement, occupational problem, problems related to abuse or neglect and pain disorders.

The invention also relates to a pharmaceutical composition as described above wherein pipamperon is provided in a unitary dose of between 5 and 15 mg of the active ingredient and wherein said selective serotonin and nor-adrenaline re-uptake inhibitor (SNRI) compound is venlafaxine, preferably provided in a unitary dose of between 75 and 300 mg of the active ingredient.

The invention also relates to a pharmaceutical composition as described above wherein pipamperon is provided in a unitary dose of between 5 and 15 mg of the active ingredient and wherein said selective serotonin and nor-adrenaline re-uptake inhibitor (SNRI) compound is tomoxetine, preferably provided in a unitary dose of between 0.475 and 3.8 mg/kg of the active ingredient.

The invention also relates to a pharmaceutical composition as described above wherein pipamperon is provided in a unitary dose of between 5 and 15 mg of the active ingredient and wherein said selective serotonin and nor-adrenaline re-uptake inhibitor (SNRI) compound is milnacipran, preferably provided in a unitary dose of between 50 and 200 mg of the active ingredient.

The invention also relates to a pharmaceutical composition as described above wherein pipamperon is provided in a unitary dose of between 5 and 15 mg of the active ingredient and wherein said selective serotonin and nor-adrenaline re-uptake inhibitor (SNRI) compound is duloxetine, preferably provided in a unitary dose of between 40 and 60 mg of the active ingredient.

92: combination therapy with a selective serotonin re-uptake inhibitor (SSRI) compound

The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with a selective serotonin re-uptake inhibitor (SSRI) compound, are chosen from the group of diseases or disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sleep disorders, adjustment disorders, impulse control disorders, substance-related disorders, personality disorders, antisocial behaviour, bereavement, occupational problem, problems related to abuse or neglect and pain disorders.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sleep disorders, adjustment disorders, impulse control disorders, substance-related disorders, personality disorders, antisocial behaviour, bereavement, occupational problem and problems related to abuse or neglect, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a selective serotonin re-uptake inhibitor (SSRI) compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said selective serotonin re-uptake inhibitor (SSRI) compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of pain disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a selective serotonin re-uptake inhibitor (SSRI) compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said selective serotonin re-uptake inhibitor (SSRI) compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

According to a preferred embodiment, the invention relates to the uses as described above, wherein said selective serotonin re-uptake inhibitor (SSRI) compound is selected from the group consisting of YM 992, VPI-013 (also known as OPC-14523), sertraline, paroxetine, LY 214.281, LU AA 21-004, Lu 35-138, litoxetine, ifoxetine, fluvoxamine (controlled release formulation), fluvoxamine, fluoxetine, femoxetine, escitalopram, EMD 68843, cyanodothepine, citalopram, venlafaxine, milnacipran, duloxetine, cericlamine and ademethionine (preferably s-adenosylmethionine), or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof. Even more preferably, said selective serotonin re-uptake inhibitor (SSRI) compound is chosen from the group consisting of litoxetine, fluvoxamine (controlled release formulation) and escitalopram, or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof. More preferably, said selective serotonin re-uptake inhibitor (SSRI) compound is fluvoxamine (controlled release formulation) and is to be administered in a daily dose ranging between 100 and 300 mg of the active ingredient. More preferably, said selective serotonin re-uptake inhibitor (SSRI) compound is escitalopram and is to be administered in a daily dose ranging between 10 and 20 mg of the active ingredient. More preferably, said selective serotonin re-uptake inhibitor (SSRI) compound is citalopram and is to be administered in a daily dose ranging between 10 and 40 mg of the active ingredient.

The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) a selective serotonin re-uptake inhibitor (SSRI) compound, preferably selected from the group consisting of YM 992, VPI-013 (also known as OPC-14523), sertraline, paroxetine, LY 214.281, LU AA 21-004, Lu 35-138, litoxetine, ifoxetine, fluvoxamine (controlled release formulation), fluvoxamine, fluoxetine, femoxetine, escitalopram, EMD 68843, cyanodothepine, citalopram, venlafaxine, milnacipran, duloxetine, cericlamine and ademethionine (preferably s-adenosylmethionine), or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of mental disease or disorder which is chosen from the group consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sleep disorders, adjustment disorders, impulse control disorders, substance-related disorders, personality disorders, antisocial behaviour, bereavement, occupational problem, problems related to abuse or neglect and pain disorders.

The invention also relates to a pharmaceutical composition as described above wherein pipamperon is provided in a unitary dose of between 5 and 15 mg of the active ingredient and wherein said selective serotonin re-uptake inhibitor (SSRI) compound is fluvoxamine (controlled release formulation), preferably provided in a unitary dose of between between
5 100 and 300 mg of the active ingredient.

The invention also relates to a pharmaceutical composition as described above wherein pipamperon is provided in a unitary dose of between 5 and 15 mg of the active ingredient and wherein said selective serotonin re-uptake inhibitor (SSRI) compound is escitalopram,
10 preferably provided in a unitary dose of between 10 and 20 mg of the active ingredient.

The invention also relates to a pharmaceutical composition as described above wherein pipamperon is provided in a unitary dose of between 5 and 15 mg of the active ingredient and wherein said selective serotonin re-uptake inhibitor (SSRI) compound is citalopram,
15 preferably provided in a unitary dose of between 10 and 40 mg of the active ingredient.

Citalopram or citalopram hydrobromide is a selective serotonin (5-hydroxytryptamine / 5-HT) re-uptake inhibitor (SSRI) and is the conventional name given for the compound of the formula (RS)-1-[3-(dimethylamino)propyl]-1-(p-fluorophenyl)-5-phthalanarbonitrile-hydro-
20 bromide. According to an embodiment, a daily doses of active ingredient of SSRI, preferably citalopram, ranges between 10 and 40 mg per day. Preferably, daily doses of active ingredient ranging between 20 and 30 mg per day are administered. More preferably, a daily dose of 10, 15, 20, 25, 30, 35 or 40 mg per day is administered.

Fluvoxamine or fluvoxamine maleate (luvox, fevarin) is a selective serotonin (5-HT) re-uptake inhibitor (SSRI) belonging to a new chemical series, the 2-aminoethyl oxime ethers of aralkylketones. It is chemically unrelated to other SSRIs and clomipramine. It is chemically designated as 5-methoxy-4'-(trifluoromethyl) valerophenone (E)-O-(2-aminoethyl) oxime maleate (1:1).
25

According to an embodiment, a daily dose of active ingredient of fluvoxamine in a controlled release mode ranges between 100 and 300 mg per day. Preferably, daily doses of active ingredient ranging between 150 and 200 mg per day are administered in a controlled release mode. More preferably, a daily dose of 100, 150, 200, 250 or 300 mg per day is administered by controlled release.
30

93: combination therapy with a substance P receptor (NK1) antagonist compound

The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with a substance P receptor (NK1) antagonist compound, are chosen from the group of diseases or disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, substance-related disorders, personality disorders, bereavement, occupational problem, problems related to abuse or neglect and pain disorders.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, substance-related disorders, personality disorders, bereavement, occupational problem and problems related to abuse or neglect, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a substance P receptor (NK1) antagonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said substance P receptor (NK1) antagonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of pain, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a substance P receptor (NK1) antagonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said substance P receptor (NK1) antagonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

According to a preferred embodiment, the invention relates to the uses as described above, wherein said substance P receptor (NK1) antagonist compound is chosen from the group consisting of vestipitant, TAK-637, R673, GW823296, GW679769, GW597599, CP-122.721, aprepitant, 823296 and 679769, or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof. More preferably, said substance P receptor (NK1) antagonist compound is aprepitant and is to be administered in a daily dose ranging between 40 and 160 mg of the active ingredient.

The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) a substance P receptor (NK1) antagonist compound, preferably chosen from the group consisting of vestipitant, TAK-637, R673, GW823296, GW679769, GW597599, CP-122.721, aprepitant, 823296 and 679769, or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of a mental disease or disorder which is chosen from the group of diseases and disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, substance-related disorders, personality disorders, bereavement, occupational problem, problems related to abuse or neglect and pain disorders.

The invention also relates to a pharmaceutical composition as described above wherein pipamperon is provided in a unitary dose of between 5 and 15 mg of the active ingredient and wherein said substance P receptor (NK1) antagonist compound is aprepitant, preferably provided in a unitary dose of between 40 and 160 mg of the active ingredient.

94: combination therapy with a sulfonamide compound

The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with a sulfonamide compound, are chosen from the group of diseases or disorders consisting of mood disorders, psychotic disorders, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sleep disorders, adjustment disorders, impulse control disorders, pervasive development disorders, disruptive behaviour disorders, substance-related disorders, personality disorders, psychological factors affecting medical conditions, malingering, antisocial behaviour, bereavement, occupational problem, identity problem, problems related to abuse or neglect, pain disorders and delirium.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting of mood disorders, psychotic disorders, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sleep disorders, adjustment disorders, impulse control disorders, pervasive development disorders, disruptive behaviour disorders, substance-related disorders, personality disorders, psychological factors affecting medical conditions, malingering, antisocial behaviour, bereavement, occupational problem, identity problem and problems related to abuse or neglect, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a sulfonamide compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said sulfonamide compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of pain disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a sulfonamide compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said sulfonamide compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a cognitive mental disease or disorder which is delirium, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a sulfonamide compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said sulfonamide compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

According to a preferred embodiment, the invention relates to the uses as described above, wherein said sulfonamide compound is zonisamide or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof. More preferably, said sulfonamide compound is zonisamide and is to be administered in a daily dose ranging between 100 and 600 mg of the active ingredient.

The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) a sulfonamide compound, preferably zonisamide, or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of a mental disease or disorder which is chosen from the group consisting mood disorders, psychotic disorders, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sleep disorders, adjustment disorders, impulse control disorders, pervasive development disorders, disruptive behaviour disorders, substance-related disorders, personality disorders, psychological factors affecting medical conditions, malingering, antisocial behaviour, bereavement, occupational problem, identity problem, problems related to abuse or neglect, pain disorders and delirium.

The invention also relates to a pharmaceutical composition as described above wherein pipamperon is provided in a unitary dose of between 5 and 15 mg of the active ingredient and wherein said sulfonamide compound is zonisamide, preferably provided in a unitary dose of between 100 and 600 mg of the active ingredient.

95: combination therapy with a tachykinin antagonist compound

The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with a tachykinin antagonist compound, are chosen from the group of diseases or disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, substance-related disorders, personality disorders, bereavement, occupational problem, problems related to abuse or neglect and pain disorders.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group

of diseases and disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, substance-related disorders, personality disorders, bereavement, occupational problem and problems related to abuse or neglect, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a tachykinin antagonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said tachykinin antagonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of pain disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a tachykinin antagonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said tachykinin antagonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

According to a preferred embodiment, the invention relates to the use as described above, wherein said tachykinin antagonist compound is SR 48968 or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof.

The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) a tachykinin antagonist compound, preferably SR 48968 or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of a mental disease or disorder which is chosen from the group consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, substance-related disorders, personality disorders, bereavement, occupational problem, problems related to abuse or neglect and pain disorders.

96: combination therapy with a compound selected from the group consisting of R228060 (YKP-10A), palanpanel, ORG 39479/PH80, ORG 34167, DP 543 and CJ-017.493

The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with a compound selected from the group consisting of R228060 (YKP-10A), palanpanel, ORG 39479/PH80, ORG 34167, DP 543 and CJ-017.493, are chosen from the group of diseases or disorders consisting of mood disorders, anxiety disorders, psychotic disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, pervasive development disorders, attention-deficit disorders, disruptive behaviour disorders, substance-related disorders, personality disorders, psychological factors affecting medical conditions, malingering, antisocial behaviour, bereavement, occupational problem, identity problem, phase of life problem, academic problem, problems related to abuse or neglect, pain disorders, delirium, Alzheimer Disease, substance-related persisting dementia, vascular dementia, dementia due to HIV disease, dementia due to head trauma, dementia due to Parkinson Disease, dementia due to Huntington Disease, dementia due to Pick Disease, dementia due to Creutzfeldt-Jacob Disease, amnesic disorders due to a general medical condition, substance-induced persisting amnesic disorder, mild cognitive impairment disorder, other cognitive disorders and Parkinson Disease.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting of mood disorders, anxiety disorders, psychotic disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, pervasive development disorders, attention-deficit disorders, disruptive behaviour disorders, substance-related disorders, personality disorders, psychological factors affecting medical conditions, malingering, antisocial behaviour, bereavement, occupational problem, identity problem, phase of life problem, academic problem and problems related to abuse or neglect, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a compound selected from the group consisting of R228060 (YKP-10A), palanpanel, ORG 39479/PH80, ORG 34167, DP 543 and CJ-017.493, to augment the therapeutic effect or to provide a

faster onset of the therapeutic effect of said compound selected from the group consisting of R228060 (YKP-10A), palanpanel, ORG 39479/PH80, ORG 34167, DP 543 and CJ-017.493, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

5

The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) a compound selected from the group consisting of R228060 (YKP-10A), palanpanel, ORG 39479/PH80, ORG 34167, DP 543 and CJ-017.493, or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of mental disease or disorder which is chosen from the group consisting of mood disorders, anxiety disorders, psychotic disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, pervasive development disorders, attention-deficit disorders, disruptive behaviour disorders, substance-related disorders, personality disorders, psychological factors affecting medical conditions, malingering, antisocial behaviour, bereavement, occupational problem, identity problem, phase of life problem, academic problem, problems related to abuse or neglect, pain disorders, delirium, Alzheimer Disease, substance-related persisting dementia, vascular dementia, dementia due to HIV disease, dementia due to head trauma, dementia due to Parkinson Disease, dementia due to Huntington Disease, dementia due to Pick Disease, dementia due to Creutzfeldt-Jacob Disease, amnesic disorders due to a general medical condition, substance-induced persisting amnesic disorder, mild cognitive impairment disorder, other cognitive disorders and Parkinson Disease.

25

97: combination therapy with a vasopressin 1B receptor (V1B) antagonist compound

The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with a vasopressin 1B receptor (V1B) antagonist compound, are chosen from the group of diseases or disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, substance-related disorders, personality disorders, bereavement, occupational problem, problems related to abuse or neglect and pain disorders.

35

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, substance-related disorders, personality disorders, bereavement, occupational problem and problems related to abuse or neglect, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a vasopressin 1B receptor (V1B) antagonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said vasopressin 1B receptor (V1B) antagonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of pain disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a vasopressin 1B receptor (V1B) antagonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said vasopressin 1B receptor (V1B) antagonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

According to a preferred embodiment, the invention relates to the uses as described above, wherein said vasopressin 1B receptor (V1B) antagonist compound is SSR149415 or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof.

The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) a vasopressin 1B receptor (V1B) antagonist compound, preferably SSR149415 or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of a mental disease or disorder which is chosen from the group consisting of mood disorders, anxiety disorders, eating disorders, premenstrual

syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, substance-related disorders, personality disorders, bereavement, occupational problem, problems related to abuse or neglect and pain disorders.

5

98: combination therapy with a voltage-gated calcium channel $\alpha(2)\delta$ subunit modulator compound

10 The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with a voltage-gated calcium channel $\alpha(2)\delta$ subunit modulator compound, are chosen from the group of diseases or disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, substance-related disorders, 15 personality disorders, bereavement, occupational problem, problems related to abuse or neglect and pain disorders.

20 The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, substance-related disorders, personality disorders, bereavement, occupational problem and problems related to abuse or neglect, 25 characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a voltage-gated calcium channel $\alpha(2)\delta$ subunit modulator compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said voltage-gated calcium channel $\alpha(2)\delta$ subunit modulator compound, further characterized in that 30 pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

35 The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of pain disorders, characterized in that pipamperon or said

pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a voltage-gated calcium channel $\alpha(2)\delta$ subunit modulator compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said voltage-gated calcium channel $\alpha(2)\delta$ subunit modulator compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

According to a preferred embodiment, the invention relates to the uses as described above, wherein said voltage-gated calcium channel $\alpha(2)\delta$ subunit modulator compound is pregabalin or PD-200,390; or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof. More preferably, said voltage-gated calcium channel $\alpha(2)\delta$ subunit modulator compound is pregabalin, and is to be administered in a daily dose ranging between 50 and 600 mg of the active ingredient.

The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) a voltage-gated calcium channel $\alpha(2)\delta$ subunit modulator compound, preferably pregabalin or PD-200,390; or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of a mental disease or disorder which is chosen from the group consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, substance-related disorders, personality disorders, bereavement, occupational problem, problems related to abuse or neglect and pain disorders.

The invention also relates to a pharmaceutical composition as described above wherein pipamperon is provided in a unitary dose of between 5 and 15 mg of the active ingredient and wherein said voltage-gated calcium channel $\alpha(2)\delta$ subunit modulator compound is pregabalin, preferably provided in a unitary dose of between 50 and 600 mg of the active ingredient.

99: combination therapy with a vomeropherin compound

The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with a vomeropherin compound, are chosen from the group of diseases or disorders consisting of

anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, personality disorders, bereavement, occupational problem and problems related to abuse or neglect.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting of anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, personality disorders, bereavement, occupational problem and problems related to abuse or neglect, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of vomeropherin compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said vomeropherin compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

According to a preferred embodiment, the invention relates to the use as described above, wherein said vomeropherin compound is PH94B or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof.

The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) vomeropherin compound, preferably PH94B or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder which is chosen from the group consisting of anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, personality disorders, bereavement, occupational problem and problems related to abuse or neglect.

Also, the invention relates in particular to the use as described before, wherein said second compound is chosen from the group consisting of fluvoxamine controlled release,

phenserine tartrate, atomoxetine hydrochloride, bupropion (controlled-release formulation), ropinirole HCL (controlled-release formulation), INN 00835, galantamine (extended release formulation), paliperidone, tomoxetine, aprepitant, rivastigmine tartrate, ORG 34517/34850, sunepitron, sumanirole, milnacipran, idazoxan, xaliproden, SR 58611, befloxatone, litoxetine, 5 tianeptine, agomelatine, SPD 503, flesinoxan, bifeprunox, ramelteon, etilevodopa, rasagiline (TVP-1012) and desvenlafaxine.

Also, the invention relates in particular to the use as described before, wherein said second compound is chosen from the group consisting of galantamine (extended release formulation), R121919, risperidone, paliperidone and R228060 (YKP-10A).

10 The disclosure of all patents, publications (including published patent publications), and database accession numbers and depository accession numbers referenced in this specification are specifically incorporated herein by reference in their entirety to the same extent as if each such individual patent, publication, and database accession number, and 15 depository accession number were specifically and individually indicated to be incorporated by reference.

The invention, now being generally described, will be more readily understood by reference to the following tables and examples, which are included merely for purposes of illustration of certain aspects and embodiments of the present invention and are not intended to limit the invention.

Short description of the Tables and Figures

Table 1: In Table 1, the pKi values of test compounds are given for each of the dopamine receptors, 5HT receptors, adrenergic receptors and the histamine₁ receptor.

5

Table 2: Set-up of a clinical trial comprising for treatment groups.

Table 3: Overview of a placebo, active and period controlled clinical trial in a fore-going pipamperon - citalopram treatment in Major Depressive Disorder.

10

Table 4: POC process for major depressive disorder.

Table 5: Summary of diseases and disorders relative to known psycho-tropics.

15

Table 6: Overview of Pharmacological grouping, indicating pharmacological profile numbering (column 2), pharmacological profile (column 3), main indication(s) (column 4), name of the compound (column 4), the dose range (column 5), and the company producing or selling said compound (column 6). Compounds indicated by hatching are preferred.

20

Figure 1: Add-on treatment with pipamperon after treatment with citalopram .

Figure 2: HDRS-17 change from baseline: combo treatment pipamperon as add-on - citalopram vs SNRI (duloxetine) in Major Depression.

25

Figure 3: Remission rates (HDRS-17 ≤ 7): combo treatment pipamperon as add-on - citalopram vs SNRI (venlafaxine) vs SSRIs vs placebo in Major Depression.

Figure 4: Fore-going treatment during 1-5 days with pipamperon followed with the combination treatment of pipamperon and citalopram.

30

Figure 5: HDRS-17 change from baseline: combo treatment pipamperon - citalopram with a fore-going treatment of 4 days with pipamperon vs SNRI (duloxetine) in Major Depression.

35

Figure 6: Remission rates (HDRS-17 ≤ 7): combo pipamperon - citalopram with a fore-going treatment of 4 days with pipamperon vs SNRI (venlafaxine) in Major Depression.

Figure 7: Fore-going treatment during 6-8 days with pipamperon followed with the combination treatment of pipamperon and citalopram.

Figure 8: HDRS-17 change from baseline: combo treatment pipamperon - citalopram with a fore-going treatment of 7 days with pipamperon vs SNRI (duloxetine) in Major Depression.

Figure 9: Fore-going and add-on treatment with pipamperon in MDD.

Figure 10: HDRS-17 change from baseline: fore-going and add-on treatment with pipamperon and citalopram in comparison with the SNRI duloxetine in Major Depression.

Figure 11: Remission rates (HDRS-17 ≤ 7): fore-going and add-on treatment with pipamperon and citalopram in comparison with the SNRI venlafaxine in Major Depression.

Figure 12: Y-BOCS total score: fore-going and add-on treatment with pipamperon and citalopram in comparison with the SSRI fluvoxamine in OCD.

Figure 13: Y-BOCS obsession score: fore-going and add-on treatment with pipamperon and citalopram in comparison with the SSRI fluvoxamine in OCD.

Figure 14: Y-BOCS compulsion score: fore-going and add-on treatment with pipamperon and citalopram in comparison with the SSRI fluvoxamine in OCD.

Table 1

Figure 1S: CGI-severity score:	D1	D2	D3	D4	SHT _{1A}	SHT _{1B}	SHT _{1D}	SHT _{1E}	SHT _{1F}	SHT _{2A}	SHT _{2B}	SHT _{2C}	SHT _{6rat}	SHT _{7rat}	Alpha1	Alpha2	Alpha2	Alpha2	Beta1	Beta2	H1
ORG5222	8-9	8-9	8-9	8-9	8-9	8-9	8-9	7-8	0	<9	<9	<9	<9	<9	<9	8-9	<9	7-8	<9	<9	<9
Zotepine	0	8-9	8-9	7-8	6-7	7-8	7-8	6-7	0	8-9	0	0	0	0	0	6-7	8-9	6-7	<9	<9	<9
Fluparoxan	0	<6	<6	6-7	6-7	<6	<6	0	0	8-9	0	<6	0	0	6-7	8-9	8-9	8-9	0	0	0
Olanzapine	7-8	7-8	7-8	7-8	<9	6-7	6-7	<6	6-7	8-9	8-9	8-9	7-8	6-7	7-8	6-7	6-7	6-7	<6	<6	<9
Clozapine	7-8	6-7	6-7	7-8	6-7	6-7	6-7	6-7	6-7	8-9	8-9	7-8	7-8	7-8	8-9	7-8	7-8	7-8	<6	<6	<9
S16924	0	7-8	7-8	7-8	8-9	0	0	0	0	<9	8-9	7-8	7-8	7-8	8-9	6-7	7-8	6-7	<6	<6	0
S18327	7-8	7-8	6-7	8-9	7-8	0	0	0	0	8-9	0	6-7	0	0	<9	6-7	0	0	0	0	0
Amperozide	6-7	6-7	6-7	6-7	<6	0	0	0	0	8-9	0	<6	0	0	7-8	<6	0	0	0	0	0
GGR218231	<6	7-8	>9	6-7	6-7	<6	<6	0	0	6-7	<6	<6	0	0	<6	<6	0	0	0	0	0
Serindole	7-8	8-9	8-9	7-8	6-7	7-8	7-8	6-7	6-7	>9	0	8-9	0	0	<9	6-7	6-7	6-7	<6	<6	6-7
MDL100,907	6-7	<6	<6	6-7	<6	0	0	0	0	<9	0	7-8	0	0	<6	<6	0	0	0	0	0
Haloperidol	8-9	>9	8-9	8-9	<6	6-7	<6	<6	<6	6-7	<9	<6	<6	6-7	8-9	<6	6-7	<6	<6	<6	6-7
Tiospirone	7-8	8-9	8-9	8-9	8-9	0	0	0	0	<9	0	8-9	0	0	<9	6-7	0	0	0	0	0
Raclopride	<6	8-9	8-9	6-7	<6	0	0	0	0	6-7	0	<6	0	0	<6	<6	0	0	0	0	0
Fluspirilene	0	<9	8-9	8-9	7-8	<6	<6	<6	0	6-7	0	0	0	0	0	6-7	7-8	7-8	6-7	6-7	7-8
Ocaperidone	7-8	>9	8-9	8-9	7-8	0	0	0	0	<9	0	7-8	0	0	<9	0	0	0	0	0	0
Risperidone	7-8	8-9	7-8	8-9	6-7	8-9	6-7	<6	<6	<9	0	7-8	0	0	<9	7-8	8-9	8-9	<6	<6	7-8
S33084	6-7	7-8	>9	6-7	<6	6-7	6-7	6-7	0	6-7	0	7-8	0	0	6-7	<6	0	0	0	0	0
L741626	6-7	8-9	7-8	6-7	<6	6-7	<6	0	0	6-7	6-7	<6	0	0	6-7	<6	0	0	0	0	0
Seroquel	6-7	6-7	6-7	6-7	6-7	<6	<6	<6	<6	6-7	6-7	6-7	0	0	7-8	<6	7-8	6-7	<6	<6	8-9
Yohimbine	0	6-7	<6	0	7-8	6-7	7-8	0	0	6-7	0	<6	0	0	6-7	7-8	<6	8-9	<6	<6	0
Ziprasidone	8-9	8-9	7-8	7-8	8-9	>9	8-9	6-7	0	6-7	8-9	8-9	7-8	8-9	8-9	6-7	7-8	7-8	<6	<6	7-8
Pipamperon	0	6-7	6-7	8-9	<6	6-7	6-7	<6	<6	8-9	0	0	0	0	0	6-7	7-8	6-7	<6	<6	<6

Table 2

173

ACUTE PHASE**					EXTENSION PHASE***										FOLLOW-UP PHASE					
VISITS	V1 Screen minus D7	V2 Baseline D0	V3 D4	V4 D7	V5 W2	V6 W3	V7 W4	V8 W6	V9 W8	V10 W10	V11 W12	V12 W16	V13 W20	V14 W24	V15 M8	V16 M10	V17 M12	V18 W1	V19 W2	
Day / Week / Month																				
TREATMENT GROUP																				
Group Pip-Active / D7	A	B	B	C														A	A	
Group Pip-Active / D4	A	B	C															A	A	
Group Pip-Active / D0	A	C																A	A	
Group Plc-Active / D0	A	D																A	A	
Informed Consent	X																			
NEC*	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	
Vital Signs / Weight	X																			
LAB	X																			
ECG	X																			
Phys Exam	X																			
Alt/Drugs Screen	X																			
CGI-S***	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	
QLESQ****	X																			
Treatment regimen:																				

Treatment regimen:

A: PLC + PLC

B: 2x(PLC+PIP(4mg))/d

C: 2x(CIT(10mg)+PIP(4mg))/d

D: 2x(CIT(10mg)+PLC)/d

*Neuronal E-Clinical Trial = Vesalius Expert Development for this Trial which includes the bottom-up measurement of:

**Entering Acute Phase: only NON-placebo responders as defined by the DSM-IV criteria of efficacy

***Entering Extension Phase: only remitters as defined by the DSM-IV criteria of efficacy

****CGI-S : Clinical Global Impressions-Improvement Scale

*****Q-LES-Q : Quality of Life, Enjoyment and Satisfaction Questionnaire

Table 3

FOREGOING PIPAMPERON - CITALOPRAM TREATMENT IN MAJOR DEPRESSIVE DISORDER : A PLACEBO, ACTIVE AND PERIODE CONTROLLED																			
CLINICAL TRIAL										FOLLOW-UP PHASE									
ACUTE PHASE**										EXTENSION PHASE***									
VISITS	V1 SCREEN minus D7	V2 BASELINE D0	V3 D4	V4 D7	V5 W2	V6 W3	V7 W4	V8 W6	V9 W8	V10 W10	V11 W12	V12 W16	V13 W20	V14 W24	V15 M8	V16 M10	V17 M12	V18 W1	V19 W2
TREATMENT GROUP																			
Group Pip - Active / Day 4		2x(PLC+PIP(4mg))id	2x(CIT(10mg)+PIP(4mg))id															PLC+PLC	PLC+PLC
Group Pip - Active / Day 0		2x(CIT(10mg)+PIP(4mg))id																PLC+PLC	PLC+PLC
Group Plc - Active / Day 0		2x(CIT(10mg)+PLC)id																PLC+PLC	PLC+PLC
Group Placebo																			
Informed Consent	x																		
NECT	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x
Vital Signs / Weight	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x
LAB	x	x																	
ECG	x																		
Phys Exam	x						x												
Atc/Drugs Screen	x						x												
CGLS	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x
****Q-LES-Q	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x

*Neuronal E-Clinical Trial = Vesalius Expert Development for this Trial which includes the bottom-up measurement of:

- . In- and exclusion criteria
- . Functional status evaluation
- . Medical history
- . (Pre-)treatment signs & symptoms
- . DSM-IV rules for diagnosis & efficacy
- . Rating Scales: HDRS-28, MADRS, HAMA
- . Medical resource utilisation
- . Pre-trial & concomitant medication
- . Drug administration
- . (Serious) Adverse events
- . Admission to the acute and extension phase of treatment
- . Right flow of the trial

**Entering Acute Phase: only NON-placebo responders as defined by the DSM-IV criteria of efficacy

***Entering Extension Phase: only remitters as defined by the DSM-IV criteria of efficacy

****Q-LES-Q: Quality of Life, Enjoyment and Satisfaction Questionnaire

175

Table 4

DAY	minus D7	D0	=>D4
TREATMENTGROUP			
Placebo (PLC)	PLC+PLC	2x(PLC+PLC)	2x(PLC+PC)
PIP - Active / Day 4	PLC+PLC	2x(PLC+PIP (4mg))/d	2x(CIT (10mg)+PIP (4mg))/d
PIP - Active / Day 0	PLC+PLC	2x(CIT (10mg)+PIP (4mg))/d	2x(CIT (10mg)+PIP (4mg))/d
PLC - Active / Day 0	PLC+PLC	2x(CIT (10mg)+PLC)/d	2x(CIT (10mg)+PLC)/d

[illegible]

MERCAPTAN											
COMPOUND											
MONO											
5-HT _{2A} /D ₄ Antagonist											
5-HT _{2A} /D ₄ Antagonist											
5-HT _{2A} /D ₄ Antagonist											
5-HT _{2A} /D ₄ Antagonist											
5-HT _{2A} /D ₄ Antagonist											
5-HT _{2A} /D ₄ Antagonist											
5-HT _{2A} /D ₄ Antagonist											
5-HT _{2A} /D ₄ Antagonist											
5-HT _{2A} /D ₄ Antagonist											
5-HT _{2A} /D ₄ Antagonist											
5-HT _{2A} /D ₄ Antagonist											
5-HT _{2A} /D ₄ Antagonist											
5-HT _{2A} /D ₄ Antagonist											
5-HT _{2A} /D ₄ Antagonist											
5-HT _{2A} /D ₄ Antagonist											
5-HT _{2A} /D ₄ Antagonist											
5-HT _{2A} /D ₄ Antagonist											
5-HT _{2A} /D ₄ Antagonist											
5-HT _{2A} /D ₄ Antagonist											
5-HT _{2A} /D ₄ Antagonist											
5-HT _{2A} /D ₄ Antagonist											
5-HT _{2A} /D ₄ Antagonist											
5-HT _{2A} /D ₄ Antagonist											
5-HT _{2A} /D ₄ Antagonist											
5-HT _{2A} /D ₄ Antagonist											
5-HT _{2A} /D ₄ Antagonist											
5-HT _{2A} /D ₄ Antagonist											
5-HT _{2A} /D ₄ Antagonist											
5-HT _{2A} /D ₄ Antagonist											
5-HT _{2A} /D ₄ Antagonist											
5-HT _{2A} /D ₄ Antagonist											
5-HT _{2A} /D ₄ Antagonist											
5-HT _{2A} /D ₄ Antagonist											
5-HT _{2A} /D ₄ Antagonist											
5-HT _{2A} /D ₄ Antagonist											
5-HT _{2A} /D ₄ Antagonist											
5-HT _{2A} /D ₄ Antagonist											
5-HT _{2A} /D ₄ Antagonist											
5-HT _{2A} /D ₄ Antagonist											
5-HT _{2A} /D ₄ Antagonist											
5-HT _{2A} /D ₄ Antagonist											
5-HT _{2A} /D ₄ Antagonist											
5-HT _{2A} /D ₄ Antagonist											
5-HT _{2A} /D ₄ Antagonist											
5-HT _{2A} /D ₄ Antagonist											
5-HT _{2A} /D ₄ Antagonist											
5-HT _{2A} /D ₄ Antagonist											
5-HT _{2A} /D ₄ Antagonist											
5-HT _{2A} /D ₄ Antagonist											
5-HT _{2A} /D ₄ Antagonist											
5-HT _{2A} /D ₄ Antagonist											
5-HT _{2A} /D ₄ Antagonist											
5-HT _{2A} /D ₄ Antagonist											
5-HT _{2A} /D ₄ Antagonist											
5-HT _{2A} /D ₄ Antagonist											
5-HT _{2A} /D ₄ Antagonist											
5-HT _{2A} /D ₄ Antagonist											
5-HT _{2A} /D ₄ Antagonist											
5-HT _{2A} /D ₄ Antagonist											
5-HT _{2A} /D ₄ Antagonist											
5-HT _{2A} /D ₄ Antagonist											
5-HT _{2A} /D ₄ Antagonist											
5-HT _{2A} /D ₄ Antagonist											
5-HT _{2A} /D ₄ Antagonist											
5-HT _{2A} /D ₄ Antagonist											
5-HT _{2A} /D ₄ Antagonist											
5-HT _{2A} /D ₄ Antagonist											
5-HT _{2A} /D ₄ Antagonist											
5-HT _{2A} /D ₄ Antagonist											
5-HT _{2A} /D ₄ Antagonist											
5-HT _{2A} /D ₄ Antagonist											
5-HT _{2A} /D ₄ Antagonist											
5-HT _{2A} /D ₄ Antagonist											
5-HT _{2A} /D ₄ Antagonist											
5-HT _{2A} /D ₄ Antagonist											
5-HT _{2A} /D ₄ Antagonist											
5-HT _{2A} /D ₄ Antagonist											
5-HT _{2A} /D ₄ Antagonist											
5-HT _{2A} /D ₄ Antagonist											
5-HT _{2A} /D ₄ Antagonist											
5-HT _{2A} /D ₄ Antagonist											
5-HT _{2A} /D ₄ Antagonist											
5-HT _{2A} /D ₄ Antagonist											
5-HT _{2A} /D ₄ Antagonist											
5-HT _{2A} /D ₄ Antagonist											
5-HT _{2A} /D ₄ Antagonist											
5-HT _{2A} /D ₄ Antagonist											
5-HT _{2A} /D ₄ Antagonist											
5-HT _{2A} /D ₄ Antagonist											
5-HT _{2A} /D ₄ Antagonist											
5-HT _{2A} /D ₄ Antagonist											
5-HT _{2A} /D ₄ Antagonist											
5-HT _{2A} /D ₄ Antagonist											
5-HT _{2A} /D ₄ Antagonist											
5-HT _{2A} /D ₄ Antagonist											
5-HT _{2A} /D ₄ Antagonist											
5-HT _{2A} /D ₄ Antagonist											
5-HT _{2A} /D ₄ Antagonist											
5-HT _{2A} /D ₄ Antagonist											
5-HT _{2A} /D ₄ Antagonist											
5-HT _{2A} /D ₄ Antagonist											
5-HT _{2A} /D ₄ Antagonist											
5-HT _{2A} /D ₄ Antagonist											
5-HT _{2A} /D ₄ Antagonist											
5-HT _{2A} /D ₄ Antagonist											
5-HT _{2A} /D ₄ Antagonist											
5-HT _{2A} /D ₄ Antagonist											
5-HT _{2A} /D ₄ Antagonist											
5-HT _{2A} /D ₄ Antagonist											
5-HT _{2A} /D ₄ Antagonist											
5-HT _{2A} /D ₄ Antagonist											
5-HT _{2A} /D ₄ Antagonist											
5-HT _{2A} /D ₄ Antagonist											
5-HT _{2A} /D ₄ Antagonist											
5-HT _{2A} /D ₄ Antagonist											
5-HT _{2A} /D ₄ Antagonist											
5-HT _{2A} /D ₄ Antagonist											
5-HT _{2A} /D ₄ Antagonist											
5-HT _{2A} /D ₄ Antagonist											
5-HT _{2A} /D ₄ Antagonist											
5-HT _{2A} /D ₄ Antagonist											
5-HT _{2A} /D ₄ Antagonist											
5-HT _{2A} /D ₄ Antagonist											
5-HT _{2A} /D ₄ Antagonist											
5-HT _{2A} /D ₄ Antagonist											
5-HT _{2A} /D ₄ Antagonist											
5-HT _{2A} /D ₄ Antagonist											
5-HT _{2A} /D ₄ Antagonist											
5-HT _{2A} /D ₄ Antagonist											
5-HT _{2A} /D ₄ Antagonist											
5-HT _{2A} /D ₄ Antagonist											
5-HT _{2A} /D ₄ Antagonist											
5-HT _{2A} /D ₄ Antagonist											
5-HT _{2A} /D ₄ Antagonist											
5-HT _{2A} /D ₄ Antagonist											
5-HT _{2A} /D ₄ Antagonist											
5-HT _{2A} /D ₄ Antagonist											
5-HT _{2A} /D ₄ Antagonist											
5-HT _{2A} /D ₄ Antagonist											
5-HT _{2A} /D ₄ Antagonist											
5-HT _{2A} /D ₄ Antagonist											
5-HT _{2A} /D ₄ Antagonist											
5-HT _{2A} /D ₄ Antagonist											
5-HT _{2A} /D ₄ Antagonist											
5-HT _{2A} /D ₄ Antagonist											
5-HT _{2A} /D ₄ Antagonist											
5-HT _{2A} /D ₄ Antagonist											
5-HT _{2A} /D ₄ Antagonist											
5-HT _{2A} /D ₄ Antagonist											
5-HT _{2A} /D ₄ Antagonist											
5-HT _{2A} /D ₄ Antagonist											
5-HT _{2A} /D ₄ Antagonist											
5-HT _{2A} /D ₄ Antagonist											
5-HT _{2A} /D ₄ Antagonist											
5-HT _{2A} /D ₄ Antagonist											
5-HT _{2A} /D ₄ Antagonist											
5-HT _{2A} /D ₄ Antagonist											
5-HT _{2A} /D ₄ Antagonist											
5-HT _{2A} /D ₄ Antagonist											
5-HT											

[illegible]

MEDICATION																			
										36	50	74	77	83	84	87	88	91	95
MONO																			
COMBO																			
5-HT2A/2C Antagonist + CNS compound																			
5-HT2A/2C Antagonist										GABA _A antagonist	venoreptin	good agent	Podagrace A2 m1 agonist	Podagrace A2 m1 agonist	Podagrace A2 m1 agonist	Podagrace A2 m1 agonist	Podagrace A2 m1 agonist	Podagrace A2 m1 agonist	Podagrace A2 m1 agonist

MEDICAMENT															53	65	23	10	43	50	38	27	75	18	81	79	18	34	15															
MONO															COMBO															5-HT2A/DA Antagonist + CNS compound														
5-HT2A/DA Antagonist															AMPA receptor modulator															GABA ^A agonist														
															opioid receptor modulator															opioid receptor modulator														
															glucocorticoid synthesis inhibitor															glucocorticoid synthesis inhibitor														
															opioid receptor modulator															opioid receptor modulator														
															opioid receptor modulator															opioid receptor modulator														
															opioid receptor modulator															opioid receptor modulator														
															opioid receptor modulator															opioid receptor modulator														
															opioid receptor modulator															opioid receptor modulator														
															opioid receptor modulator															opioid receptor modulator														
															opioid receptor modulator															opioid receptor modulator														
															opioid receptor modulator															opioid receptor modulator														
															opioid receptor modulator															opioid receptor modulator														
															opioid receptor modulator															opioid receptor modulator														
															opioid receptor modulator															opioid receptor modulator														
															opioid receptor modulator															opioid receptor modulator														
															opioid receptor modulator															opioid receptor modulator														
															opioid receptor modulator															opioid receptor modulator														
															opioid receptor modulator															opioid receptor modulator														
															opioid receptor modulator															opioid receptor modulator														
															opioid receptor modulator															opioid receptor modulator														
															opioid receptor modulator															opioid receptor modulator														
															opioid receptor modulator															opioid receptor modulator														
															opioid receptor modulator															opioid receptor modulator														
															opioid receptor modulator															opioid receptor modulator														
															opioid receptor modulator															opioid receptor modulator														
															opioid receptor modulator															opioid receptor modulator														
															opioid receptor modulator															opioid receptor modulator														
															opioid receptor modulator															opioid receptor modulator														
															opioid receptor modulator															opioid receptor modulator														
															opioid receptor modulator															opioid receptor modulator														
															opioid receptor modulator															opioid receptor modulator														
															opioid receptor modulator															opioid receptor modulator														
															opioid receptor modulator															opioid receptor modulator														
															opioid receptor modulator															opioid receptor modulator														
															opioid receptor modulator															opioid receptor modulator														
															opioid receptor modulator															opioid receptor modulator														
															opioid receptor modulator															opioid receptor modulator														
															opioid receptor modulator															opioid receptor modulator														
															opioid receptor modulator															opioid receptor modulator														
															opioid receptor modulator															opioid receptor modulator														
															opioid receptor modulator															opioid receptor modulator														
															opioid receptor modulator															opioid receptor modulator														
															opioid receptor modulator															opioid receptor modulator														
															opioid receptor modulator															opioid receptor modulator														
															opioid receptor modulator															opioid receptor modulator														
															opioid receptor modulator															opioid receptor modulator														
															opioid receptor modulator															opioid receptor modulator														
															opioid receptor modulator															opioid receptor modulator														
															opioid receptor modulator															opioid receptor modulator														
															opioid receptor modulator															opioid receptor modulator														
															opioid receptor modulator															opioid receptor modulator														
															opioid receptor modulator															opioid receptor modulator														
															opioid receptor modulator															opioid receptor modulator														
															opioid receptor modulator															opioid receptor modulator														
															opioid receptor modulator															opioid receptor modulator														
															opioid receptor modulator															opioid receptor modulator														
															opioid receptor modulator															opioid receptor modulator														
															opioid receptor modulator															opioid receptor modulator														
															opioid receptor modulator															opioid receptor modulator														
															opioid receptor modulator															opioid receptor modulator														
															opioid receptor modulator															opioid receptor modulator														
															opioid receptor modulator															opioid receptor modulator														
															opioid receptor modulator															opioid receptor modulator														
															opioid receptor modulator															opioid receptor modulator														
															opioid receptor modulator															opioid receptor modulator														
															opioid receptor modulator															opioid receptor modulator														
															opioid receptor modulator															opioid receptor modulator														
															opioid receptor modulator															opioid receptor modulator														
															opioid receptor modulator															opioid receptor modulator														
															opioid receptor modulator															opioid receptor modulator														
															opioid receptor modulator															opioid receptor modulator														
															opioid receptor modulator															opioid receptor modulator														
															opioid receptor modulator															opioid receptor modulator														
															opioid receptor modulator															opioid receptor modulator														
															opioid receptor modulator															opioid receptor modulator														
															opioid receptor modulator															opioid receptor modulator														
															opioid receptor modulator															opioid receptor modulator														
															opioid receptor modulator															opioid receptor modulator														
															opioid receptor modulator															opioid receptor modulator														
															opioid receptor modulator															opioid receptor modulator														
															opioid receptor modulator															opioid receptor modulator														
															opioid receptor modulator															opioid receptor modulator														
															opioid receptor modulator															opioid receptor modulator														
															opioid receptor modulator															opioid receptor modulator														
															opioid receptor modulator															opioid receptor modulator														
															opioid receptor modulator															opioid receptor modulator														
															opioid receptor modulator															opioid receptor modulator														
															opioid receptor modulator															opioid receptor modulator														
															opioid receptor modulator															opioid receptor modulator														
															opioid receptor modulator															opioid receptor modulator														
															opioid receptor modulator															opioid receptor modulator														
															opioid receptor modulator															opioid receptor modulator														
															opioid receptor modulator															opioid receptor modulator														
															opioid receptor modulator															opioid receptor modulator														
															opioid receptor modulator															opioid receptor modulator														
															opioid receptor modulator</																													

[illegible]

[illegible]

GLOSSARY	
5-HT = serotonin	
5-HT1 = serotonin 1 receptor	
5-HT1A = serotonin 1A receptor	
5-HT1B = serotonin 1B receptor	
5-HT2AD4 = serotonin 2A en dopamine D4 receptor	
5-HT2B = serotonin 2B receptor	
5-HT2C = serotonin 2C receptor	
5HT3 = serotonin 3 receptor	
5HT6 = serotonin 6 receptor	
AMPA = alpha-amino-3-hydroxy-5-methyl-4-isoxazole propionate	
CB1 = cannabinoid receptor 1	
CINODs = COX-inhibiting nitric oxide donors	
COX = cyclooxygenase	
COX-2 = cyclooxygenase 2	
CRF-1 = Corticotropin-Releasing Factor Receptor 1	
D1 = Dopamine 1	
D2 = Dopamine 2	
D2 = Dopamine 3	
DA = Dopamine	
ERK = extracellular signal-related kinase	
GABA = gamma-aminobutyric acid	
GABA-A = gamma-aminobutyric acid A receptor	
GABA-B = gamma-aminobutyric acid B receptor	
GPCR = G-Protein-Coupled Receptor	
GR = glucocorticoid receptor	
H3 = histamine H3 receptor	
MAO = mono-amine oxidase	
MAO-A = mono-amine oxidase A	
MAO-B = mono-amine oxidase B	
MC4 = melanocortin-4 receptor	
MCH = Melanin concentrating hormone	
MgluR = metabotropic glutamate receptor	
MT = melatonin receptor	
NARI = selective nor-adrenaline re-uptake inhibitor	
NasSA = noradrenergic/specific serotonergic antidepressant	
NDRI = selective nor-adrenaline and dopamine re-uptake inhibitor	
NGF = Nerve Growth Factor	
NGF = nerve growth factor	
NK1 = neurokinin 1 receptor	
NK2 = neurokinin 2 receptor	
NK3 = neurokinin 3 receptor	
NMDA = N-Methyl-D-aspartate	
NSAID = Non-steroidal anti-inflammatory drugs	
PDE4 = phosphodiesterase-4	
RIMA = reversible inhibitor of mono-amine oxidase A	
SC1-11 = G protein-coupled receptor	
SDA = Serotonin/Dopamine Antagonist	
SDRI = selective serotonin and dopamine reuptake inhibitor	
SDRI = selective serotonin, nor-adrenaline and dopamine reuptake int	
SNRI = selective serotonin and nor-adrenaline reuptake inhibitor	
SSRI = selective serotonin reuptake inhibitor	
V1B = vasopressin 1B receptor	

184

[illegible]

[illegible]

PHARMAC. GROUP (see overview hereunder)	nr. PH. PROF.	PHARMACOLOGICAL PROFILE	MAIN INDICATIONS	COMPOUND	DOSE RANGE	COMPANY
Enzymatic System	32	ERK activator	Alzheimer's Disease	CPH-1189	50 to 100mg daily	Centaur Pharmaceuticals
	33	GABA agonist	Alzheimer's Disease	Mefenazine	unknown	Danach Sciences, JPI Natemann, BRD
	34	GABA-A agonist	Insomnia	Gabacadol	5 to 20mg daily	Lundbeck
	35	GABA-A modulator	Insomnia	esazepam	2 to 3mg daily	Sepracor
	36	GABA-A modulator	Insomnia	Zapiden MR sustained-release version	10 to 20mg daily	Sanofi-Synthelabo
	37	GABA-A modulator	Insomnia	Indinon	10 to 20mg daily	DOV/Neurocrine
	38	GABA-A modulator	Anxiety	Pagocaine	30mg daily	Indevus
	39	GABA-A modulator	Insomnia	Zaleplon extended-release	10mg daily	Abbott Pharmaceuticals
	40	GABA-A modulator	Anxiety	ISEP-174559		Sepracor
	41	GABA-A modulator	Anxiety, muscular contractions	SL 65,1469		Sanofi-Synthelabo
Inhibitory Amino Acid System	42	GABA-A modulator	Insomnia	CP-730,330 (IGD 96-3)		Neurogen
	43	GABA-A modulator	Insomnia	IGD 96-3		Neurogen
	44	GABA-A modulator	Anxiety	Oxapropion	10 to 60mg daily	DOV
	45	GABA-B agonist	Depression / Anxiety	AVE 7388	15mg daily	Aventis
	46	GABA-B agonist	Parkinson's Disease	GNF	unknown	Angion
	47	GABA-B agonist	Parkinson's Disease	Meperidine	15mg daily	National Institute on Drug Abuse
	48	GABA-B agonist	Anxiety	VN34740		EL Lilly
	49	GABA-B agonist	Depression / Anxiety	R1204		Roche
	50	GABA-B agonist	Depression (paralytic)	Mefenazine	600 to 1200mg daily	Concept
	51	GABA-B agonist	Depression	ORG 34617/4850	unknown	Abbott
Monoaminergic Transmitter Systems	52	GABA-B agonist	Alzheimer's Disease	ABT-238		Abbott
	53	GABA-B agonist	Alzheimer's Disease	ABT-834		Abbott
	54	Paradoxical Substance	Paradoxical Substance	Paradoxical Substance	see formula	Baxter Laboratories
	55	Paradoxical Substance	Paradoxical Substance	Paradoxical Substance	see formula	Procter & Gamble Pharmaceuticals
	56	Paradoxical Substance	Paradoxical Substance	Paradoxical Substance	see formula	Baxter Laboratories
	57	Paradoxical Substance	Paradoxical Substance	Paradoxical Substance	see formula	Procter & Gamble Pharmaceuticals
	58	Paradoxical Substance	Paradoxical Substance	Paradoxical Substance	see formula	Baxter Laboratories
	59	Paradoxical Substance	Paradoxical Substance	Paradoxical Substance	see formula	Procter & Gamble Pharmaceuticals
	60	Paradoxical Substance	Paradoxical Substance	Paradoxical Substance	see formula	Baxter Laboratories
	61	Paradoxical Substance	Paradoxical Substance	Paradoxical Substance	see formula	Procter & Gamble Pharmaceuticals
Monoaminergic Transmitter Systems	62	Paradoxical Substance	Paradoxical Substance	Paradoxical Substance	see formula	Baxter Laboratories
	63	Paradoxical Substance	Paradoxical Substance	Paradoxical Substance	see formula	Procter & Gamble Pharmaceuticals
	64	Paradoxical Substance	Paradoxical Substance	Paradoxical Substance	see formula	Baxter Laboratories
	65	Paradoxical Substance	Paradoxical Substance	Paradoxical Substance	see formula	Procter & Gamble Pharmaceuticals
	66	Paradoxical Substance	Paradoxical Substance	Paradoxical Substance	see formula	Baxter Laboratories
	67	Paradoxical Substance	Paradoxical Substance	Paradoxical Substance	see formula	Procter & Gamble Pharmaceuticals
	68	Paradoxical Substance	Paradoxical Substance	Paradoxical Substance	see formula	Baxter Laboratories
	69	Paradoxical Substance	Paradoxical Substance	Paradoxical Substance	see formula	Procter & Gamble Pharmaceuticals
	70	Paradoxical Substance	Paradoxical Substance	Paradoxical Substance	see formula	Baxter Laboratories
	71	Paradoxical Substance	Paradoxical Substance	Paradoxical Substance	see formula	Procter & Gamble Pharmaceuticals
Monoaminergic Transmitter Systems	72	Paradoxical Substance	Paradoxical Substance	Paradoxical Substance	see formula	Baxter Laboratories
	73	Paradoxical Substance	Paradoxical Substance	Paradoxical Substance	see formula	Procter & Gamble Pharmaceuticals
	74	Paradoxical Substance	Paradoxical Substance	Paradoxical Substance	see formula	Baxter Laboratories
	75	Paradoxical Substance	Paradoxical Substance	Paradoxical Substance	see formula	Procter & Gamble Pharmaceuticals
	76	Paradoxical Substance	Paradoxical Substance	Paradoxical Substance	see formula	Baxter Laboratories
	77	Paradoxical Substance	Paradoxical Substance	Paradoxical Substance	see formula	Procter & Gamble Pharmaceuticals
	78	Paradoxical Substance	Paradoxical Substance	Paradoxical Substance	see formula	Baxter Laboratories
	79	Paradoxical Substance	Paradoxical Substance	Paradoxical Substance	see formula	Procter & Gamble Pharmaceuticals
	80	Paradoxical Substance	Paradoxical Substance	Paradoxical Substance	see formula	Baxter Laboratories
	81	Paradoxical Substance	Paradoxical Substance	Paradoxical Substance	see formula	Procter & Gamble Pharmaceuticals
Monoaminergic Transmitter Systems	82	Paradoxical Substance	Paradoxical Substance	Paradoxical Substance	see formula	Baxter Laboratories
	83	Paradoxical Substance	Paradoxical Substance	Paradoxical Substance	see formula	Procter & Gamble Pharmaceuticals
	84	Paradoxical Substance	Paradoxical Substance	Paradoxical Substance	see formula	Baxter Laboratories
	85	Paradoxical Substance	Paradoxical Substance	Paradoxical Substance	see formula	Procter & Gamble Pharmaceuticals
	86	Paradoxical Substance	Paradoxical Substance	Paradoxical Substance	see formula	Baxter Laboratories
	87	Paradoxical Substance	Paradoxical Substance	Paradoxical Substance	see formula	Procter & Gamble Pharmaceuticals
	88	Paradoxical Substance	Paradoxical Substance	Paradoxical Substance	see formula	Baxter Laboratories
	89	Paradoxical Substance	Paradoxical Substance	Paradoxical Substance	see formula	Procter & Gamble Pharmaceuticals
	90	Paradoxical Substance	Paradoxical Substance	Paradoxical Substance	see formula	Baxter Laboratories
	91	Paradoxical Substance	Paradoxical Substance	Paradoxical Substance	see formula	Procter & Gamble Pharmaceuticals
Monoaminergic Transmitter Systems	92	Paradoxical Substance	Paradoxical Substance	Paradoxical Substance	see formula	Baxter Laboratories
	93	Paradoxical Substance	Paradoxical Substance	Paradoxical Substance	see formula	Procter & Gamble Pharmaceuticals
	94	Paradoxical Substance	Paradoxical Substance	Paradoxical Substance	see formula	Baxter Laboratories
	95	Paradoxical Substance	Paradoxical Substance	Paradoxical Substance	see formula	Procter & Gamble Pharmaceuticals
	96	Paradoxical Substance	Paradoxical Substance	Paradoxical Substance	see formula	Baxter Laboratories
	97	Paradoxical Substance	Paradoxical Substance	Paradoxical Substance	see formula	Procter & Gamble Pharmaceuticals
	98	Paradoxical Substance	Paradoxical Substance	Paradoxical Substance	see formula	Baxter Laboratories
	99	Paradoxical Substance	Paradoxical Substance	Paradoxical Substance	see formula	Procter & Gamble Pharmaceuticals
	100	Paradoxical Substance	Paradoxical Substance	Paradoxical Substance	see formula	Baxter Laboratories
	101	Paradoxical Substance	Paradoxical Substance	Paradoxical Substance	see formula	Procter & Gamble Pharmaceuticals

PHARMAC. GROUP (see overview hereunder)	nr.	PH. PROF.	PHARMACOLOGICAL PROFILE	MAIN INDICATIONS	COMPOUND	DOSE RANGE	COMPANY
Monoaminergic Transmitter Systems							
Monoaminergic Transmitter Systems	60	NASDA	Insomnia		ORG 4420	unknown	Organon
Monoaminergic Transmitter Systems	61	NADRI	Depression (bipolar disorder)		GW395762	20 to 80mg daily	Gilead Sciences
Neurotransmitter System	62	PASOD	Parkinson's Disease		GP 1485	200 to 1000mg daily	Gallard Pharmaceuticals
Amino Acid Transmitter System	63	PARMODOL	Schizophrenia		salsoline		Schering-Plough
Peptide/Tyrosine Transmitter System	64	PRNMD	Alzheimer's Disease		SR 4862	90 to 300mg daily	Sandoz-Synthelabo
Neurotrophic System	65	PRNMD	new growth factor (NGF) gene therapy		SEP 14539	unknown	Cengen
Excitatory Amino Acid System	66	PRNMD	Anxiety		ABT 489	4 to 40mg daily	Abbott
Excitatory Amino Acid System	67	PRNMD	Alzheimer's Disease		serendipin	100 mg daily	Sandoz-Synthelabo
Peptide/Tyrosine Transmitter System	68	PRNMD	Depression / Anxiety		isozaxat		Gilead Sciences
Peptide/Tyrosine Transmitter System	69	PRNMD	Schizophrenia		Lactant	6mg daily	Sandoz
Excitatory Amino Acid System	70	PRNMD	Schizophrenia / BSS / Overactive bladder		SEP 14539	20mg daily	Lundbeck / Forest Laboratories
Excitatory Amino Acid System	71	PRNMD	Alzheimer's Disease		mennamine	20mg daily	Lundbeck / Forest Laboratories
Excitatory Amino Acid System	72	PRNMD	Depression		mennamine	20mg daily	Lundbeck / Forest Laboratories
Excitatory Amino Acid System	73	PRNMD	Pain		mennamine	20mg daily	Lundbeck / Forest Laboratories
Enzymatic System	74	NSAID	Depression		Deleucopine		NPS
Enzymatic System	75	NSAID	Pain		prolopan		Biochemie-Hyphem Pharmaceuticals
Enzymatic System	76	NSAID	Alzheimer's Disease		Furizan (pure Renalizer form of Furiproden)	unknown	Myriad Genetics
Enzymatic System	77	NSAID	Alzheimer's Disease		JAX-1094	132 to 84mg	Medix
Enzymatic System	78	NSAID	Alzheimer's Disease		nallone depot	unknown	Drug Abuse Sciences
Enzymatic System	79	NSAID	Alzheimer's Disease		depo nallone microcapsules	unknown	Boehringer-Ingelheim
Enzymatic System	80	NSAID	Alzheimer's Disease		Seraxina	unknown	Lundbeck / Forest Laboratories
Enzymatic System	81	NSAID	Alzheimer's Disease		pyrazochne		Ester
Enzymatic System	82	NSAID	Alzheimer's Disease		pyrazochne		Ester
Enzymatic System	83	NSAID	Alzheimer's Disease		pyrazochne		Ester
Enzymatic System	84	NSAID	Alzheimer's Disease		pyrazochne		Ester
Enzymatic System	85	NSAID	Alzheimer's Disease		pyrazochne		Ester
Enzymatic System	86	NSAID	Alzheimer's Disease		pyrazochne		Ester
Enzymatic System	87	NSAID	Alzheimer's Disease		pyrazochne		Ester
Enzymatic System	88	NSAID	Alzheimer's Disease		pyrazochne		Ester
Enzymatic System	89	NSAID	Alzheimer's Disease		pyrazochne		Ester
Enzymatic System	90	NSAID	Alzheimer's Disease		pyrazochne		Ester
Enzymatic System	91	NSAID	Alzheimer's Disease		pyrazochne		Ester
Enzymatic System	92	NSAID	Alzheimer's Disease		pyrazochne		Ester
Enzymatic System	93	NSAID	Alzheimer's Disease		pyrazochne		Ester
Enzymatic System	94	NSAID	Alzheimer's Disease		pyrazochne		Ester
Enzymatic System	95	NSAID	Alzheimer's Disease		pyrazochne		Ester
Enzymatic System	96	NSAID	Alzheimer's Disease		pyrazochne		Ester
Enzymatic System	97	NSAID	Alzheimer's Disease		pyrazochne		Ester
Enzymatic System	98	NSAID	Alzheimer's Disease		pyrazochne		Ester
Enzymatic System	99	NSAID	Alzheimer's Disease		pyrazochne		Ester
Enzymatic System	100	NSAID	Alzheimer's Disease		pyrazochne		Ester
Enzymatic System	101	NSAID	Alzheimer's Disease		pyrazochne		Ester
Enzymatic System	102	NSAID	Alzheimer's Disease		pyrazochne		Ester
Enzymatic System	103	NSAID	Alzheimer's Disease		pyrazochne		Ester
Enzymatic System	104	NSAID	Alzheimer's Disease		pyrazochne		Ester
Enzymatic System	105	NSAID	Alzheimer's Disease		pyrazochne		Ester
Enzymatic System	106	NSAID	Alzheimer's Disease		pyrazochne		Ester
Enzymatic System	107	NSAID	Alzheimer's Disease		pyrazochne		Ester
Enzymatic System	108	NSAID	Alzheimer's Disease		pyrazochne		Ester
Enzymatic System	109	NSAID	Alzheimer's Disease		pyrazochne		Ester
Enzymatic System	110	NSAID	Alzheimer's Disease		pyrazochne		Ester
Enzymatic System	111	NSAID	Alzheimer's Disease		pyrazochne		Ester
Enzymatic System	112	NSAID	Alzheimer's Disease		pyrazochne		Ester
Enzymatic System	113	NSAID	Alzheimer's Disease		pyrazochne		Ester
Enzymatic System	114	NSAID	Alzheimer's Disease		pyrazochne		Ester
Enzymatic System	115	NSAID	Alzheimer's Disease		pyrazochne		Ester
Enzymatic System	116	NSAID	Alzheimer's Disease		pyrazochne		Ester
Enzymatic System	117	NSAID	Alzheimer's Disease		pyrazochne		Ester
Enzymatic System	118	NSAID	Alzheimer's Disease		pyrazochne		Ester
Enzymatic System	119	NSAID	Alzheimer's Disease		pyrazochne		Ester
Enzymatic System	120	NSAID	Alzheimer's Disease		pyrazochne		Ester
Enzymatic System	121	NSAID	Alzheimer's Disease		pyrazochne		Ester
Enzymatic System	122	NSAID	Alzheimer's Disease		pyrazochne		Ester
Enzymatic System	123	NSAID	Alzheimer's Disease		pyrazochne		Ester
Enzymatic System	124	NSAID	Alzheimer's Disease		pyrazochne		Ester
Enzymatic System	125	NSAID	Alzheimer's Disease		pyrazochne		Ester
Enzymatic System	126	NSAID	Alzheimer's Disease		pyrazochne		Ester
Enzymatic System	127	NSAID	Alzheimer's Disease		pyrazochne		Ester
Enzymatic System	128	NSAID	Alzheimer's Disease		pyrazochne		Ester
Enzymatic System	129	NSAID	Alzheimer's Disease		pyrazochne		Ester
Enzymatic System	130	NSAID	Alzheimer's Disease		pyrazochne		Ester
Enzymatic System	131	NSAID	Alzheimer's Disease		pyrazochne		Ester
Enzymatic System	132	NSAID	Alzheimer's Disease		pyrazochne		Ester
Enzymatic System	133	NSAID	Alzheimer's Disease		pyrazochne		Ester
Enzymatic System	134	NSAID	Alzheimer's Disease		pyrazochne		Ester
Enzymatic System	135	NSAID	Alzheimer's Disease		pyrazochne		Ester
Enzymatic System	136	NSAID	Alzheimer's Disease		pyrazochne		Ester
Enzymatic System	137	NSAID	Alzheimer's Disease		pyrazochne		Ester
Enzymatic System	138	NSAID	Alzheimer's Disease		pyrazochne		Ester
Enzymatic System	139	NSAID	Alzheimer's Disease		pyrazochne		Ester
Enzymatic System	140	NSAID	Alzheimer's Disease		pyrazochne		Ester
Enzymatic System	141	NSAID	Alzheimer's Disease		pyrazochne		Ester
Enzymatic System	142	NSAID	Alzheimer's Disease		pyrazochne		Ester
Enzymatic System	143	NSAID	Alzheimer's Disease		pyrazochne		Ester
Enzymatic System	144	NSAID	Alzheimer's Disease		pyrazochne		Ester
Enzymatic System	145	NSAID	Alzheimer's Disease		pyrazochne		Ester
Enzymatic System	146	NSAID	Alzheimer's Disease		pyrazochne		Ester
Enzymatic System	147	NSAID	Alzheimer's Disease		pyrazochne		Ester
Enzymatic System	148	NSAID	Alzheimer's Disease		pyrazochne		Ester
Enzymatic System	149	NSAID	Alzheimer's Disease		pyrazochne		Ester
Enzymatic System	150	NSAID	Alzheimer's Disease		pyrazochne		Ester
Enzymatic System	151	NSAID	Alzheimer's Disease		pyrazochne		Ester
Enzymatic System	152	NSAID	Alzheimer's Disease		pyrazochne		Ester
Enzymatic System	153	NSAID	Alzheimer's Disease		pyrazochne		Ester
Enzymatic System	154	NSAID	Alzheimer's Disease		pyrazochne		Ester
Enzymatic System	155	NSAID	Alzheimer's Disease		pyrazochne		Ester
Enzymatic System	156	NSAID	Alzheimer's Disease		pyrazochne		Ester
Enzymatic System	157	NSAID	Alzheimer's Disease		pyrazochne		Ester
Enzymatic System	158	NSAID	Alzheimer's Disease		pyrazochne		Ester
Enzymatic System	159	NSAID	Alzheimer's Disease		pyrazochne		Ester
Enzymatic System	160	NSAID	Alzheimer's Disease		pyrazochne		Ester
Enzymatic System	161	NSAID	Alzheimer's Disease		pyrazochne		Ester
Enzymatic System	162	NSAID	Alzheimer's Disease		pyrazochne		Ester
Enzymatic System	163	NSAID	Alzheimer's Disease		pyrazochne		Ester
Enzymatic System	164	NSAID	Alzheimer's Disease		pyrazochne		Ester
Enzymatic System	165	NSAID	Alzheimer's Disease		pyrazochne		Ester
Enzymatic System	166	NSAID	Alzheimer's Disease		pyrazochne		Ester
Enzymatic System	167	NSAID	Alzheimer's Disease		pyrazochne		Ester
Enzymatic System	168	NSAID	Alzheimer's Disease		pyrazochne		Ester
Enzymatic System	169	NSAID	Alzheimer's Disease		pyrazochne		Ester
Enzymatic System	170	NSAID	Alzheimer's Disease		pyrazochne		Ester
Enzymatic System	171	NSAID	Alzheimer's Disease		pyrazochne		Ester
Enzymatic System	172	NSAID	Alzheimer's Disease		pyrazochne		Ester
Enzymatic System	173	NSAID	Alzheimer's Disease		pyrazochne		Ester
Enzymatic System	174	NSAID	Alzheimer's Disease		pyrazochne		Ester
Enzymatic System	175	NSAID	Alzheimer's Disease		pyrazochne		Ester
Enzymatic System	176	NSAID	Alzheimer's Disease		pyrazochne		Ester
Enzymatic System	177	NSAID	Alzheimer's Disease		pyrazochne		Ester
Enzymatic System	178	NSAID	Alzheimer's Disease		pyrazochne		Ester
Enzymatic System	179	NSAID	Alzheimer's Disease		pyrazochne		Ester
Enzymatic System	180	NSAID	Alzheimer's Disease		pyrazochne		Ester
Enzymatic System	181	NSAID	Alzheimer's Disease		pyrazochne		Ester
Enzymatic System	182	NSAID	Alzheimer's Disease		pyrazochne		Ester
Enzymatic System	183	NSAID	Alzheimer's Disease		pyrazochne		Ester
Enzymatic System	184	NSAID	Alzheimer's Disease		pyrazochne		Ester
Enzymatic System	185	NSAID	Alzheimer's Disease		pyrazochne		Ester
Enzymatic System	186	NSAID	Alzheimer's Disease		pyrazochne		Ester
Enzymatic System	187	NSAID	Alzheimer's Disease		pyrazochne		Ester
Enzymatic System	188	NSAID	Alzheimer's Disease		pyrazochne		Ester
Enzymatic System	189	NSAID	Alzheimer's Disease		pyrazochne		Ester
Enzymatic System	190	NSAID	Alzheimer's Disease		pyrazochne		Ester
Enzymatic System	191	NSAID	Alzheimer's Disease		pyrazochne		Ester
Enzymatic System	192	NSAID	Alzheimer's Disease		pyrazochne		Ester
Enzymatic System	193	NSAID	Alzheimer's Disease		pyrazochne		Ester
Enzymatic System	194	NSAID	Alzheimer's Disease		pyrazochne		Ester
Enzymatic System	195	NSAID	Alzheimer's Disease		pyrazochne		Ester
Enzymatic System	196	NSAID	Alzheimer's Disease		pyrazochne		Ester
Enzymatic System	197	NSAID	Alzheimer's Disease		pyrazochne		Ester
Enzymatic System	198	NSAID	Alzheimer's Disease		pyrazochne		Ester
Enzymatic System	199	NSAID	Alzheimer's Disease		pyrazochne		Ester
Enzymatic System	200	NSAID	Alzheimer's Disease		pyrazochne		Ester
Enzymatic System	201	NSAID	Alzheimer's Disease		pyrazochne		Ester
Enzymatic System	202	NSAID	Alzheimer's Disease		pyrazochne		Ester
Enzymatic System	203	NSAID	Alzheimer's Disease		pyrazochne		Ester
Enzymatic System	204	NSAID	Alzheimer's Disease		pyrazochne		Ester
Enzymatic System	205	NSAID	Alzheimer's Disease		pyrazochne		Ester
Enzymatic System	206	NSAID	Alzheimer's Disease		pyrazochne		Ester
Enzymatic System	207	NSAID	Alzheimer's Disease		pyrazochne		Ester
Enzymatic System	208	NSAID	Alzheimer's Disease		pyrazochne		Ester
Enzymatic System	209	NSAID	Alzheimer's Disease		pyrazochne		Ester
Enzymatic System	210	NSAID	Alzheimer's Disease		pyrazochne		Ester
Enzymatic System	211	NSAID	Alzheimer's Disease		pyrazochne		Ester
Enzymatic System	212	NSAID	Alzheimer's Disease		pyrazochne		Ester
Enzymatic System	213	NSAID	Alzheimer's Disease		pyrazochne		Ester
Enzymatic System	214	NSAID	Alzheimer's Disease		pyrazochne		Ester
Enzymatic System	215	NSAID	Alzheimer's Disease		pyrazochne		Ester
Enzymatic System	216	NSAID	Alzheimer's Disease		pyrazochne		Ester
Enzymatic System	217	NSAID	Alzheimer's Disease		pyrazochne		Ester
Enzymatic System	218	NSAID	Alzheimer's Disease		pyrazochne		Ester
Enzymatic System	219	NSAID	Alzheimer's Disease		pyrazochne		Ester
Enzymatic System	220	NSAID	Alzheimer's Disease		pyrazochne		Ester
Enzymatic System	221	NSAID	Alzheimer's Disease		pyrazochne		Ester
Enzymatic System	222	NSAID	Alzheimer's Disease		pyrazochne		Ester
Enzymatic System	223	NSAID	Alzheimer's Disease		pyrazochne		Ester
Enzymatic System	224	NSAID	Alzheimer's Disease		pyrazochne		Ester
Enzymatic System	225	NSAID	Alzheimer's Disease		pyrazochne		Ester
Enzymatic System	226	NSAID	Alzheimer's Disease		pyrazochne		Ester
Enzymatic System	227	NSAID	Alzheimer's Disease		pyrazochne		Ester
Enzymatic System	228	NSAID	Alzheimer's Disease		pyrazochne		Ester
Enzymatic System	229	NSAID	Alzheimer's Disease		pyrazochne		Ester
Enzymatic System	230	NSAID	Alzheimer's Disease		pyrazochne		Ester
Enzymatic System	231	NSAID	Alzheimer's Disease		pyrazochne		Ester
Enzymatic System	232	NSAID	Alzheimer's Disease		pyrazochne		Ester
Enzymatic System	233	NSAID	Alzheimer's Disease		pyrazochne		Ester
Enzymatic System	234	NSAID	Alzheimer's Disease		pyrazochne		Ester
Enzymatic System	235	NSAID	Alzheimer's Disease		pyrazochne		Ester
Enzymatic System	236	NSAID	Alzheimer's Disease		pyrazochne		Ester
Enzymatic System	237	NSAID	Alzheimer's Disease		pyrazochne		Ester
Enzymatic System	238	NSAID	Alzheimer's Disease		pyrazochne		Ester
Enzymatic System	239	NSAID	Alzheimer's Disease		pyrazochne		Ester
Enzymatic System	240	NSAID	Alzheimer's Disease		pyrazochne		Ester
Enzymatic System	241	NSAID	Alzheimer's Disease		pyrazochne		Ester
Enzymatic System	242	NSAID	Alzheimer's Disease		pyrazochne		Ester
Enzymatic System	243	NSAID	Alzheimer's Disease		pyrazochne		Ester
Enzymatic System	244	NSAID	Alzheimer's Disease		pyrazochne		Ester
Enzymatic System	245	NSAID	Alzheimer's Disease		pyrazochne		Ester
Enzymatic System	246	NSAID	Alzheimer's Disease		pyrazochne		Ester
Enzymatic System	247	NSAID	Alzheimer's Disease		pyrazochne		Ester
Enzymatic System	248	NSAID	Alzheimer's Disease		pyrazochne		Ester
Enzymatic System	249	NSAID	Alzheimer's Disease		pyrazochne		Ester
Enzymatic System	250	NSAID	Alzheimer's Disease		pyrazochne		Ester
Enzymatic System	251	NSAID	Alzheimer's Disease		pyrazochne		Ester
Enzymatic System	252	NSAID	Alzheimer's Disease		pyrazochne		Ester
Enzymatic System	253	NSAID	Alzheimer's Disease		pyrazochne		Ester
Enzymatic System	254	NSAID	Alzheimer's Disease		pyrazochne		Ester
Enzymatic System	255	NSAID	Alzheimer's Disease		pyrazochne		Ester
Enzymatic System	256	NSAID	Alzheimer's Disease		pyrazochne		Ester
Enzymatic System	257	NSAID	Alzheimer's Disease		pyrazochne		Ester
Enzymatic System	258						

[illegible]

PHARMAC. GROUP (see overview hereunder)	nr. PH. PROF.	PHARMACOLOGICAL PROFILE	MAIN INDICATIONS	COMPOUND	DOSE RANGE	COMPANY
Noncompetitive Transmitter Systems	92	SSRI	Depression / Anxiety	paroxetine	10 to 20mg daily	GlaxoSmithKline
	93	SSRI	Depression / Anxiety	escitalopram	10 to 20mg daily	Lundbeck / Forest Laboratories
	94	SSRI	Depression / Anxiety	clomipramine	10 to 40 mg daily	off patent
Monoaminergic Transmitter Systems	95	SSRI	Depression / Anxiety	fluoxetine	10 to 40 mg daily	off patent
	96	SSRI	Depression / Anxiety	fluoxetine	10 to 40 mg daily	off patent
	97	SSRI	Depression / Anxiety	fluoxetine	10 to 40 mg daily	off patent
Peptidergic Transmitter Systems	98	SSRI	Depression / Anxiety	fluoxetine	10 to 40 mg daily	off patent
	99	SSRI	Depression / Anxiety	fluoxetine	10 to 40 mg daily	off patent
	100	SSRI	Depression / Anxiety	fluoxetine	10 to 40 mg daily	off patent
Adrenergic Transmitter Systems	101	SSRI	Depression / Anxiety	fluoxetine	10 to 40 mg daily	off patent
	102	SSRI	Depression / Anxiety	fluoxetine	10 to 40 mg daily	off patent
	103	SSRI	Depression / Anxiety	fluoxetine	10 to 40 mg daily	off patent
Endocrine System	104	SSRI	Depression / Anxiety	fluoxetine	10 to 40 mg daily	off patent
	105	SSRI	Depression / Anxiety	fluoxetine	10 to 40 mg daily	off patent
	106	SSRI	Depression / Anxiety	fluoxetine	10 to 40 mg daily	off patent
Neurotrophic System	107	SSRI	Depression / Anxiety	fluoxetine	10 to 40 mg daily	off patent
	108	SSRI	Depression / Anxiety	fluoxetine	10 to 40 mg daily	off patent
	109	SSRI	Depression / Anxiety	fluoxetine	10 to 40 mg daily	off patent
Nerve Cell Function System	110	SSRI	Depression / Anxiety	fluoxetine	10 to 40 mg daily	off patent
	111	SSRI	Depression / Anxiety	fluoxetine	10 to 40 mg daily	off patent
	112	SSRI	Depression / Anxiety	fluoxetine	10 to 40 mg daily	off patent
Pathogenic Mechanisms of Dementia of the Alzheimer Type	113	SSRI	Depression / Anxiety	fluoxetine	10 to 40 mg daily	off patent
	114	SSRI	Depression / Anxiety	fluoxetine	10 to 40 mg daily	off patent
	115	SSRI	Depression / Anxiety	fluoxetine	10 to 40 mg daily	off patent
Other / Unknown Systems	116	SSRI	Depression / Anxiety	fluoxetine	10 to 40 mg daily	off patent
	117	SSRI	Depression / Anxiety	fluoxetine	10 to 40 mg daily	off patent
	118	SSRI	Depression / Anxiety	fluoxetine	10 to 40 mg daily	off patent
PHARMACOLOGICAL GROUPS: COMPOUNDS WORKING ON THE	119	SSRI	Depression / Anxiety	fluoxetine	10 to 40 mg daily	off patent
	120	SSRI	Depression / Anxiety	fluoxetine	10 to 40 mg daily	off patent
	121	SSRI	Depression / Anxiety	fluoxetine	10 to 40 mg daily	off patent
Amino Acid Transmitter Systems	122	SSRI	Depression / Anxiety	fluoxetine	10 to 40 mg daily	off patent
	123	SSRI	Depression / Anxiety	fluoxetine	10 to 40 mg daily	off patent
	124	SSRI	Depression / Anxiety	fluoxetine	10 to 40 mg daily	off patent
Monoaminergic Transmitter Systems	125	SSRI	Depression / Anxiety	fluoxetine	10 to 40 mg daily	off patent
	126	SSRI	Depression / Anxiety	fluoxetine	10 to 40 mg daily	off patent
	127	SSRI	Depression / Anxiety	fluoxetine	10 to 40 mg daily	off patent
Excitatory Amino Acid System	128	SSRI	Depression / Anxiety	fluoxetine	10 to 40 mg daily	off patent
	129	SSRI	Depression / Anxiety	fluoxetine	10 to 40 mg daily	off patent
	130	SSRI	Depression / Anxiety	fluoxetine	10 to 40 mg daily	off patent
Inhibitory Amino Acid System	131	SSRI	Depression / Anxiety	fluoxetine	10 to 40 mg daily	off patent
	132	SSRI	Depression / Anxiety	fluoxetine	10 to 40 mg daily	off patent
	133	SSRI	Depression / Anxiety	fluoxetine	10 to 40 mg daily	off patent
Peptidergic Transmitter Systems	134	SSRI	Depression / Anxiety	fluoxetine	10 to 40 mg daily	off patent
	135	SSRI	Depression / Anxiety	fluoxetine	10 to 40 mg daily	off patent
	136	SSRI	Depression / Anxiety	fluoxetine	10 to 40 mg daily	off patent
Adrenergic Transmitter Systems	137	SSRI	Depression / Anxiety	fluoxetine	10 to 40 mg daily	off patent
	138	SSRI	Depression / Anxiety	fluoxetine	10 to 40 mg daily	off patent
	139	SSRI	Depression / Anxiety	fluoxetine	10 to 40 mg daily	off patent
Endocrine System	140	SSRI	Depression / Anxiety	fluoxetine	10 to 40 mg daily	off patent
	141	SSRI	Depression / Anxiety	fluoxetine	10 to 40 mg daily	off patent
	142	SSRI	Depression / Anxiety	fluoxetine	10 to 40 mg daily	off patent
Neurotrophic System	143	SSRI	Depression / Anxiety	fluoxetine	10 to 40 mg daily	off patent
	144	SSRI	Depression / Anxiety	fluoxetine	10 to 40 mg daily	off patent
	145	SSRI	Depression / Anxiety	fluoxetine	10 to 40 mg daily	off patent
Nerve Cell Function System	146	SSRI	Depression / Anxiety	fluoxetine	10 to 40 mg daily	off patent
	147	SSRI	Depression / Anxiety	fluoxetine	10 to 40 mg daily	off patent
	148	SSRI	Depression / Anxiety	fluoxetine	10 to 40 mg daily	off patent
Pathogenic Mechanisms of Dementia of the Alzheimer Type	149	SSRI	Depression / Anxiety	fluoxetine	10 to 40 mg daily	off patent
	150	SSRI	Depression / Anxiety	fluoxetine	10 to 40 mg daily	off patent
	151	SSRI	Depression / Anxiety	fluoxetine	10 to 40 mg daily	off patent
Other / Unknown Systems	152	SSRI	Depression / Anxiety	fluoxetine	10 to 40 mg daily	off patent
	153	SSRI	Depression / Anxiety	fluoxetine	10 to 40 mg daily	off patent
	154	SSRI	Depression / Anxiety	fluoxetine	10 to 40 mg daily	off patent

Examples

Example 1: Measuring pKi values of test compounds

- 5 In Table 1, the pKi values of test compounds are given for each of the dopamine receptors, 5HT receptors, adrenergic receptors and the histamine1 receptor. The affinity of test compounds for the respective receptors has been performed according to conventional procedures known in the art.
- An indication "0" means that no affinity has been measured between the test compound and
- 10 the receptor.
- The columns displaying the pKi values for the D4 and the 5-HT2A receptor are filled with dark grey. pKi values between 8 and 9 and higher than 9 are represented by light grey boxes.

15 Example 2: Foregoing pipamperon-citalopram treatment in major depressive disorder: a placebo and active controlled period finding clinical trial

- Table 2 represents the set-up of a clinical trial comprising for treatment groups:
- Group Plc – Active / Day 0 represents the group receiving 10 mg citalopram, twice a day, starting the first day (Day 0) of active treatment in the clinical trial. This administration regime
- 20 is also indicated as the mono therapy.
- Group Pip - Active / Day 0 represents the group receiving a combination of 4 mg pipamperon and 10 mg citalopram, twice a day, starting the first day (Day 0) of active treatment in the clinical trial. This administration regime is also indicated as the non-foregoing combo therapy.
- 25 Group Pip - Active / Day 4 represents the group receiving 4 mg pipamperon, twice a day, starting the first day (Day 0) of active treatment in the clinical trial, followed by a combination of 4 mg pipamperon and 10 mg citalopram, twice a day, starting the fifth (Day 4) day of active treatment in the clinical trial. This administration regime is also indicated as the foregoing therapy with combination therapy starting after 4 days of active treatment.
- 30 Group Pip - Active / Day 7 represents the group receiving 4 mg pipamperon, twice a day, starting the first day (Day 0) of active treatment in the clinical trial, followed by a combination of 4 mg pipamperon and 10 mg citalopram, twice a day, starting the eight (Day 7) day of

active treatment in the clinical trial. This administration regime is also indicated as the foregoing therapy with combination therapy starting after 7 days of active treatment.

All subjects also undergo a placebo (PLC) run-in therapy, administered during a period of about 7 days before the active treatment starts.

5 During daily (D), weekly (W) or monthly (M) visits, several parameters are measured.

Under NECT is to be understood: Neuronal E-clinical Trial = Vesalius Expert development for this trial which includes the bottom-up measurement of:

- In- and exclusion-criteria
- Functional status evaluation
- 10 - Medical history
- (Pre-)treatment signs & symptoms
- DSM-IV rules for diagnosis & efficacy
- HDRS-28 (Hamilton Depression Rating Scale - 28 items)
- Medical resource utilisation
- 15 - Pre-trial & Concomittant medication
- Drug administration
- (Serious) Adverse events
- Admission to the acute and extension phase of treatment
- Right flow of the trial

20

Example 3: combo pipamperon-citalopram: therapeutic use in Major Depression

25 Purpose

Pipamperon (1'-[3-(p-Fluorobenzoyl)propyl]-[1,4'-bipiperidine]-4'-carboxamide), the active ingredient of Dipiperon (Janssen-Cilag B.V), administered to patients in a dose ranging between 8 and 12 mg is claimed via its specific pharmacological properties to be a booster of the antidepressant effect of the selective serotonin re-uptake inhibitor

30 citalopram. Preferably, pipamperon is administered daily at least 4-5 days before administering said antidepressant. The mechanism of boosting of pipamperon has to deal with (i) the selective affinity for the dopamine-4 (D4) receptor with a pKi value equal to or higher than 8 towards the D4 receptor and less than 8 towards other dopamine receptors, and (ii) the selective affinity for the 5-HT2A receptor with a pKi value equal to or higher

35 than 8 towards the 5-HT2A receptor and less than 8 towards other 5HT receptors. This semi-naturalistic open label study investigated the efficacy and tolerability of the combo pipamperon - citalopram in the treatment of patients with major depression.

Details

Design: Semi-naturalistic i.e. inclusion of every 'natural' patient in an outpatient practice but without concomitant use of mood enhancing drugs, open label

Control: No

Phase: Phase IIa – preliminary Proof of Concept

Location: Belgium – Research Centre ANIMA, Alken

End Points : Assessment scale scores, Hamilton Depression Rating Scale 17 items, Reduction, Response, Remission

Medication: Exclusion of mood stabilisers, antipsychotics (typical and atypical) and other antidepressants

Subjects

Type	No.	Sex	Age
Patients	23	10 male & 13 female	23-80 (mean 47) years

- 5 Characteristics: patients had a major depressive disorder according to DSM-IV criteria, with or without a chronic course and a treatment refractory state towards another SSRI then citalopram.

Treatments

- 10 PIP-CIT¹ add-on: citalopram from day minus 60-20 – pipamperon from DAY 0

Drug/Treatment	Dose	Route	Frequency	Duration
Pipamperon ¹	+ Pip.: 8-12 mg/day – Cit.: 20-	PO	bid	8 weeks
Citalopram ¹	40 mg/day			

1. Pipamperon (Pip) and citalopram (Cit) dosage was adjusted according to clinical response.

15

PIP-CIT¹ fore-going 1-5: pipamperon from day 0 – cital from day 1-5

Drug/Treatment	Dose	Route	Frequency	Duration
Pipamperon ¹	+ Pip.: 8-12 mg/day – Cit.: 20-	PO	bid	8 weeks
Citalopram ¹	40 mg/day			

20 1. Pipamperon (Pip) and citalopram (Cit) dosage was adjusted according to clinical response.

20

PIP-CIT¹ fore-going 6-8: pipamperon from day 0 – citalopram from day 6-8

25

193

Drug/Treatment	Dose	Route	Frequency	Duration
Pipamperon ¹	+Pip.: 8-12 mg/day – Cit.: 20-	PO	bid	8 weeks
Citalopram ¹	40 mg/day			

1. Pipamperon (Pip) and citalopram (Cit) dosage was adjusted according to clinical response.

5 Results

		<u>PIP-CIT add-on</u>	<u>PIP-CIT foregoing</u>	
		After 20-60 DAYS (mean 33) (n = 5)	1-5 DAYS (mean 4) (n = 15)	6-8 DAYS (mean 7) (n = 3)
Mean	Used			
<u>Medication</u>				
Pipamperone	9mg/day		10mg/day	11mg/day
Citalopram	30mg/day		26mg/day	30mg/day
<u>Depression scale scores</u>				
HDRS 17-item total score				
baseline	29		23	28
endpoint (week 8)	4		5	11
diminishment at week 8	-25 (+8/-9)		-18 (+8/-8)	-17 (+17/-17)
% reduction at week 8	86 (+14/-12)		80 (+20/-30)	61 (+39/-61)
response ¹ at week 8	5 (100%)		15 (100%)	2 (67%)
remission ² at week 8	4 (80%)		10 (67%)	1 (33%)

1. Response = $\geq 50\%$ reduction in HDRS 17-item score;

2. Remission = HDRS 17-item score < 8

- 10 Notably, the results obtained are highly significant since the variability in every group is distributed evenly around the mean.

Add-on PIP-CIT

- 15 Figure 1 schematically depicts the "add-on" treatment with pipamperon 8-12 (mean 9) mg (bid) after treatment with citalopram 10-20 (mean 30) mg (bid) during 20-60 (mean 33) days (PIPCIT ADD-ON) with HDRS-17. Totalscore is 29 at baseline in MDD in comparison with the standard efficacy of antidepressants in clinical trials according to Khan *et al.* (2000), in "Symptom Reduction and Suicide Risk in Patients Treated With Placebo in Antidepressant Clinical Trials" (Arch. of General Psychiatry, Vol. 57, April 2000).

Figure 2 schematically depicts the HDRS-17 change from baseline in the combo pipamperon as "add-on" to citalopram vs SNRI (duloxetine) in Major Depression. Treatment with pipamperon 8-12 (mean 9 mg/day) during 20-60 (mean 33) days after treatment with SSRI (n=5). The SNRI (duloxetine) treatment was 40-120 mg/day (n = 152) according to Goldstein *et al.*, (Clin. Psychiatry, in press).

Figure 3 schematically depicts the remission rates (HDRS-17 ≤ 7) with the combo pipamperon as "add-on" to citalopram vs SNRI (venlafaxine) vs SSRIs vs placebo in Major Depression. Treatment with pipamperon 8-12 (mean 9 mg/day) during 20-60 (mean 33) days after treatment with SSRI (n=5). Treatment with the SNRI venlafaxine is according to a meta-analysis of Thase *et al.* (Br. J. Psychiatry (2001) 178:234-241). Treatment with SSRIs is according to a meta-analysis of Thase *et al.* (Br. J. Psychiatry (2001) 178:234-241). Treatment with placebo is according to a meta-analysis of Thase *et al.* (Br. J. Psychiatry (2001) 178:234-241).

Fore-going 1-5 PIP-CIT

Figure 4 schematically depicts the "fore-going" treatment during 1-5 (mean 4) days with pipamperon 8-12 (mean 10) mg (bid), followed with the combination treatment of pipamperon and citalopram 20-50 (mean 26) mg/day (bid) (PIPCIT FG 1-5) in MDD (HDRS-17 at BL = 23) in comparison with the standard efficacy of antidepressants in clinical trials according to Khan *et al.* (2000), in "Symptom Reduction and Suicide Risk in Patients Treated With Placebo in Antidepressant Clinical Trials" (Arch. of General Psychiatry, Vol. 57, April 2000).

Figure 5 schematically depicts the HDRS-17 change from baseline in the combo pipamperon-citalopram treatment with a "fore-going" treatment of 4 days with pipamperon (10 mg/day) vs SNRI (duloxetine) in Major Depression. Treatment with the combo pipamperon-citalopram with pipamperon 8-12 (mean 10 mg/day) (bid) 1-5 (mean 4) days before treatment with SSRI (n=15). The SNRI (duloxetine) treatment was 40-120 mg/day (n = 152) according to Goldstein *et al.*, (Clin. Psychiatry, in press).

Figure 6 schematically depicts the remission rates (HDRS-17 ≤ 7) with the combo pipamperon with a "fore-going" treatment of 4 days with pipamperon (10 mg/day) vs SNRI (venlafaxine) in Major Depression. Treatment with the combo pipamperon-citalopram was with pipamperon 8-12 (mean 10 mg/day) during 1-5 (mean 4) days before treatment with the

SSRI (n=5). Treatment with the SNRI venlafaxine is according to a meta-analysis of Thase et al. (Br. J. Psychiatry (2001) 178:234-241). Treatment with SSRIs is according to a meta-analysis of Thase et al. (Br. J. Psychiatry (2001) 178:234-241). Treatment with placebo is according to a meta-analysis of Thase et al. (Br. J. Psychiatry (2001) 178:234-241).

5

Fore-going 6-8 PIP-CIT

Figure 7 schematically depicts the "fore-going" treatment during 6-8 (mean 7) days with pipamperon 8-12 (mean 11) mg/day (bid), followed with the combination treatment of pipamperon and citalopram 20-40 (mean 30) mg/day (bid) (PIPCIT FG 6-8) in MDD (HDRS-17 at BL = 28) in comparison with the standard efficacy of antidepressants in clinical trials according to Khan et al. (2000), in "Symptom Reduction and Suicide Risk in Patients Treated With Placebo in Antidepressant Clinical Trials" (Arch. of General Psychiatry, Vol. 57, April 2000).

10

15

Figure 8 schematically depicts the HDRS-17 change from baseline in the combo pipamperon-citalopram treatment with a "fore-going" treatment of 7 days with pipamperon (11 mg/day) vs SNRI (duloxetine) in Major Depression. Treatment with the combo pipamperon-citalopram with pipamperon 8-12 (mean 11 mg/day) (bid) 6-8 (mean 7) days before treatment with SSRI (n=3). The SNRI (duloxetine) treatment was 40-120 mg/day (n = 152) according to Goldstein et al., (Clin. Psychiatry, in press).

20

Comparison "add-on" vs "fore-going"

Figure 9 schematically depicts a comparison between "fore-going" and "add-on" treatments with pipamperon (8-12 mg/day; bid) and citalopram (20-40 mg/day; bid) in MDD in comparison with the standard efficacy of antidepressants in clinical trials according to Khan et al. (2000), in "Symptom Reduction and Suicide Risk in Patients Treated With Placebo in Antidepressant Clinical Trials" (Arch. of General Psychiatry, Vol. 57, April 2000).

25

Figure 10 schematically depicts a comparison between "fore-going" and "add-on" treatments. In particular, the HDRS-17 change from baseline between "fore-going" and "add-on" treatment with pipamperon (8-12 mg/day; bid) and citalopram (20-40 mg/day; bid) in comparison with the SNRI duloxetine in Major Depression is depicted. Treatment with the combo pipamperon as "add-on" to citalopram, with pipamperon 8-12 mg/day (mean 9 mg/day) 20-60 (mean 33) days after treatment with the SSRI (n=5). Treatment with the combo pipamperon-citalopram, with pipamperon 8-12 mg/day (mean 11 mg/day; bid) 6-8 days (mean 7 days) before treatment with the SSRI (n = 15). Treatment with the combo

30

35

pipamperon-citalopram, with pipamperon 8-12 mg/day (mean 10 mg/day; bid) 1-5 days (mean 4 days) before treatment with the SSRI (n = 15). The SNRI (duloxetine) treatment was 40-120 mg/day (n = 152) according to Goldstein *et al.*, (Clin. Psychiatry, in press).

- 5 Figure 11 schematically depicts the remission rates (HDRS-17 ≤ 7) in a comparison between "fore-going" and "add-on" treatment with pipamperon (8-12 mg/day; bid) and citalopram (20-40 mg/day; bid) in comparison with the SNRI venlafaxine in Major Depression. Treatment with the combo pipamperon-citalopram was with pipamperon 8-12 (mean 10 mg/day) during 1-5 (mean 4) days before treatment with the SSRI (n = 15).
- 10 Treatment with the SNRI venlafaxine is according to a meta-analysis of Thase *et al.* (Br. J. Psychiatry (2001) 178:234-241). Treatment with pipamperon as "add-on" to citalopram, with pipamperon 8-12 (mean 9 mg/day) during 20-60 (mean 33) days after treatment with SSRI (n = 5).
- 15 The intention-to-treat / last-observation-carried-forward analysis showed a high therapeutic efficacy according HDRS 17-item in all the treatment groups. This was especially true for the 'add-on' group probably caused by the longer treatment with an active antidepressant (+33 days). The huge therapeutic effect observed in the 'PIP-CIT 1-5' group present for at a mean dosage of pipamperon of 10 mg per day and administered
- 20 the first four days of treatment without an active antidepressant, indicates the boosting effect of pipamperon on the SSRI citalopram at an extremely and thus unconventional low dose. Only 1 patient discontinued treatment due to a lost of follow-up.

Adverse Events

Side effects (patients)	PIP-CIT	add-on	PIP-CIT foregoing
	After 20-60 DAYS (mean 33) (n = 5)	1-5 DAYS (mean 4) (n = 15)	6-8 DAYS (mean 7) (n = 3)
Discontinued treatment due to adverse events	0	0	0
By system:			
body as a whole	0	0	0
central and peripheral nervous system	1(20%)	4(26.6%)	0
gastrointestinal	1(20%)	5(33%)	2(66.6%)
musculoskeletal	1(20%)	3(20%)	0
psychiatric	0	0	0
respiratory	0	1(6.6%)	0
skin and appendages	1(20%)	2(13.3%)	1(33.3%)
vascular	0	1(6.6%)	0
urinary	0	1(6.6%)	0

Laboratory parameters, ECG, bodyweight and vital signs were not measured since this was a naturalistic study.

5

AssessmentOutcome

Efficacy: the 4-day fore-going combo pipamperon 8-12mg/d - citalopram 20-40mg/day is comparable to the add-on combo pipamperon-citalopram.

10 **Efficacy:** the 4-day fore-going combo pipamperon 8-12mg/d - citalopram 20-40mg/day is larger than the 7-day fore-going combo pipamperon 8-12mg/d - citalopram 20-40mg/day.

Efficacy: the combo pipamperon 8-12mg/d - citalopram 20-40mg/day is larger than the in the art known antidepressants SSRIs.

15 Tolerability

Tolerability: the 4-day fore-going treatment is comparable to the 7-day fore-going combo is comparable to add-on combo pipamperon-citalopram.

Tolerability: no discontinued treatment due to adverse events.

Study Messages

5 The boosting effect of pipamperon at an extremely unconventional low dose on a SSRI is indicated since the efficacy of the 'add-on' and '4-day fore-going' combo 'pipamperon 8 - 12 mg/d - citalopram 20 - 40 mg/day' is in this study as twice higher as known in the art in the treatment of patients with major depression.

10 The combo pipamperon-citalopram is generally well tolerated in patients with depression i.e. at least no specific added adverse events were occurring by adding pipamperon at the doses used in the study.

Example 4: combo pipamperon-citalopram: therapeutic use in Obsessive-Compulsive Disorder (OCD).

Purpose

- 5 Pipamperon (1'-[3-(p-Fluorobenzoyl)propyl]-[1,4'-bipiperidine]-4'-carboxamide), the active ingredient of Dipiperon (Janssen-Cilag B.V), administered to a patient in a dose ranging between 8 and 12 mg is claimed via its specific pharmacological properties to be a booster of the effect of the selective serotonin re-uptake inhibitor citalopram towards OCD. Preferably, pipamperon is administered daily at least 4-5 days before administering
10 said antidepressant. The mechanism of boosting of pipamperon has to deal with (i) the selective affinity for the dopamine-4 (D4) receptor with a pKi value equal to or higher than 8 towards the D4 receptor and less than 8 towards other Dopamine receptors, and (ii) the selective affinity for the 5-HT2A receptor with a pKi value equal to or higher than 8 towards the 5-HT2A receptor and less than 8 towards other 5HT receptors. This semi-
15 naturalistic open label study investigated the efficacy and tolerability of the combo pipamperon - citalopram in the treatment of patients with OCD.

Details

- Design: Semi-naturalistic i.e. inclusion of every 'natural' patient in an outpatient practice but without concomitant use of mood enhancing drugs, open label
- Control: No
- Phase: Phase IIa – preliminary Proof of Concept
- Location: Belgium – Research Centre ANIMA, Alken
- End Points : Assessment scale scores, Yale-Brown Obsessive-Compulsive Scale, Reduction, Remission
- Medication: Exclusion of mood stabilisers, antipsychotics (typical and atypical) and other antidepressants

20 **Subjects**

Type	No.	Sex	Age
Patients	7	1 male & 7 female	20-63 (mean 33) years

Characteristics: patients had an obsessive-compulsive disorder according to DSM-IV criteria, with or without a chronic course and a treatment refractory state towards another SSRI then citalopram.

200

TreatmentsPIP-CIT¹ ADD-ON: citalopram from DAY minus 730-60 – pipamperon from DAY 0

Drug/Treatment	Dose	Route	Frequency	Duration
Pipamperone ¹	+Pip.: 8-16 mg/day – Cit.: 30-80	PO	bid	12 weeks
Citalopram ¹	mg/day			

- 5 1. Pipamperone (Pip) and Citalopram (Cit) dosage was adjusted according to clinical response.

PIP-CIT¹ FORE-GOING 4-6: pipamperon from DAY 0 – citalopram from DAY 4-6

10

Drug/Treatment	Dose	Route	Frequency	Duration
Pipamperone ¹	+Pip.: 8-16 mg/day – Cit.: 30-80	PO	bid	12 weeks
Citalopram ¹	mg/day			

1. Pipamperone (Pip) and Citalopram (Cit) dosage was adjusted according to clinical response.

Results

		<u>PIP-CIT add-on</u>
		after 730-60 DAYS (mean 241) (n = 6)
		with mean Cit. 54mg/d and Pip. 11mg/d
		<u>PIP-CIT foregoing</u>
		4-6 DAYS (mean 5) (n = 2)
		with mean Cit. 60mg/d and Pip. 10mg/d
Y-BOCS score		
<u>Baseline</u>		
Total	31	
Obsessions	18	
Compulsions	13	
Endpoint (week 12)		
<u>Total</u>	<u>15</u>	
diminishment	-16 (+16/-11)	
% reduction	53	
<u>Obsessions</u>		
total	8	
diminishment	-10 (+9/-7)	
% reduction	57	
<u>Compulsions</u>		
total	7	
diminishment	-6 (+7/-6)	
% reduction	45	
<u>% Remission</u>		
YBOCS score ≤8	29	
BOCS score ≤16	57	

Notably, the results obtained are highly significant since the variability in every group is distributed evenly around the mean.

Figure 12 schematically depicts the Y-BOCS total score: "fore-going" and "add-on" treatment with pipamperon (8-15 mg/day; bid) and citalopram (30-80 mg/day; bid) in comparison with the SSRI fluvoxamine in OCD. Treatment with the combo pipamperon-citalopram (n = 7). Treatment with fluvoxamine (controlled release) mean 271 mg/day (n = 253) is according to Hollander et al. (2003).

202

Figure 13 schematically depicts the Y-BOCS obsession score: "fore-going" and "add-on" treatment with pipamperon (8-15 mg/day; bid) and citalopram (30-80 mg/day; bid) in comparison with the SSRI fluvoxamine in OCD. Treatment with the combo pipamperon-citalopram (n = 7). Treatment with fluvoxamine (controlled release) mean 271 mg/day (n = 253) is according to Hollander et al. (2003).

Figure 14 schematically depicts the Y-BOCS compulsion score: "fore-going" and "add-on" treatment with pipamperon (8-16 mg/day; bid) and citalopram (30-80 mg/day; bid) in comparison with the SSRI fluvoxamine in OCD. Treatment with the combo pipamperon-citalopram (n = 7). Treatment with fluvoxamine (controlled release) mean 271 mg/day (n = 253) is according to Hollander et al. (2003).

The intention-to-treat / last-observation-carried-forward analysis showed a high therapeutic efficacy according Y-BOCS total score, obsession and compulsion scores. This indicates the boosting effect of pipamperon on the SSRI citalopram at an extremely and thus unconventional low dose. No patient discontinued treatment.

Assessment

Efficacy: the combo pipamperone 8-16mg/d - citalopram 30-80mg/day > the in the art known compounds effective towards OCD (Hollander E, Koran LM, Goodman WK, Greist JH, Ninan PT, et al. A double-blind, placebo-controlled study of the efficacy and safety of controlled-release fluvoxamine in patients with obsessive-compulsive disorder. Journal of Clinical Psychiatry 64: 640-647, Jun 2003 Mount Sinai School of Medicine, New York, New York, USA; Solvay Pharmaceuticals Inc., Marietta, Georgia, USA).

Study Messages

The boosting effect of pipamperon at an extremely unconventional low dose on a SSRI is indicated since the efficacy of the 'add-on' and 'fore-going' combo 'pipamperon 8-15 mg/d - citalopram 30-80 mg/day' is in this study as twice higher as known in the art in the treatment of patients with obsessive-compulsive disorder.

Example 5: combo pipamperon-citalopram: therapeutic use in Panic Disorder.**Purpose**

Preliminary examination of a "fore-going" and "add-on" treatment with pipamperon and citalopram in comparison with the SSRI in Panic Disorder.

Results

The results are indicated in Figure 15. Figure 15 schematically depicts the CGI-severity score: "fore-going" and "add-on" treatment with pipamperon (8 mg/day; bid) and citalopram (20-40 mg/day; bid) in comparison with the SSRI in Panic Disorder. Treatment with the combo pipamperon-citalopram (n = 3). Treatment with paroxetine is according to the Journal of Clinical Psychiatry (2004) 65: 405-413. Treatment with Sertraline is according to the Journal of Clinical Psychiatry (2004) 65: 405-413.

Conclusion

Notably, although a small test group has been used (n = 3), the distribution around the mean is good. It will further be apparent from Figure 15 that the effect of the combo treatment of pipamperon and citalopram is twice as high as the standard treatments with paroxetine or sertraline.

Example 6: POC process for mayor depressive disorder

Concept: Combo of the high selective 5-HT_{2A/D4} antagonist pipamperon with:

- a compound active towards the Amino Acid Transmitter, Peptidergic Transmitter, Adenosine Transmitter, Endocrine and/ or Enzymatic System;
- a fore-going admission during 4 days of pipamperon;
- a dose of pipamperon of 12 mg/day

Objectives: Demonstrating that this combo therapy has:

- the potency of being a treatment standard for depression by having an added value of reducing the total score of the Hamilton Depression Rating Scale – 17 items (HDRS-17) after 8 weeks of therapy with a least 20% more than reached with the conventional known antidepressants, i.e. 60% versus 40%. This stands for an added medium demission of 5 points on the total score of the HDRS-17 and by this will be very highly

significant since the mean difference in all recent clinical trials between placebo and active treatment is 2.5;

- a more sustained therapeutic effect than the conventional mono therapy by preventing significant more relapses during 48 weeks following the acute treatment; and/or
- 5 - a complete neutral safety profile, e.g. there are no more adverse events in the combo therapy than in mono admission of the in the combo used antidepressant compound.

Process: the following different steps were implemented to reach out for these objectives (see also Tables 3 and 4):

- 10 (1) an naturalistic open label study ($n \geq 20$) on a depressive population with a normal variability of medical and psychiatric history, course of depression, earlier and concomitant therapy admitting the golden standard antidepressant citalopram 20-40 mg/day and a dose of 8-12 mg/day of pipamperon in a foregoing, simultaneous or add-on use.

- 15 (2) a 16 weeks placebo controlled randomised four armed study of each 36 patients with a mayor depressive disorder admitting:

- from day 0: placebo or pipamperon (PIP) 10 mg/day or an active antidepressant compound or the combination of the last two;
- from day 4: placebo or pipamperon 10 mg/day combined with an active antidepressant compound or an active antidepressant compound without pipamperon.
- 20

By including rigorous control groups (placebo and active comparator; see Tables 3 and 4) this clinical trial is evaluated as a proof of concept of the added value of the combo and the foregoing treatment method since the inclusion/exclusion of:

- 25 - a negative trial, i.e. no significant difference between the placebo and active treatment with the comparator;
- a failed trial, i.e. no significant difference between the active and the studied treatment i.e. the combo.

- 30 (3) an active controlled randomised relapse prevention study following the POC trial during another 36 weeks with three arms of each 36 patients which is formed by:

- continuation of the active mono therapy;
- randomising the patients with a combo therapy in a group with an active mono therapy and with a continuation of the combo treatment.

Claims

1. Use of pipamperon for the preparation of a medicament for treating a disease or disorder with an underlying dysregulation of the emotional functionality, wherein said pipamperon is administered to a patient in a dose ranging between 5 and 15 mg of the active ingredient, and wherein a second compound is administered simultaneously with, separate
5 from or sequential to said pipamperon to augment the therapeutic effect of said second compound.
2. Use according to claim 1, wherein said pipamperon is administered daily at least one
10 day before administering said second compound.
3. Use according to any of claims 1 to 2, wherein said second compound affects the monoaminergic transmitter system.
- 15 4. Use according to claim 3, wherein said second compound is selected from the group comprising: 5-HT reuptake enhancer (1), 5-HT1 autoreceptor agonist (2), 5HT1A receptor agonist (3), 5-HT1A receptor antagonist (4), 5-HT1B receptor antagonist (5), 5-HT2B receptor antagonist (6), 5-HT2C receptor antagonist (7), 5-HT3 receptor antagonist (8), 5-HT6 receptor antagonist (9), adrenergic transmitter releaser (12), α 1 adrenoreceptor antagonist (13), α 2 adrenoreceptor antagonist (14), β 3 adrenoreceptor agonist (19),
20 cannaboid receptor antagonist (21), D1 receptor agonist (27), D2 receptor antagonist (28), D3 receptor antagonist (29), DA uptake inhibitor (30), dopamine receptor agonist (31), H3 receptor antagonist (42), compounds which increase brain concentrations of 5-HT (44), levodopa (48), MAO reuptake inhibitor (50), MAO-A & MAO-B reuptake inhibitor (51), MAO-B inhibitor (52), MAO-B re-uptake inhibitor (53), NARI (60), NaSSA (61), NDRI (62), RIMA (82), SDA (84), SDRI (85), Second messenger beta agonist (86), SNDRI (90), SNRI (91) and SSRI (92).
- 25 5. Use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases or disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders,
30 substance related disorder, personality disorders, antisocial behaviour, bereavement,
35

206

occupational problem and problems related to abuse or neglect and pain disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a 5-HT (serotonin) reuptake enhancer compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said 5-HT (serotonin) reuptake enhancer compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

6. Use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases or disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, substance related disorder, personality disorders, antisocial behaviour, bereavement, occupational problem, problems related to abuse or neglect and pain disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a 5-HT₁ autoreceptor agonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said 5-HT₁ autoreceptor agonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

7. Use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender disorders, sleep disorders, adjustment disorders, impulse control disorders, attention-deficit disorders, substance-related disorder, personality disorders, antisocial behaviour, bereavement, occupational problem, problems related to abuse or neglect, pain disorders, delirium, Alzheimer Disease, substance-induced persisting dementia, vascular dementia, dementia due to HIV disease, dementia due to head trauma, dementia due to Parkinson Disease, dementia due to Huntington Disease, dementia due to Pick Disease, dementia due to Creutzfeldt-Jacob Disease, amnesic disorders due to a general medical condition, substance-induced persisting amnesic disorder, mild cognitive impairment

disorder, other cognitive disorders and Parkinson Disease, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a 5-HT_{1A} (serotonin 1A) receptor agonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said 5-HT_{1A} (serotonin) 1A receptor agonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

8. Use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender disorders, adjustment disorders, impulse control disorders, attention-deficit disorders, substance-related disorder, personality disorders, antisocial behaviour, bereavement, occupational problem and problems related to abuse or neglect, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a 5-HT_{1A} receptor antagonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said 5-HT_{1A} receptor antagonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

9. Use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender disorders, sleep disorders, adjustment disorders, impulse control disorders, attention-deficit disorders, substance-related disorder, personality disorders, antisocial behaviour, bereavement, occupational problem and problems related to abuse or neglect, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a 5-HT_{1B} receptor antagonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said 5-HT_{1B} receptor antagonist compound, further characterized

in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

10. Use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation
5 of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental
disease or disorder selected from the group of diseases and disorders consisting of mood
disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform
disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and
10 gender disorders, sleep disorders, adjustment disorders, impulse control disorders,
substance-related disorder, personality disorders, antisocial behaviour, bereavement,
occupational problem, problems related to abuse or neglect and pain disorders
characterized in that pipamperon or said pharmaceutically acceptable salt thereof is
administered simultaneously with, separate from or prior to the administration of a 5-HT2B
15 receptor antagonist compound to augment the therapeutic effect or to provide a faster onset
of the therapeutic effect of said 5-HT2B receptor antagonist compound, further characterized
in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and
15 mg of the active ingredient.

11. Use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation
20 of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental
disease or disorder selected from the group of diseases and disorders consisting of mood
disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform
disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and
gender disorders, sleep disorders, adjustment disorders, impulse control disorders,
25 substance-related disorder, personality disorders, antisocial behaviour, bereavement,
occupational problem, problems related to abuse or neglect and pain disorders,
characterized in that pipamperon or said pharmaceutically acceptable salt thereof is
administered simultaneously with, separate from or prior to the administration of a 5-HT2C
receptor antagonist compound to augment the therapeutic effect or to provide a faster onset
30 of the therapeutic effect of said 5-HT2C receptor antagonist compound, further characterized
in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and
15 mg of the active ingredient.

12. Use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation
35 of a medicament for treating the underlying emotion dysregulation of substance-related
disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof

is administered simultaneously with, separate from or prior to the administration of a 5-HT₃ receptor antagonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said 5-HT₃ receptor antagonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

13. Use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a cognitive disorder selected from the group of diseases and disorders consisting of Alzheimer Disease, substance-related persisting dementia, vascular dementia, dementia due to HIV disease, dementia due to head trauma, dementia due to Parkinson Disease, dementia due to Huntington Disease, dementia due to Pick Disease, dementia due to Creutzfeldt-Jacob Disease, amnesic disorders due to a general medical condition, substance-induced persisting amnesic disorder, mild cognitive impairment disorder and other cognitive disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a 5-HT₆ receptor antagonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said 5-HT₆ receptor antagonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

14. Use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, adjustment disorders, impulse control disorders, personality disorders, bereavement, occupational problem, problems related to abuse or neglect and pain disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of an adrenergic transmitter releaser compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said adrenergic transmitter releaser compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

15. Use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, personality disorders, bereavement, occupational problem, problems related to abuse or neglect and pain disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a alpha 1 adrenoreceptor antagonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said alpha 1 adrenoreceptor antagonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.
16. Use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting of mood disorders, anxiety disorders, psychotic disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, substance related disorders, personality disorders, bereavement, occupational problem, problems related to abuse or neglect and pain disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a alpha 2 adrenoreceptor antagonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said alpha 2 adrenoreceptor antagonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.
17. Use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, substance related disorders, personality disorders, bereavement, occupational problem,

problems related to abuse or neglect and pain disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a beta 3 adrenoreceptor agonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said beta 3
5 adrenoreceptor agonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

18. Use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation
10 of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting mood disorders, anxiety disorders, psychotic disorders, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sleep disorders, adjustment disorders, impulse control disorders, pervasive development disorders, disruptive behaviour disorders,
15 substance-related disorders, personality disorders, psychological factors affecting medical conditions, malingering, antisocial behaviour, bereavement, occupational problem, identity problem and problems related to abuse or neglect, pain disorders and delirium, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a
20 cannabiod receptor 1 antagonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said cannabiod receptor 1 antagonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

25 19. Use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of substance related disorders and Parkinson disease, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a D1 receptor receptor agonist compound to augment the therapeutic effect
30 or to provide a faster onset of the therapeutic effect of said D1 receptor receptor agonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

20. Use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation
35 of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting mood

212

disorders, psychotic disorders, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sleep disorders, adjustment disorders, impulse control disorders, pervasive development disorders, disruptive behaviour disorders, substance-related disorders, personality disorders, psychological factors affecting medical conditions, malingering, antisocial behaviour, bereavement, occupational problem, identity problem and problems related to abuse or neglect, pain disorders and delirium, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a D2 receptor antagonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said D2 receptor antagonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

21. Use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting of psychotic disorders, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sleep disorders, adjustment disorders, impulse control disorders, pervasive development disorders, disruptive behaviour disorders, substance-related disorders, personality disorders, psychological factors affecting medical conditions, malingering, antisocial behaviour, bereavement, occupational problem, identity problem, problems related to abuse or neglect, pain disorders and delirium, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a D3 receptor antagonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said D3 receptor antagonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

22. Use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of substance related disorders and Parkinson disease, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a DA (dopamine) uptake inhibitor compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said DA (dopamine) uptake

inhibitor compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

23. Use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, adjustment disorders, impulse control disorders, attention-deficit disorders, substance-related disorders, personality disorders and problems related to abuse or neglect, pain disorder and Parkinson disease, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a dopamine receptor agonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said dopamine receptor agonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

24. Use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a cognitive mental disease or disorder selected from the group of diseases and disorders consisting of Alzheimer Disease, substance-related persisting dementia, vascular dementia, dementia due to HIV disease, dementia due to head trauma, dementia due to Parkinson Disease, dementia due to Huntington Disease, dementia due to Pick Disease, dementia due to Creutzfeldt-Jacob Disease, amnesic disorders due to a general medical condition, substance-induced persisting amnesic disorder, mild cognitive impairment disorder and other cognitive disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a histamine H3receptor antagonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said histamine H3receptor antagonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

25. Use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non cognitive mental disease or disorder selected from the group of diseases and disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform

disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, substance-related disorders, personality disorders, bereavement, occupational problem, problems related to abuse or neglect and pain disorder, characterized in that pipamperon or
5 said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a compound which increases brain concentrations of 5-HT (serotonin) to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said compound which increases brain concentrations of 5-HT (serotonin), further characterized in that pipamperon is to be administered to a patient in a
10 daily dose ranging between 5 and 15 mg of the active ingredient.

26. Use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of Parkinson Disease, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is
15 administered simultaneously with, separate from or prior to the administration of a levodopa/decarboxylase inhibitor compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said levodopa/decarboxylase inhibitor compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

27. A pharmaceutical composition comprising (a) pipamperon, and (b) a levodopa/decarboxylase inhibitor compound, preferably is (eti)levodopa/carbidopa, or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof in combination with entacapone, which is an inhibitor of catechol-O-methyltransferase (COMT),
25 or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of Parkinson Disease.

28. A pharmaceutical composition according to claim 27, wherein pipamperon is
30 provided in a unitary dose of between 5 and 15 mg of the active ingredient and wherein said levodopa/decarboxylase inhibitor compound is levodopa/carbidopa, preferably provided in a unitary dose of between 100 mg and 10 mg of the active ingredient.

29. A pharmaceutical composition according to claim 27 or 28, wherein pipamperon is
35 provided in a unitary dose of between 5 and 15 mg of the active ingredient and wherein said levodopa/decarboxylase inhibitor compound is levodopa/carbidopa or etilevodopa/ carbidopa

in combination with entacapone, of which the latter is preferably provided in a unitary dose of between 500 mg and 100 mg of the active ingredient.

30. Use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation
5 of a medicament for treating the underlying emotion dysregulation of non-cognitive mental
disease or disorder which are substance related disorders and Parkinson disease,
characterized in that pipamperon or said pharmaceutically acceptable salt thereof is
administered simultaneously with, separate from or prior to the administration of a
10 monoamine oxidase (MAO) reuptake inhibitor compound to augment the therapeutic effect
or to provide a faster onset of the therapeutic effect of said monoamine oxidase (MAO)
reuptake inhibitor compound, further characterized in that pipamperon is to be administered
to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

31. Use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation
15 of a medicament for treating the underlying emotion dysregulation of a non cognitive mental
disease or disorder selected from the group of diseases and disorders consisting of mood
disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform
disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and
gender identity disorders, adjustment disorders, impulse control disorders, attention-deficit
20 disorders, personality disorders, antisocial behaviour, bereavement, occupational problem,
problems related to abuse or neglect and pain disorder, characterized in that pipamperon or
said pharmaceutically acceptable salt thereof is administered simultaneously with, separate
from or prior to the administration of a monoamine oxidase A (MAO-A) and a monoamine
oxidase B (MAO-B) reuptake inhibitor compound to augment the therapeutic effect or to
25 provide a faster onset of the therapeutic effect of said monoamine oxidase A (MAO-A) and a
monoamine oxidase B (MAO-B) reuptake inhibitor compound, further characterized in that
pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg
of the active ingredient.

30 32. Use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation
of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental
disease or disorder selected from the group of diseases and disorders consisting of mood
disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform
disorders (excluding pain disorders), factitious disorders, dissociative disorders, adjustment
35 disorders, impulse control disorders, attention-deficit disorders, substance-related disorders,
personality disorders, problems related to abuse or neglect, pain disorder and Parkinson

disease, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a monoamine oxidase B (MAO-B) inhibitor compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said monoamine oxidase B (MAO-B) inhibitor compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

33. Use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of Parkinson Disease, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a monoamine oxidase B (MAO-B) reuptake inhibitor to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said monoamine oxidase B (MAO-B) reuptake inhibitor, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

34. Use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting of mood disorders, anxiety disorders, adjustment disorders, attention-deficit disorders, personality disorders, antisocial behaviour, bereavement, occupational problem, problems related to abuse or neglect and pain disorder, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a selective nor-adrenaline re-uptake inhibitor (NARI) compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said selective nor-adrenaline re-uptake inhibitor (NARI) compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

35. Use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, personality disorders, antisocial behaviour, bereavement, occupational problem, problems

related to abuse or neglect and pain disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a noradrenergic/specific serotonergic antidepressant (NaSSA) compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said noradrenergic/specific serotonergic antidepressant (NaSSA) compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

36. Use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting of mood disorders, anxiety disorders, adjustment disorders, attention-deficit disorders, personality disorders, antisocial behaviour, bereavement, occupational problem, problems related to abuse or neglect and pain disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a selective nor-adrenaline and dopamine re-uptake inhibitor (NDRI) compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said selective nor-adrenaline and dopamine re-uptake inhibitor (NDRI) compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

37. Use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, adjustment disorders, impulse control disorders, personality disorders, antisocial behaviour, bereavement, occupational problem, problems related to abuse or neglect and pain disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a compound which is a reversible inhibitor of mono-amine oxydase A (RIMA) to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said compound which is a reversible inhibitor of mono-amine oxydase A (RIMA), further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

38. Use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting of mood disorders, anxiety disorders, psychotic disorders, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sleep disorders, adjustment disorders, impulse control disorders, pervasive development disorders, disruptive behaviour disorders, substance-related disorders, personality disorders, psychological factors affecting medical conditions, malingering, antisocial behaviour, bereavement, occupational problem, identity problem, problems related to abuse or neglect, pain disorder and delirium, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a serotonin/dopamine antagonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said serotonin/dopamine antagonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

39. Use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sleep disorders, adjustment disorders, impulse control disorders, substance-related disorders, personality disorders, antisocial behaviour, bereavement, occupational problem, problems related to abuse or neglect, pain disorders, delirium, Alzheimer Disease, substance-related persisting dementia, vascular dementia, dementia due to HIV disease, dementia due to head trauma, dementia due to Parkinson Disease, dementia due to Huntington Disease, dementia due to Pick Disease, dementia due to Creutzfeldt-Jacob Disease, amnesic disorders due to a general medical condition, substance-induced persisting amnesic disorder, mild cognitive impairment disorder and other cognitive disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a selective serotonin and dopamine re-uptake inhibitor (SDRI) compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said selective serotonin and dopamine re-uptake inhibitor (SDRI) compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

40. Use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, substance-related disorders, personality disorders, bereavement, occupational problem, problems related to abuse or neglect and pain disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a second messenger beta agonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said second messenger beta agonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

41. Use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sleep disorders, adjustment disorders, impulse control disorders, attention-deficit disorders, substance-related disorders, personality disorders, antisocial behaviour, bereavement, occupational problem, problems related to abuse or neglect and pain disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a selective serotonin, nor-adrenaline and dopamine re-uptake inhibitor (SNDRI) compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said selective serotonin, nor-adrenaline and dopamine re-uptake inhibitor (SNDRI) compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

42. Use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sleep

disorders, adjustment disorders, impulse control disorders, attention-deficit disorders, substance-related disorders, personality disorders, antisocial behaviour, bereavement, occupational problem, problems related to abuse or neglect and pain disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a selective serotonin and nor-adrenaline re-uptake inhibitor (SNRI) compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said selective serotonin and nor-adrenaline re-uptake inhibitor (SNRI) compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

43. Use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sleep disorders, adjustment disorders, impulse control disorders, substance-related disorders, personality disorders, antisocial behaviour, bereavement, occupational problem, problems related to abuse or neglect and pain disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a selective serotonin re-uptake inhibitor (SSRI) compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said selective serotonin re-uptake inhibitor (SSRI) compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

44. Use according to claim 1 or 2, wherein said disease or disorder is Alzheimer disease.

45. Use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of Alzheimer Disease, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a 5-HT_{1A} (serotonin 1A receptor) agonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said 5-HT_{1A} (serotonin 1A receptor) agonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

46. Use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of Alzheimer Disease, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a 5-HT6 antagonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said 5-HT6 antagonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

47. Use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of Alzheimer Disease, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of an acetylcholinesterase inhibitor compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said acetylcholinesterase inhibitor compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

48. Use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of Alzheimer Disease, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of an AMPA receptor mediator compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said AMPA receptor mediator compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

49. Use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of Alzheimer Disease, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of an amyloid aggregation-inhibitor compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said amyloid aggregation-inhibitor compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

50. Use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of Alzheimer Disease, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a calcium channel modulator compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said calcium channel modulator compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

51. Use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of Alzheimer Disease, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a choline uptake enhancer compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said choline uptake enhancer compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

52. Use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of Alzheimer Disease, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a compound that activates ERK (extracellular signal-related kinase) to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said compound that activates ERK (extracellular signal-related kinase), further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

53. Use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of Alzheimer Disease, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a GABA agonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said GABA agonist compound, further characterized in that pipamperon

is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

54. Use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of Alzheimer Disease, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a histamine H3-receptor antagonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said histamine H3-receptor antagonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

55. Use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of Alzheimer Disease, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a compound which increases insulin sensitivity to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said compound which increases insulin sensitivity, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

56. Use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of Alzheimer Disease, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a compound which mimics the effect of nerve growth factor (NGF) to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said compound which mimics the effect of nerve growth factor (NGF), further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

57. Use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of Alzheimer Disease, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a muscarinic receptor partial agonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said muscarinic receptor partial agonist compound,

further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

58. Use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of Alzheimer Disease, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a selective nor-adrenaline and dopamine re-uptake inhibitor (NDRI) compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said selective nor-adrenaline and dopamine re-uptake inhibitor (NDRI) compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

59. Use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of Alzheimer Disease, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from nerve growth factor (NGF) gene therapy, to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said nerve growth factor (NGF) gene therapy, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

60. Use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of Alzheimer Disease, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a nicotinic receptor agonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said nicotinic receptor agonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

61. Use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of Alzheimer Disease, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of an N-Methyl-D-aspartate (NMDA) antagonist compound to augment the therapeutic effect or to

provide a faster onset of the therapeutic effect of said N-Methyl-D-aspartate (NMDA) antagonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

5 62. Use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation
of a medicament for treating the underlying emotion dysregulation of Alzheimer Disease,
characterized in that pipamperon or said pharmaceutically acceptable salt thereof is
administered simultaneously with, separate from or prior to the administration of a non-
steroidal anti-inflammatory drug to augment the therapeutic effect or to provide a faster
10 onset of the therapeutic effect of said a non-steroidal anti-inflammatory drug, further
characterized in that pipamperon is to be administered to a patient in a daily dose ranging
between 5 and 15 mg of the active ingredient.

15 63. Use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation
of a medicament for treating the underlying emotion dysregulation of Alzheimer Disease,
characterized in that pipamperon or said pharmaceutically acceptable salt thereof is
administered simultaneously with, separate from or prior to the administration of a
phosphodiesterase-4 (PDE4) inhibitor compound to augment the therapeutic effect or to
provide a faster onset of the therapeutic effect of said phosphodiesterase-4 (PDE4) inhibitor
20 compound, further characterized in that pipamperon is to be administered to a patient in a
daily dose ranging between 5 and 15 mg of the active ingredient.

25 64. Use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation
of a medicament for treating the underlying emotion dysregulation of Alzheimer Disease,
characterized in that pipamperon or said pharmaceutically acceptable salt thereof is
administered simultaneously with, separate from or prior to the administration of a peptidic
compound to augment the therapeutic effect or to provide a faster onset of the therapeutic
effect of said peptidic compound, further characterized in that pipamperon is to be
administered to a patient in a daily dose ranging between 5 and 15 mg of the active
30 ingredient.

35 65. Use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation
of a medicament for treating the underlying emotion dysregulation of Alzheimer Disease,
characterized in that pipamperon or said pharmaceutically acceptable salt thereof is
administered simultaneously with, separate from or prior to the administration of a compound
which protects dopaminergic and cholinergic neurons to augment the therapeutic effect or to

provide a faster onset of the therapeutic effect of said compound which protects dopaminergic and cholinergic neurons, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

5

66. Use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of Alzheimer Disease, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a selective serotonin and dopamine reuptake inhibitor (SDRI) compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said selective serotonin and dopamine reuptake inhibitor (SDRI) compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

10
15

67. Use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of Alzheimer Disease, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a selective serotonin and dopamine reuptake inhibitor (SDRI) compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said selective serotonin and dopamine reuptake inhibitor (SDRI) compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

20
25

68. Use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of Alzheimer Disease, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a selective serotonin, nor-adrenaline and dopamine re-uptake inhibitor (SNDRI) compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said selective serotonin, nor-adrenaline and dopamine re-uptake inhibitor (SNDRI) compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

30
35

69. Use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, substance related disorders, personality disorders, bereavement, occupational problem and problems related to abuse or neglect, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a CRF1 antagonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said CRF1 antagonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

70. Use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, substance-related disorders, personality disorders, bereavement, occupational problem and problems related to abuse or neglect, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a GR antagonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said GR antagonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

71. Use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, substance-related disorders, personality disorders, bereavement, occupational problem and problems related to abuse or neglect, characterized in that pipamperon or said

pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a melatonin receptor (MT) agonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said melatonin receptor (MT) agonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

72. Use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting of mood disorders, anxiety disorders, psychotic disorders, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sleep disorders, adjustment disorders, impulse control disorders, pervasive development disorders, disruptive behaviour disorders, substance-related disorders, personality disorders, psychological factors affecting medical conditions, malingering, antisocial behaviour, bereavement, occupational problem, identity problem and problems related to abuse or neglect, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a neurotensin receptor antagonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said neurotensin receptor antagonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

73. Use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, substance-related disorders, personality disorders, bereavement, occupational problem and problems related to abuse or neglect, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a neurokinin 2 receptor (NK2) antagonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said neurokinin 2 receptor (NK2) antagonist compound, further characterized in that pipamperon

is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

74. Use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting of mood disorders, anxiety disorders, psychotic disorders, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sleep disorders, adjustment disorders, impulse control disorders, pervasive development disorders, disruptive behaviour disorders, substance-related disorders, personality disorders, psychological factors affecting medical conditions, malingering, antisocial behaviour, bereavement, occupational problem, identity problem and problems related to abuse or neglect, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a neurokinin 3 receptor (NK3) antagonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said neurokinin 3 receptor (NK3) antagonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

75. Use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, adjustment disorders, impulse control disorders, personality disorders, bereavement, occupational problem and problems related to abuse or neglect, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a peptidic compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said peptidic compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

76. Use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting of mood

disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, substance-related disorders, personality disorders, bereavement, occupational problem and
5 problems related to abuse or neglect, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a substance P receptor (NK1) antagonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said substance P receptor (NK1) antagonist compound, further characterized in that pipamperon
10 is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

77. Use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental
15 disease or disorder selected from the group of diseases and disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, substance-related disorders, personality disorders, bereavement, occupational problem and
20 problems related to abuse or neglect, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a tachykinin antagonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said tachykinin antagonist compound, further characterized in that pipamperon is to be administered to a
25 patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

78. Use according to any of claims 1 to 4 or 7, wherein said second compound is gepirone.

79. A method for preparing a compound having a selective D4 and 5-HT2A antagonist, reverse agonist or partial agonist activity comprising the following steps: (a) measuring the selective affinity of a test compound to the D4 receptor and selecting a compound that has a pKi value equal to or greater than 8 towards the D4 receptor in respect to all the other D receptors, and measuring the selective efficacy of the selected compound to the D4 receptor
30 and selecting a compounds which is a selective antagonist, inverse agonist or partial agonist of the D4 receptor; (b) measuring the selective affinity of a test compound to the 5-HT2A
35

231

receptor and selecting a compound that has a pKi value equal to or greater than 8 towards the 5-HT_{2A} receptor in respect to all the other 5HT receptors, and measuring the selective efficacy of the selected compound to the 5-HT_{2A} receptor and selecting a compounds which is a selective antagonist, inverse agonist or partial agonist of the 5-HT_{2A} receptor; (c) identifying a compound which is selected in (a) and (b), (d) preparing the compound identified in (c).


80. Compound prepared by the method of claim 79.

81. Use according to any of claims 1, 2 or 7, wherein said second compound is chosen from the group consisting of fluvoxamine controlled release, phenserine tartrate, atomoxetine hydrochloride, bupropion (controlled-release formulation), ropinirole HCL (controlled-release formulation), INN 00835, galantamine (extended release formulation), paliperidone, tomoxetine, aprepitant, rivastigmine tartrate, ORG 34517/34850, sunepitron, sumanirole, milnacipran, idazoxan, xaliproden, SR 58611, befloxatone, litoxetine, tianeptine, agomelatine, SPD 503, flesinoxan, bifeprunox, ramelteon, etilevodopa, rasagiline (TVP-1012) and desvenlafaxine.

82. Use according to any of claims 1, 2 or 7, wherein said second compound is chosen from the group consisting of galantamine (extended release formulation), R121919, risperidone, paliperidone and R228060 (YKP-10A).

CGI-SEVERITY SCORE: Foregoing & Add-On Treatment with Pipamperon 8 mg/day (bid) and Citalopram 20-40 mg/day (bid) in comparison with the

SSRI in Panic Disorder




—○— Paroxetine in the treatment of panic disorder, Journal of Clinical Psychiatry 65: 405-413, No. 3, Mar 2004

-*- COMBO PIPAMPERON-CITALOPRAM (n=3)

Setraline in the treatment of panic disorder, Journal of Clinical Psychiatry 65: 405-413, No. 3, Mar 2004


**Y-BOCS COMPULSION SCORE: Foregoing & Add-On Treatment
with Pipamperon 8-16 mg/day (bid) and Citalopram 30-80 mg/day
(bid) in comparison with the SSRI fluvoxamine in OCD**



—●— COMBO PIPAMPERON/CITALOPRAM (n=7)

FLUVOXAMINE Controlled Release mean
271 mg/day (n=253) (Hollander E, Journal of
Clinical Psychiatry 64: 640-647, Jun 2003)


**Y-BOCS OBSESSION SCORE: Foregoing & Add-On Treatment
with Pipamperon 8-16 mg/day (bid) and Citalopram 30-80 mg/day
(bid) in comparison with the SSRI fluvoxamine in OCD**



○ COMBO PIPAMPERON-CITALOPRAM (n=7)

FLUVOXAMINE Controlled Release mean
271 mg/day (n=253) (Hollander E, Journal of
Clinical Psychiatry 64: 640-647, Jun 2003)

**Y-BOCS TOTAL SCORE: Foregoing & Add-On Treatment with
Pipamperon 8-16 mg/day (bid) and Citalopram 30-80 mg/day (bid)
in comparison with the SSRI fluvoxamine in OCD**




COMBO PIPAMPERON-CITALOPRAM (n=7)

FLUVOXAMINE Controlled Release mean
271 mg/day (n=253) (Hollander E, Journal of
Clinical Psychiatry 64: 640-647, Jun 2003)


REMISSION RATES (HDRS-17 ≤ 7): Foregoing & Add-On

Treatment with Pipamperon 8-12 mg/day (bid) and Citalopram 20-40 mg/day (bid) in comparison with the


SNRI venlafaxine in MAJOR DEPRESSION



HDRS-17 CHANGE FROM BASELINE: Foregoing & Add-On Treatment with Pipamperon 8-12 mg/day (bid) and Citalopram 20-40 mg/day (bid) in comparison with the SNRI duloxetine in MAJOR DEPRESSION




Foregoing & Add-On Treatment with Pipamperon 8-12 mg/day (bid) and Citalopram 20-40 mg/day (bid) in MDD in Comparison with the Standard Efficacy of Antidepressants in Clinical Trials*



* A. KHAN et al, Symptom Reduction and Suicide Risk in Patients Treated With Placebo in Antidepressant Clinical Trials, ARCH. OF GENERAL PSYCHIATRY / VOL 57, APR 2000)


**HDRS-17 CHANGE FROM BASELINE: COMBO PIPAMPERON-
CITALOPRAM WITH A FORE-GOING TREATMENT OF 7 DAYS
WITH PIPAMPERON 11mg/day vs SNRI (duloxetine) in MAJOR
DEPRESSION**



COMBO PIPAMPERON-CITALOPRAM WITH PIPAMPERON 8-12 (mean 11) mg/day (bid) 6-8 (mean 7) days before treatment with the SSRI (n=3)


DULOXETINE 40-120 mg/day (n=152) (Goldstein et al, J. Clin. Psychiatry (in press))

Foregoing Treatment During 6-8 (mean 7) days with Pipamperon 8-12 (mean 11) mg/day (bid) Followed With the Combination Treatment of Pipamperon and Citalopram 20-40 (mean 30) mg/day (bid) (PICIT FG 6-8) in MDD (HDRS-17 at BL = 28 in Comparison with the Standard Efficacy of Antidepressants in Clinical Trials*



* A. KHAN et al, Symptom Reduction and Suicide Risk in Patients Treated With Placebo in Antidepressant Clinical Trials, ARCH. OF GENERAL PSYCHIATRY / VOL 57, APR 2000)

**REMISSION RATES (HDRS-17 ≤ 7): COMBO PIPAMPERON-CITALOPRAM
WITH A FORE-GOING TREATMENT OF 4 DAYS WITH PIPAMPERON 10mg/day
vs SNRI (venlafaxine) in MAJOR DEPRESSION**




COMBO PIPAMPERON-CITALOPRAM WITH PIPAMPERON 8-12 (mean 10) mg/day 1-5 (mean 4) days before treatment with the SSRI (n=15) SNRI Venlafaxine - Meta-analysis of Thase et al, Br J Psychiatry 2001; 178: 234-241


SSRI's - Meta-analysis of Thase et al, Br J Psychiatry 2001; 178: 234-241

PLACEBO - Meta-analysis of Thase et al, Br J Psychiatry 2001; 178: 234-241

**HDRS-17 CHANGE FROM BASELINE: COMBO PIPAMPERON-
CITALOPRAM WITH A FORE-GOING TREATMENT OF 4 DAYS
WITH PIPAMPERON 10mg/day vs SNRI (duloxetine) in MAJOR
DEPRESSION**




Foregoing Treatment During 1-5 (mean 4) days with Pipamperon 8-12 (mean 10) mg/day (bid) Followed With the Combination Treatment of Pipamperon and Citalopram 20-50 (mean 26) mg/day (bid) (PICIT FG 1-5) in MDD (HDRS-17 at BL = 23) in Comparison with the Standard Efficacy of Antidepressants in Clinical Trials*



* A. KHAN et al, Symptom Reduction and Suicide Risk in Patients Treated With Placebo in Antidepressant Clinical Trials, ARCH. OF GENERAL PSYCHIATRY / VOL 57, APR 2000)

REMISSION RATES (HDRS-17 ≤ 7): COMBO PIPAMPERON AS ADD- ON - CITALOPRAM vs SNRI (venlafaxine) vs SSRI's vs PLACEBO in MAJOR DEPRESSION




COMBO PIPAMPERON AS ADD-ON - CITALOPRAM WITH PIPAMPERON 8-12mg/day (mean 9mg/day) 20-60 (mean 33) days after the treatment with the SSRI (n=5)
SNRI Venlafaxine - Meta-analysis of Thase et al, Br J Psychiatry 2001; 178: 234-241

SSRI's - Meta-analysis of Thase et al, Br J Psychiatry 2001; 178: 234-241

PLACEBO - Meta-analysis of Thase et al, Br J Psychiatry 2001; 178: 234-241


HDRS-17 CHANGE FROM BASELINE: COMBO PIPAMPERON AS ADD-ON - CITALOPRAM vs SNRI (duloxetine) in MAJOR DEPRESSION



■ COMBO PIPAMPERON AS ADD-ON -
 CITALOPRAM WITH PIPAMPERON 8-
 12mg/day (mean 9mg/day) 20-60
 (mean 33) days after the treatment with
 the SSRI (n=5)

DULOXETINE 40-120 mg/day (n=152)
 (Goldstein et al, J. Clin. Psychiatry (in
 press))

Add-On Treatment with *Pipamperon* 8-12 (mean 9) mg (*bid*) after Treatment with *Citalopram* 10-20 (mean 30) mg (*bid*) during 20-60 (mean 33) days (*PIPCIT ADD-ON*) with *HDRS-17* Totalscore = 29 at Baseline in *MDD* in Comparison with the Standard Efficacy of Antidepressants in Clinical Trials*



* A. KHAN et al, Symptom Reduction and Suicide Risk in Patients Treated With Placebo in Antidepressant Clinical Trials, ARCH. OF GENERAL PSYCHIATRY / VOL 57, APR 2000)



